FROM THE FUND BEQUEATHED BY
ROBERT CHARLES BILLINGS 1819 1899
TO THE
BOSTON MEDICAL LIBRARY
Nº 8 THE FENWAY

DISEASES OF THE EYE

ANREP AND HARRIS' PRACTICAL PHYSIOLOGY.

By G. V. ANREP, M.D., D.Sc., Senior Assistant in Physiology, and D. T. HARRIS, M.B., B.S., Assistant Professor of Physiology, University College, London. 197 Illustrations. 10*s.* 6*d.*

STARLING'S PRINCIPLES OF HUMAN PHYSIOLOGY.

By E. H. STARLING, C.M.G., M.D., F.R.C.P., F.R.S., Jodrell Professor of Physiology in University College, London. Third Edition. 579 Figures. 25*s.*

CLARK'S APPLIED PHARMACOLOGY.

By A. J. CLARK, M.C., M.D., F.R.C.P., Professor of Pharmacology in the University of London; Examiner in the Universities of London, Edinburgh, and Wales. 46 Illustrations. 15*s.*

MOORE'S MEDICAL OPHTHALMOLOGY.

By R. FOSTER MOORE, O.B.E., B.Ch., F.R.C.S., Assistant Ophthalmic Surgeon, St. Bartholomew's Hospital; Surgeon Moorfields Eye Hospital. 80 Illustrations. 15*s.*

TAYLOR'S PRACTICE OF MEDICINE.

Twelfth Edition. *By* E. P. POULTON, M.D., F.R.C.P., with the assistance of C. PUTNAM SYMONDS, M.D., M.R.C.P., and H. W. BARBER, M.B., F.R.C.P., Physicians, Guy's Hospital. 30*s.*

COLLIS AND GREENWOOD'S HEALTH OF THE INDUSTRIAL WORKER.

By E. L. COLLIS, B.Ch., M.D., Mansel Talbot Professor of Preventive Medicine, Welsh National School of Medicine; Member of Industrial Fatigue Research Board; and MAJOR GREENWOOD, M.R.C.P., M.R.C.S., Member of Industrial Fatigue Research Board; Reader in Medical Statistics, University of London. 30*s.*

J. & A. CHURCHILL

DISEASES OF THE EYE

BY

SIR JOHN HERBERT PARSONS,

C.B.E., D.Sc., F.R.C.S., F.R.S.

OPHTHALMIC SURGEON, UNIVERSITY COLLEGE HOSPITAL; SURGEON, ROYAL LONDON (MOORFIELDS) OPHTHALMIC HOSPITAL; LATE OPHTHALMIC SURGEON, HOSPITAL FOR SICK CHILDREN, GREAT ORMOND STREET.

FOURTH EDITION

With 21 Plates and 326 Text Figures

LONDON

J. & A. CHURCHILL

7 GREAT MARLBOROUGH STREET

1923

28812 Bi

First Edition 1907
Second Edition 1912
Third Edition 1918
Fourth Edition 1923

28, D. 366



Printed in Great Britain.

PREFACE TO THE FOURTH EDITION

The text in this edition has been revised throughout. Considerable additions have been made, including three new coloured plates and several figures. By a freer use of small print the relative importance of the various subjects to the student has been thrown into relief and increase in the size of the book obviated. A short section on Preventive Ophthalmology has been added. It is hoped that the *résumé* of the Causes and Prevention of Blindness will emphasise the need of special attention to these matters, not only by the student, but also by his teachers and the examining bodies. The remarks on the Hygiene of Vision will indicate the widespread ramifications of applied ophthalmology.

I am indebted to Mr. E. E. Henderson for kindly reading the proofs.

J. HERBERT PARSONS.

Queen Anne Street,
W. 1.

Digitized by the Internet Archive
in 2026

https://archive.org/details/diseasesofeye0000sirj

CONTENTS

SECTION I.

ANATOMY AND PHYSIOLOGY.

CHAP.		PAGE
I.	Anatomy	1
II.	Physiology	14
III.	Elementary Optics	19
IV.	Elementary Physiological Optics	32
V.	The Neurology of Vision	65

SECTION II.

THE EXAMINATION OF THE EYE.

VI.	External Examination	74
VII.	Ophthalmoscopic Examination	93
VIII.	The Fundus Oculi	117
IX.	Functional Examination	124

SECTION III.

DISEASES OF THE EYE.

X.	Diseases of the Conjunctiva	139
XI.	Diseases of the Cornea	187
XII.	Diseases of the Sclerotic	236
XIII.	Diseases of the Iris and Ciliary Body	241
XIV.	Glaucoma	263
XV.	The Lens	285
XVI.	Diseases of the Vitreous	306
XVII.	Diseases of the Choroid and Retina	312
XVIII.	Diseases of the Optic Nerve	357
XIX.	Symptomatic Disturbances of Vision	370
XX.	Intra-Ocular Tumours	385
XXI.	Injuries to the Eye, Panophthalmitis, and Sympathetic Ophthalmia	393
XXII.	Operations upon the Eyeball	431

SECTION IV.

ERRORS OF REFRACTION AND ANOMALIES OF ACCOMMODATION

CHAP.

XXIII. Retinoscopy . . . . . .
XXIV. Errors of Refraction . . . . .
XXV. Anomalies of Accommodation . . . .

SECTION V.

DISORDERS OF MOTILITY OF THE EYE.

XXVI. Anatomy and Physiology of the Extrinsic Ocular Muscles . . . . . .
XXVII. Paralytic and Kinetic Strabismus. Nystagmus . . . . . .
XXVIII. Concomitant Strabismus. Heterophoria. Congenital Defects . . . . .

SECTION VI.

SYMPTOMATIC DISEASES OF THE EYE.

XXIX. Ocular Manifestations of Diseases of the Nervous System . . . . . .
XXX. Ocular Manifestations of Other Diseases .

SECTION VII.

DISEASES OF THE ADNEXA OF THE EYE.

XXXI. Diseases of the Lids 5
XXXII. Diseases of the Lacrymal Apparatus . 7
XXXIII. Diseases of the Orbit 6

SECTION VIII.

PREVENTIVE OPHTHALMOLOGY.

XXXIV. The Causes and Prevention of Blindness 6
XXXV. The Hygiene of Vision 6

APPENDICES.

APPENDIX

I. Preliminary Investigation of the Patient 6
II. Therapeutic Notes 6
III. Requirements of Candidates for Admission into the Public Services . . . 6

INDEX 6

BOSTON MEDICAL
JUN 28 1923
LIBRARY

DISEASES OF THE EYE.

SECTION I.

ANATOMY AND PHYSIOLOGY.

CHAPTER I.

ANATOMY.

The sensory nerves of the body are provided with end organs, by means of which they receive specific physical stimuli and transform them into nerve impulses. The nerves of the special senses are no exceptions to the rule, and the eye is the highly differentiated and complex end organ of the sense of sight.

The wall of the globe is composed of a dense, elastic supporting membrane (Fig. 1). The anterior part of the membrane is transparent—the cornea; the remainder is opaque—the sclerotic. The anterior part of the sclerotic is covered by mucous membrane the conjunctiva which is reflected from its surface on to the lids.

The *cornea* consists of three layers: the epithelium, the substantia propria, and Descemet's membrane. The epithelium, which is stratified, may be regarded as the continuation of the conjunctiva over the cornea proper. It lies upon a homogeneous lamina of the substantia propria, called Bowman's membrane. The substantia propria may be regarded as the continuation forwards of the sclerotic. Descemet's membrane is a thin elastic membrane, covered on its posterior surface by endothelium: it may be regarded as the continua-

tion forwards of the uveal tract. We shall see that the relationship of the three layers is of some pathological importance, for when as is often the case the cornea suffers secondarily to some conjunctival complaint the epithelium and superficial layers are most likely to be affected: similarly, in diseases of the sclerotic the substantia propria suffers most, and in diseases of the uveal tract the endothelium, Descemet's



FIG. 1.—Diagrammatic horizontal section of the eye. C, cornea; *a.c.*, anterior chamber; I, iris; C.B., ciliary body; *p.c.*, posterior chamber; L, lens; V, vitreous; R, retina; Ch., choroid; Scl., sclerotic; *f*, fovea centralis; O.N., optic nerve.

membrane, and the adjoining posterior layers of the substantia propria.

The cornea is set into the sclerotic like a watch glass, *i.e.*, the sclerotic overlaps the cornea all round the periphery. The cornea is very richly supplied with nerve fibres derived from the trigeminal. It has no blood vessels with the exception of minute festoons, about 1 mm. broad, at the periphery; the cornea is therefore dependent for its nourishment upon diffusion of lymph, which is supplied from the conjunctival vessels.

Lining the sclerotic are two membranes: an outer, highly

vascular, concerned chiefly in the nutrition of the eye, and comprising the greater part of the uveal tract; and an inner, nervous, the true visual nerve ending, concerned in the reception and transformation of light stimuli, and called the retina.

The uveal tract consists of three parts, of which the two posterior, the choroid and ciliary body, line the sclerotic,

FIG. 2.—Angle of the anterior chamber. A, cornea; B, canal of Schlemm, which appears as several small spaces in the sclerotic just outside the ligamentum pectinatum iridis; it thus consists of irregular anastomosing venous channels which are cut across in the section; C, ciliary muscle; D, circulus arteriosus iridis major; E, iris; F, ciliary processes.

while the anterior forms a free circular diaphragm, the iris. The plane of the iris is approximately coronal: the aperture of the diaphragm is the pupil. Situated behind the iris and in contact with the pupillary margin is the crystalline lens.

The *anterior chamber* is a space filled with lymph, the *aqueous humour;* it is bounded in front by the cornea, behind by the iris and the part of the anterior surface of the lens which is exposed in the pupil. The sclerotic enters into the boundaries of the anterior chamber at the part which is known as the angle of the anterior chamber (Fig. 2). In the

inner layers of the sclerotic at this part there is a network of venous spaces which is called the *canal of Schlemm*. At the periphery, just anterior to the canal of Schlemm, Descemet's membrane splits up into fibrillæ, which are continuous with a meshwork of fibres stretching between the sclerotic and the iris, and known as the *ligamentum pectinatum iridis*. These fibres are covered by endothelium, which is continuous with that lining the cornea and also with that covering the iris. The spaces in the network of the ligamentum pectinatum iridis are called the *spaces of Fontana*: they are much better developed in lower mammals than in man. The tissue separating the ligamentum pectinatum from the canal of Schlemm is somewhat denser, and there is no free communication between the anterior chamber and the venous plexus, a thin membrane, covered on each surface by endothelium, being interposed. We shall see that a thorough knowledge of the anatomy of angle of the anterior chamber is essential to the proper understanding of several pathological problems, especially that of glaucoma.

The anterior chamber is about 2·5 mm. deep in the centre in the normal adult: it is shallower in very young children and also in old people.

The *uveal tract*, as already mentioned, consists of the iris, the ciliary body, and the choroid, from before backwards.

The *iris* is composed of a stroma, consisting of branched connective tissue cells, usually pigmented, but unpigmented in blue irides, and containing a rich supply of blood vessels which run in a radial direction. The stroma is covered on its posterior surface by two layers of pigmented epithelium, which properly belong to the retina and are therefore called the pars iridica retinæ, or pars retinalis iridis. The anterior layer consists of flattened cells, which are very firmly attached to the stroma, the posterior of cubical cells, not so firmly attached to the anterior layer. Near the pupillary margin and concentric with it is a bundle of unstriped muscle fibres, the *sphincter iridis*. There is also a set of cells, elongated in the radial direction, which acts as a *dilatator iridis*.

The anterior surface of the iris is covered with a single layer of endothelium, except at some minute depressions or crypts which are found most at the ciliary border. Here the lymph spaces between the stroma cells communicate directly with the anterior chamber: this is probably a device for ensuring rapid transference of lymph from the iris to the anterior chamber and *vice versâ*, so as to facilitate quick movements of

the pupil in response to variations in the intensity of the light falling upon the eye. The iris is thinnest at its attachment to the ciliary body, so that if torn it tends to give way here.

The iris is richly supplied by sensory nerve fibres derived from the trigeminal, a fact which it is important to remember, since touching or cutting the iris, especially if it is inflamed, is intensely painful. The sphincter iridis is supplied by motor nerve fibres derived from the oculomotor nerve, whilst the motor fibres of the dilatator iridis are derived from the cervical sympathetic nerve.



FIG. 3.—Diagrammatic meridional section of the ciliary body. Note that the sclerotic overlaps the cornea, as shown by the shading.

The *ciliary body* in antero-posterior section is shaped roughly like an isosceles triangle, with the base forwards. The iris is attached to about the middle of the base, so that a small portion of the ciliary body enters into the posterior boundary of the anterior chamber at the angle (Fig. 3). The chief mass of the ciliary body is composed of unstriped muscle fibres, the *ciliary muscle*. This consists of two parts, an outer, in contact with the sclerotic, consisting of antero-posterior or meridional fibres, and an inner, consisting of fibres running at right angles to the former, and therefore arranged in a circle concentric with the base of the iris. The meridional fibres can be traced far back, well into the choroid, whereas the circular fibres are confined to the anterior part.

If an eye is cut in half in an antero-posterior direction and the inner surface of the ciliary body is inspected, it will be noticed that the anterior part has a number of folds upon it,

while the posterior part is smooth. The anterior part is, therefore, called the pars plicata, the posterior, the pars plana. If the plications are counted with the naked eye or under slight magnification, it will be found that there are about seventy in the whole circumference. If microscopical sections are examined, innumerable smaller plications and processes, the *ciliary processes*, will be seen upon the pars plicata. These contain no part of the ciliary muscle, but consist essentially of tufts of blood vessels, not unlike the glomeruli of the kidney. They are covered upon the inner surface by two layers of epithelium, which belong properly to the retina, and are hence called the pars ciliaris retinæ. As in the pars iridica retinæ, the outer layer, corresponding with the anterior in the iris, consists of flattened cells, the inner of cubical cells, but unlike what obtains in the iris, they are not both pigmented, but only the outer layer.

The ciliary body extends backwards as far as the *ora serrata*, at which point the retina proper begins abruptly; the transition from ciliary body to choroid, on the other hand, is gradual, though this line is conveniently accepted as the limit of the two structures. The ora serrata is slightly more anterior on the nasal than on the temporal side.

The ciliary body is richly supplied with sensory nerve fibres derived from the trigeminal, so that great pain results from injury or acute inflammation. The ciliary muscle is supplied with motor fibres from the oculomotor nerve.

The *choroid* is an extremely vascular membrane in contact everywhere with the sclerotic, though not firmly adherent to it, so that there is a potential space between the two structures, which acts as a lymph space (Fig. 4). On the inner side, the choroid is covered by a thin elastic membrane, the *lamina vitrea*, or *membrane of Bruch*. The blood vessels of the choroid increase in size from within outwards, so that immediately beneath the membrane of Bruch there is a capillary plexus, the *choriocapillaris*. Following upon this is the layer of medium-sized vessels, while most external are the large vessels. The vessels are held together by a stroma consisting of branched pigmented connective tissue cells. It is easy to remember that the capillaries are innermost, because one of the chief functions of the choroid is to nourish the outer layers of the retina.

The choroid is supplied with sensory nerve fibres from the trigeminal.

The *retina* proper corresponds in extent with the choroid,

which it lines. As already mentioned, however, and as shown by embryological research, it is continued forwards as a double layer of epithelium as far as the edge of the pupil. If the two layers of epithelium are traced backwards, the anterior layer in the iris is found to be continuous with the outer layer in the ciliary body, and this again is continued into the hexagonal pigment epithelium, which covers the membrane of Bruch. Similarly, the posterior layer in the iris, although pigmented, passes into the inner unpigmented layer of the ciliary body, and this suddenly changes at the ora serrata into the highly complex retina proper.



Fig. 4.—Diagrammatic section of retina, choroid, and sclerotic at posterior pole of the eye. *R*, retina, with *f*, fovea centralis: *a*, internal limiting membrane; 1, nerve fibre layer; 2, ganglion cell layer; 3, internal reticular layer; 4, internal nuclear layer; 5, external reticular layer; 6, external nuclear layer; *b*, external limiting membrane; 7, rods and cones; 8, retinal pigment epithelium. Ch., choroid: *c*, membrana vitrea or membrane of Bruch; 9, choriocapillaris; 10, layer of medium-sized vessels or Sattler's layer; 11, layer of large vessels or Haller's layer. Scl., sclerotic, with 12, lamina fusca on its inner surface.

The retina proper consists of a number of layers. Most external, in contact with the pigment epithelium, is a neural epithelium, the rods and cones (Fig. 4). Following this, in order from without inwards, are the outer nuclear, the outer reticular, the inner nuclear, the inner reticular, the ganglion cell, and the nerve fibre layers. These special nervous constituents are bound together by neuroglia, the better developed vertical strands being called the fibres of Müller. The interlacement of neuroglial fibrils on the outer side form a sort of membrane which acts as a basement membrane for the rods and cones, the outer limiting membrane. Similarly on the inner side the bases of Müller's fibres spread out and form an inner limiting membrane upon the inner surface of the nerve fibre layer. Here the retina is in contact with the vitreous, which also has an extremely delicate bounding membrane, the hyaloid membrane: this is regarded by

some as identical with the inner limiting membrane of the retina.

At the optic disc the fibres of the nerve fibre layer pass into the optic nerve, the other layers of the retina stopping short abruptly at the edge of the porus opticus.

At the posterior pole of the eye, which is situated about 3 mm. to the temporal side of the optic disc, a specially differentiated spot is found in the retina of higher mammals (man and monkeys), the *fovea centralis*. As its name implies, it is a depression or pit, and here only cones are present in the neuro-epithelial layer and the other layers are almost completely absent. The fovea is the most sensitive part of the retina, and it is surrounded by a small area, the macula lutea, or yellow spot, which, though not so sensitive, is more so than other parts of the retina. It is here that the nuclear layers become gradually thinned out, while on the other hand parts of the reticular layers are specially in evidence: the ganglion cells too, instead of consisting of a single row of cells, are heaped up into several layers. There are no blood vessels in the retina at the macula, so that its nourishment here is entirely dependent upon the choroid (Fig. 5): by way of compensation the meshes of the capillary network of the choriocapillaris are particularly small here.



Fig. 5.—Blood vessels in the macular region of the human retina. (Nettleship.) The central gap corresponds with the fovea centralis. N, nasal side; T, temporal side; A, arteries; V, veins. The meshes are much smaller than at the periphery of the retina.

The so-called *optic nerve* is really a lobe of the brain, and therefore belongs properly to the central nervous system. Embryological and morphological investigations show that the bipolar cells of the retina, the nuclei of which are in the inner nuclear layer, probably correspond with the cells in the dorsal root ganglion of an ordinary sensory nerve (neurons of the first order). Similarly, the ganglion cells correspond morphologically with the cells of the nucleus gracilis, or nucleus cuneatus (neurons of the second order). Hence the part of the visual nervous mechanism which corresponds with

an ordinary peripheral sensory nerve is a microscopic cell and its processes situated within the retina itself. All the remainder is really part of the central nervous system, and we shall see that it responds to pathological processes more like the central than the peripheral nervous system.

The porus opticus is the aperture in the sclerotic through which the optic nerve passes. It varies much in shape in different cases, but in all it is traversed by a transverse network of connective tissue fibres containing much elastic tissue, the lamina cribrosa. The fibres of the nerve fibre layer of the retina pass through the meshes of the lamina cribrosa, and on the posterior side they suddenly become surrounded by medullary sheaths. These nerve fibres, which comprise the greater number of the nerve fibres in the so-called optic nerve, are the axis cylinder processes of the ganglion cells of the retina. They are therefore afferent or centripetal fibres, but the optic nerve also contains a few efferent or centrifugal fibres.

The *lens* is a biconvex mass of peculiarly differentiated epithelium. It will be remembered that it is developed from an invagination of the epidermal epiblast of the fœtus, so that what was originally the surface of the epithelium comes to lie in the centre of the lens, the peripheral cells corresponding with the basal cells of the epidermis. Just as the epidermis grows by the proliferation of the basal cells, the old superficial cells being cast off, so the lens grows by the proliferation of the peripheral cells. The old cells, however, cannot be cast off, but become massed together in the centre or nucleus: moreover the newly formed cells elongate into fibres, the lens fibres, which have a rather complicated arrangement. Without going into details, it is important to bear in mind that the nucleus of the lens consists of the oldest cells and the periphery or cortex of the youngest. Further, it must be pointed out that at an early stage the productive basal cells become limited to a single row of cubical cells covering the anterior surface. The mass of epithelium which constitutes the lens is surrounded by a hyaline membrane, the lens capsule, which is thicker over the anterior than over the posterior surface: it is a cuticular deposit secreted by the epithelial cells.

The lens in fœtal life is almost spherical; it gradually becomes flattened so as to assume the biconvex shape. It is held in place by the suspensory ligament or zonule of Zinn. This is not a complete membrane, but consists of bundles of fibrils which pass from the surface of the ciliary body to the

capsule. The flattening of the lens is due to these fibrils becoming more and more stretched as the eye grows. The fibrils pass in various directions and the various bundles often cross one another. Thus the most posterior arise from the pars plana of the ciliary body almost as far back as the ora serrata; these lie in contact for a considerable distance with the ciliary body and then curve towards the equator of the lens to be inserted into the capsule: most are inserted slightly anterior to the equator. A second group of bundles springs from the summits and sides of the ciliary processes, *i.e.*, far forwards, and passes backwards to be inserted into the lens capsule, slightly posterior to the equator. A third group passes from the summits of the processes almost directly inwards to be inserted at the equator.

It will be noticed that there is a somewhat triangular space between the back of the iris and the anterior surface of the lens, having its apex at the point where the pupillary margin comes in contact with the lens; it is bounded on the outer side by the ciliary body. This is the *posterior chamber:* it contains lymph of the same nature as the aqueous.

Behind the lens is the large vitreous chamber, containing the *vitreous humour*. This is a jelly-like material, intersected by delicate fibrils with a few cells and wandering leucocytes. It is bounded by the excessively delicate hyaloid membrane, but it is doubtful whether this is present at the anterior part where the vitreous is in contact with the lens.

The Blood Supply of the Eye.

The arrangement of the blood vessels which supply the eye is peculiar and is of great importance in considering pathological conditions.

The arteries of the eye in man are all derived from the ophthalmic artery, which is a branch of the internal carotid. The ophthalmic artery has very few and insignificant anastomoses, so that on the arterial side the ocular circulation is an offshoot of the intracranial circulation. This is not the case to so marked a degree of the venous outflow of the eye. While in man most of the blood passes to the cavernous sinus by way of the ophthalmic veins, yet it must be remembered that these anastomose freely in the orbit, the superior ophthalmic vein communicating with the angular vein at the root of the nose, and the inferior ophthalmic vein with the pterygoid plexus. Hence too great stress must not be laid upon the



THE RETINAL VESSELS (after Leber).

a.t.s., superior temporal artery. *a.n.s.*, superior nasal artery. *a.t.i.*, inferior temporal artery. *a.n.i.*, inferior nasal artery. *a.me.*, macular artery. *v.t.s.*, superior temporal vein. *v.n.s.*, superior nasal vein. *v.t.i.*, inferior temporal vein. *v.n.i.*, inferior nasal vein. *v.m.*, macular vein.

[*To face p.* 11.



THE CILIARY SYSTEM OF VESSELS (after Leber).

s.p.c., short posterior ciliary arteries. *l.p.c.*, long posterior ciliary artery. *a.c.*, anterior ciliary vessels. *C. of S.*, canal of Schlemm. *c.a.i.ma.*, circulus arteriosus iridis major. *v.v.*, venæ vorticosæ. *a.conj.*, anterior conjunctival vessels. *p.conj.*, posterior conjunctival vessels.

[*To face p.* 11.

circulation in the retina as a guide to the condition of the intracranial circulation, as has been done in the past.

The retina is supplied by the central artery, which enters the nerve on its lower surface, 15–20 mm. behind the globe. The central artery divides on or slightly below the surface of the disc into the main retinal trunks, which will be considered in detail later (Plate I.). The retinal arteries are end-arteries and have no anastomoses at the ora serrata. The only place where the retinal system anastomoses with any other is in the neighbourhood of the lamina cribrosa. The veins of the retina do not accurately follow the course of the arteries, but they behave similarly at the disc, uniting on or slightly below the surface of the disc to form the central vein of the retina, which accurately follows the course of the corresponding artery.

The uveal tract is supplied by the ciliary arteries, which are divided into three groups – the short posterior, the long posterior, and the anterior (Plate II. : Fig. 6). The short posterior ciliary arteries, about twenty in number, pierce the sclerotic in a ring around the optic nerve, running perpendicularly through the sclera, to which fine branches are given off. The long posterior ciliary arteries, two in number, pierce the sclerotic slightly farther away from the nerve, in the horizontal meridian, one on the nasal, the other on the temporal side. They traverse the sclerotic very obliquely, running in it for a distance of 4 mm. The anterior ciliary arteries are derived from the muscular branches of the ophthalmic artery to the four recti. They pierce the sclerotic 5 or 6 mm. behind the limbus or corneo-scleral margin, giving off twigs to this region, to the conjunctiva and sclerotic.

The ciliary veins also form three groups—the short posterior ciliary, the venæ vorticosæ, and the anterior ciliary. The short posterior ciliary veins are unimportant ; they do not receive any blood from the choroid, but only from the sclerotic. The venæ vorticosæ are the most important, consisting usually of four large trunks which open into the ophthalmic vein. They enter the sclerotic rather behind the equator of the globe. two above and two below : they pass very obliquely through the sclera. The anterior ciliary veins are smaller than the corresponding arteries, since they receive blood only from the outer part of the ciliary muscle.

Of these ciliary vessels the short posterior ciliary arteries supply the whole of the choroid, being reinforced anteriorly by anastomosis with recurrent branches from the ciliary body.



FIG. 6.—Blood vessels of the human uveal tract, injected, from the optic disc to the edge of the iris. (After Leber.) Arteries, black; veins, paler. Running up the centre is a long posterior ciliary artery. Two vorticose veins and their tributaries are seen. The capillaries are only partially filled in.

The ciliary body and iris are supplied by the long posterior and anterior ciliary arteries. The blood from the whole of the uveal tract, with the exception of the ciliary muscle, normally leaves the eye by the venæ vorticosæ only.

The two long posterior ciliary arteries pass forwards between the choroid and the sclerotic, without dividing, as far as the posterior part of the ciliary body. Here each divides into two branches (Fig. 6): they run forwards in the ciliary muscle, and at its anterior part bend round in a circular direction, anastomosing with each other and thus forming the circulus arteriosus iridis major. This is situated in the ciliary body at the base of the iris: from it the ciliary processes and iris are supplied. Other branches from the major arterial circle run radially through the iris, dividing dendritically and ending in loops at the pupillary margin. A circular anastomosis takes place a little outside the pupillary margin, the circulus arteriosus iridis minor.

The tributaries of the vorticose veins, which receive the whole of the blood from the choroid, are arranged radially, the radii being bent, so as to give a whorled appearance hence their name. The veins of the iris are collected into radial bundles which pass backwards through the ciliary body, receiving tributaries from the ciliary processes. Thus reinforced, they form an immense number of veins running backwards parallel to each other through the smooth part of the ciliary body. After reaching the choroid they converge to form the large anterior tributaries of the vorticose veins. The veins from the outer part of the ciliary muscle on the other hand pass forwards and unite with others to form a plexus, part of which is the so-called canal of Schlemm. These vessels drain into the anterior ciliary veins. The marginal loops of the cornea and the conjunctival vessels are branches of the anterior ciliary (Plate II.).

CHAPTER II.

PHYSIOLOGY.

In order that the eye may satisfactorily perform its duties as an organ of vision it is essential that a sharp image of objects in the outer world shall be formed upon the retina. This is effected by means of a series of curved surfaces, and the curvature of these surfaces and their relative positions to each other must be kept constant. For this purpose it is necessary that the walls of the globe should be kept stretched.

If a small canula connected with a narrow-bored mercury manometer is pushed into the anterior chamber or into the vitreous of an animal it will be found that the mercury in the manometer will rise about 25 or 30 mm. (Fig. 7). In other words, the contents of the eyeball, which are for the most part fluid, exert a pressure upon the inner side of the walls which is about 25 mm. of mercury greater than the atmospheric pressure which falls upon the outside of the walls; the walls are thus kept well stretched.

This pressure inside the eye is called the intraocular pressure, or the *tension*, of the eye. (These terms are used indiscriminately, though it is not quite accurate to do so.) In order that the tension of the eye may be kept up, work must be done by the organism, and a moment's consideration will make it evident that the source of energy must be the blood pressure. This conjecture has been proved conclusively true by experiments on animals, in which it is found that the intraocular pressure generally follows passively the changes in the general blood pressure (Fig. 8).

It is obvious that it must be the fluid constituent of the contents of the globe which keeps up the internal pressure. This fluid fills the anterior and posterior chamber and permeates the vitreous. It is comparable to the lymph which bathes the tissues in other parts of the body, and it is indeed the lymph of the eye. In other parts of the body the chief function of the lymph is to carry food material to the tissue

cells, and to carry away the effete products of the cell metabolism. It has a further function of keeping up the normal tissue tension. Both of these functions attain an unusual degree of importance in the case of the eye. We have already seen the necessity for keeping up the normal tension of the eye. As regards the nourishment of the cells, our review of the anatomy of the eye has shown that there are large areas, notably the whole of the lens and the vitreous, which possess no blood supply. They are dependent entirely for their nutrition upon the lymph.

If water were to be forced into an impermeable elastic bag, the internal pressure might be kept indefinitely above the pressure upon the outside of the bag. Such an arrangement for keeping up the intraocular pressure would be unsatisfactory owing to the function of the lymph in nutrition. The stagnant fluid would soon lose all its food material, which would be used up, and it would become loaded with the excreted products of the cells, which would have a very deleterious effect upon them. Hence it is essential that the lymph shall be constantly renewed. This occurs in the eye. The lymph is continually being renewed, but at the same time it is equally rapidly being removed, and thus the amount present is kept constant.



FIG. 7.—Diagram of manometer placed in communication with anterior chamber, showing that the normal intraocular pressure is about 25 mm. of mercury.

It has been shown that in other parts of the body lymph is formed by filtration out of the capillaries: the amount poured out and its chemical composition are dependent upon the difference in pressure on the two sides of the filtering membrane, *i.e.*, the capillary wall. On the one side there is the relatively high intracapillary pressure, on the other the low tissue tension. Exactly the same is found to be the case in the eye. Here the intracapillary pressure is higher than in other parts of the body, but the extracapillary pressure is also higher than usual, being as much as 25 mm. of mercury, *i.e.*, the intraocular pressure. The difference between the two pressures is, however, less in the eye than elsewhere. Con-

sequently we may expect to find, on the one hand, that the amount of lymph produced in a given time is small, and, on the other hand, that its chemical composition will differ materially



FIG. 8.—Tracing of right carotid blood pressure, intraocular pressure from canula in anterior chamber of left eye of a dog, which was fully under the influence of morphia, A.C.E. mixture, and curare. Stimulation of the vasomotor centre, showing that the intraocular pressure follows passively all the changes in the general blood pressure as produced by Traube-Hering curves and constriction of the arterioles of the splanchnic area.

from that of lymph elsewhere. Both these conjectures have been substantiated. The rate of production is so slow that it probably takes nearly an hour for the aqueous in the anterior

chamber to be renewed. The chemical composition of the aqueous is consistent with the view that filtration is going on at a small difference of pressures, for under these circumstances we shall expect very little of the large moleculed proteids of the blood to be able to get through the capillary wall. As a matter of fact the aqueous contains excessively little proteid (0·045 per cent.). In spite of the fact that the conditions of the production of aqueous are consistent with the view that it is a simple process of filtration, it is more generally spoken of as a secretion. Though the process is essentially a filtration, it is not to be expected that membranes covered with living endothelial and epithelial cells will behave like dead inert membranes. That these cells exercise a selective activity is shown by the fact that while certain substances, *e.g.*, agglutinins and precipitins, are allowed to pass freely, others, such as hæmolysins, do not pass into the aqueous. There is therefore justification for the use of the term secretion.

We have already noticed the resemblance of the ciliary processes to the glomeruli of the kidney. It is always dangerous to deduce physiological function from anatomical structure alone, but in the case of the ciliary body experiment and clinical and pathological observation alike confirm the conjecture that the processes are the chief if not the only site of the production of the aqueous in the human eye. If the iris and ciliary body are removed, as can be done in the rabbit without immediate destruction of the eye, no more aqueous is formed and the eye rapidly becomes soft and shrinks. That the effect is not due to the absence of the iris is shown by the fact that the iris is sometimes almost completely absent as a congenital defect, yet the aqueous is normally secreted. We may therefore conclude that the aqueous is secreted by the ciliary processes.

The greater part of the lymph thus produced passes into the posterior chamber. Thence it passes forwards through the pupil into the anterior chamber, the free communication between the two chambers being facilitated by the perpetual movements of the iris. A small quantity—not more than one-fiftieth part—of the aqueous secreted by the ciliary body passes backwards into the vitreous, which it nourishes. Situated in the vitreous, running directly forwards from the optic disc almost as far as the posterior pole of the lens, is a small tubular lymph space, the canal of Cloquet. It is the lymph space which originally surrounded the fœtal hyaloid artery; it persists after the latter disappears. The lymph of the vitreous

finds its way into this space, which communicates with the perivascular lymph spaces surrounding the central vessels of the retina. By this route the lymph passes out of the eye through the optic nerve into the vaginal space. Only a very small proportion of the lymph produced by the ciliary processes—as mentioned probably not a fiftieth part—finds its exit from the eye here. What becomes of the remainder? We have already said that it passes forwards into the anterior chamber. It was at one time thought that it transuded through the cornea and escaped into the conjunctival sac. This is not so: it is now known that it filters away through the angle of the anterior chamber into the venous plexus which is called the canal of Schlemm. Having passed into the canal of Schlemm the effete lymph is carried away by the anterior ciliary veins.

It might be expected that so vascular a tissue as the choroid would be concerned in the production of lymph. It is probable, however, that it produces only sufficient to nourish its own tissues and the outer layers of the retina. The retinal capillaries extend no deeper into the retina than the outer reticular layer; consequently the rods and cones and the pigment epithelium are dependent upon the choroid for their nutrition. The choroidal lymph passes out of the eye chiefly by way of perivascular lymph spaces around the vorticose veins, a small quantity passing into the subchoroidal space and so into the lymph sheaths of the posterior ciliary vessels. The retinal lymph passes away by perivascular lymph sheaths through the optic nerve into the vaginal space. The retina and choroid, then, produce a sufficiency of lymph for their own requirements only, and have nothing to do with sustaining the intraocular tension.

Similarly it might be thought that the vascular iris provided some of the aqueous. It is not easy to disprove this hypothesis; but it is easy to show that the iris is to some extent concerned in absorption, and it is highly improbable that the same structure carries on such opposite functions.

It has already been stated that sharp images of external objects must be formed upon the retina if the latter are to be seen clearly. Before considering how this is effected it will be advisable to refresh the reader's memory of the elementary principles of optics. I would seriously impress upon the student that success in the diagnosis, and hence in the treatment, of diseases of the eye is impossible if such elementary principles of optics as are set forth here are not thoroughly mastered.

CHAPTER III.

ELEMENTARY OPTICS.

If white light, such as sunlight, is passed through a suitable prism or diffraction grating a spectrum is formed, consisting of rays differing from each other in wave-length. Of these certain are visible and appear to the majority of people as pure colours, viz., red, orange, yellow, green, blue, and violet in the order named, the red having the longest and the violet the shortest wave-length. The visible spectrum extends from about 723 $\mu\mu$ at the red end to 397 $\mu\mu$ at the violet end, or roughly from 700 $\mu\mu$ to 400 $\mu\mu$. Beyond the red end are infra-red rays of greater length which when absorbed cause a rise in temperature and are commonly known as heat rays. Beyond the violet end are waves of smaller length, the ultra-violet rays, which are capable of causing chemical action. The longer visible rays also cause a rise in temperature, and the visible rays are also actinic, though less so than the infra-red and ultra-violet respectively. Glass absorbs some of the heat rays and many of the ultra-violet. Prisms and lenses made of quartz allow most of the ultra-violet rays to pass unimpeded. The media of the eye are uniformly permeable to the visible rays between 660 $\mu\mu$ and 390 $\mu\mu$. With regard to ultra-violet rays, the cornea absorbs all rays beyond 295 $\mu\mu$, the lens all rays beyond 350 $\mu\mu$, the vitreous shows an absorption band with its maximum at 270 $\mu\mu$ (E. K. Martin). Rays between 400 $\mu\mu$ and 295 $\mu\mu$ can therefore reach the lens, those between 400 $\mu\mu$ and 350 $\mu\mu$ can reach the retina in the normal eye, and those between 400 $\mu\mu$ and 295 $\mu\mu$ can reach the retina in the aphakic eye. Whenever absorption occurs there is the possibility of pathological changes resulting. Sunlight at the lower sea-levels is poor in ultra-violet rays, which fall off rapidly in intensity beyond 380 $\mu\mu$. Ordinary glass used for spectacles absorbs rays beyond 350 $\mu\mu$. Heat radiation from 1,100 $\mu\mu$ to 700 $\mu\mu$ passes into the eye almost unchecked, and a large amount of it reaches the retina (Hartridge and Hill). The pigment epithelium on the back of the iris absorbs heat radiation of all wave-lengths, and the same is probable of the retinal epithelium at the back of the eye.

It is a familiar fact that a candle flame emits light in all directions. The light is transmitted in straight lines, so that

we may imagine the light coming from the candle as consisting of an immense number of straight lines, all intersecting in some part of the flame. If we consider a minute point in the flame, then all the straight lines which cross in this point may be said to diverge from it. Each of these hypothetical straight lines is called a ray.

Now, every point on such a ray represents, or is the image of, the point of light from which it springs. This is shown very clearly by a simple experiment carried out in a dark room. Make a pinhole in a piece of cardboard (Fig. 9, A), and hold the cardboard in front of the candle (C) at a little distance from it. Beyond the cardboard hold up a white screen (B), so that the cardboard is between the screen and the candle. A dim image (D) of the flame will be thrown upon the screen, and it will be noticed that it is upside down, *i.e.*, an inverted image of the flame is formed. This is due to the fact that the carboard cuts off all the rays of light from the candle except such as can pass through the hole. The only rays from the top of the flame which can pass through the hole are those which are caught upon the lower part of the screen. They represent the top of the flame; hence they reproduce its shape accurately. The image is very dim because only a few rays of light can pass through the small hole. Now make another hole a little distance away from the first. Another inverted image of the flame is seen. If a dozen holes are made, a dozen images appear. If two holes are very close together the images will overlap. If a large hole is made, so many images overlap that all resemblance to the original flame is lost, and part of the screen becomes uniformly illuminated. If we take away the cardboard altogether the whole screen becomes illuminated, and we now know that this is because we have an infinite number of images of the flame all overlapping each other.



FIG. 9.

Light travels with different velocities in different media. If the velocity is less in one medium than another the first medium is said to be optically denser than the second.

When light, travelling in one medium, meets another

medium it breaks up into two parts: part is *reflected* back into the first medium: part is *refracted* into the second medium. If the second medium is opaque none of the light is refracted.



FIG. 10.—The ray from P which strikes the mirror AB at Q is reflected to R, so that PQ and QR are in the same plane, viz., that of the paper, and the angle of incidence, i, is equal to the angle of reflection, r.

REFLECTION.

Let us now consider what happens to a ray of light when, travelling in one medium, it is reflected from the surface of a denser medium. We have already said that its direction is altered. Before it meets the surface it is called an incident ray; after it leaves the surface it is called the reflected ray. If a line is drawn at right angles to the surface at the point where the incident ray meets it, it is found to be an invariable rule that the incident ray makes the same angle with this line, which is called the normal, that the reflected ray makes with it. Put in formal language, this law of reflection is that *for all surfaces the angle of incidence is equal to the angle of reflection, and is in the same plane with it* (Fig. 10).



FIG. 11.

Plane Mirrors.—Let us apply this rule to an ordinary flat mirror (Fig. 11). If P is a luminous point in front of the mirror AB, the ray PQ will be reflected towards R, and the ray PS towards T; *i.e.*, the reflected rays QR and ST appear to come from p, a point as far behind the mirror as P is in front of it. As the rays QR and ST have to be produced backwards in order that they may meet, no real image is formed, and such an image is called a virtual image. Note that the rays reflected from a plane mirror are divergent. The same reasoning holds good for every point on the object PV, its image being pv as far behind the mirror

as the object is in front of it: moreover, the size of the image is equal to that of the object.

Concave Mirrors.—Here the normal to the surface is the radius of the sphere. If AH (Fig. 12) is part of the section of a concave mirror and PB is an object, K being the centre of the sphere, then the line HKB is called the axis, and H the apex of the mirror. The ray PK through the centre of the sphere will obviously be reflected along itself, so that the image of P must be on PK. The ray PA, parallel to the axis, will meet PK in *p*. Hence *p* is the image of P. Now it is found that all rays parallel to the axis and not very far removed from it cut the axis in the same point, F, and this point bisects the line HK. This point is called the principal focus of the mirror. If the object PB were removed a very great distance away from the mirror all the rays which fell upon a small portion of the mirror near H would diverge so little from each



Fig. 12.

other that they would all be practically parallel to BH, and the image of PB would be extremely small and situated at F. In each of these cases the image is an inverted one of the object.

It is an axiom of optics that the direction of the rays is reversible. Hence, if *pb* were an object, it would have its image at PB, and if there were an object at F, all the rays from it reflected by the mirror would be parallel to the axis, and the image would be infinitely large and situated at infinity.

What would happen if the object were situated between F and H? In that case (Fig. 13) the rays would diverge after reflection as if they came from an object behind the mirror, much as they do with a plane mirror. The image would therefore be a virtual one, situated behind the mirror: it would be erect and larger than the object.

The important fact to remember with regard to concave

mirrors is that if the object is farther away from the mirror than its focal distance, *i.e.*, than half its radius of curvature, the image is a real inverted one situated also in front of the mirror. This is the condition which is almost always present in the ordinary use of ophthalmic instruments.



FIG. 13.—The ray from P parallel to the axis is reflected through F, the principal focus. The ray FP is reflected parallel to the axis. The ray KP is normal to the surface, and is therefore reflected on itself. Any two of these rays will give the situation of *p*, the image of P.

Convex Mirrors. We are not accustomed to use convex mirrors in ophthalmic instruments, but it is necessary to know what happens with them, since the cornea acts as a convex mirror. Here, as will be seen from Fig. 14, the image is always virtual, erect, and smaller than the object. As with the concave mirror, if the object is a long way off, the image will be



FIG. 14.—Reflection by a convex mirror. The description of Fig. 13 applies equally to this case.

situated at the principal focus, *i.e.*, at a distance equal to half the radius of curvature behind the mirror.

REFRACTION.

We have now to consider what happens to the refracted ray when the incident ray, travelling in one medium, *e.g.*, air, meets an optically denser medium, *e.g.*, glass. We have already said that the light will now travel more slowly. It follows directly from this fact that it will be deviated towards

the normal to the surface, and it will be more deviated the greater the difference in optical density between the two media. If the density of air is taken as unity, then the ratio of its density to that of the second medium is called the *index of refraction* of the medium.



Fig. 15.—Refraction by a plane lamina.

Plane Lamina.—Let us see what happens when an incident ray, such as PQ (Fig. 15), meets the surface of a plate of glass with parallel sides. It will be deflected towards the normal, *ab*. When the ray passes out of the glass on the other side, it will obviously be deflected away from the new normal, *cd*, just as much as PQ was deflected towards it. Hence the emergent ray RS will be parallel to the incident ray PQ. If the plate of glass is very thin, RS will be practically continuous with PQ.

Prisms.—If we imagine one side (CD) of the plate in the last figure to revolve round R, we shall be able to understand the nature of refraction by a prism. QR will now make a larger angle with the new normal *ef* than it did with the old one *cd*. Consequently the angle of refraction will also be larger, *i.e.*, the new direction of the emergent ray will be RT. In other words, the ray is deviated towards the base of the prism.



Fig. 16.—Refraction by a prism.

The ray PQRS in Fig. 16 is said to pass symmetrically through the prism. Under these circumstances, if the prism is made of crown glass, the deviation of the ray is approximately equal to half the refracting angle of the prism, *a*.

We are accustomed to project objects along the direction of the rays of light as they enter the eye, and in doing so we ignore the effect of refraction, since it enters relatively little into our everyday experience. If, therefore, we look at a candle P through a prism, as in Fig. 17, the light will appear to come from *p*. Objects, then, seen through a prism, appear displaced towards the apex of the prism.



FIG. 17.—Displacement of objects seen through a prism. The object P appears to be situated at *p*.

Lenses. — Ordinary lenses are pieces of glass with spherical surfaces. The line passing through the centres of curvature of the surfaces is called the axis of the lens. Fig. 18 shows the chief varieties of lenses, viz., (1) biconvex, (2) biconcave, (3) plano-convex, (4) plano-concave, (5) convexo-concave or meniscus: these names require no further explanation.

The effect of a biconvex lens upon rays of light meeting it is very similar to what would occur if it were replaced by two prisms set base to base (Fig. 19).



FIG. 18.—Types of lenses.

If the incident rays are parallel to the axis they will be refracted in such a manner that they all cross the axis in a single point upon the other side of the lens. This point is called the principal focus of the lens, and its distance from the lens is called the focal distance or length of the lens. When the lens has the same medium, *e.g.*, air, on each side of it, the

two principal foci, one on each side of the lens, are situated at equal distances from it. For thin glass lenses of low power the focal distance is equal to the radius of curvature of the two surfaces when these are equally curved. If there is an object



Fig. 19.

a very long distance away from the lens, the rays which come from it are practically parallel. Hence in this case an image of the object will be formed by the lens at its principal focus: it will be inverted and very small. If the object is gradually brought nearer and nearer to the lens (Fig. 20) the image will



Fig. 20.—The ray PE, parallel to the axis, is refracted through the second principal focus F_2. The ray PF, through the first principal focus, is refracted parallel to the axis. The ray PO, through the optical centre of the lens, is not deflected. Any two of these rays give the situation of *p*, the image of P.

recede farther and farther from it: from being very small it will grow larger and larger, until, when the object is at the principal focus, the image will have receded to infinity, and it will be infinitely large, *i.e.*, all the rays coming from the object at the principal focus are parallel to the axis and to each other after retraction. If the object is brought still closer to the lens than its focal distance (Fig. 21) it will be found that its image is a virtual one behind the object, and that it is erect and

larger than the object. The positions of the object and image bear a constant relationship to each other and are called conjugate foci.

There is a point in the middle of a biconvex lens which is



FIG. 21.—The description of Fig. 20 applies equally to this figure.

called its optical centre. With thin lenses any ray which passes through this point is practically not deviated at all. It is easy to understand why this is so. If PQRS (Fig. 22) is such a ray and tangents are drawn to the two surfaces at the points Q and R, these two tangents will be parallel to each other. Consequently, the lens acts for such a ray exactly as if it were a plate with parallel sides, and we have already seen that in such a case the emergent ray is parallel to its original direction. If the lens is very thin the refracted ray will be practically continuous with the incident ray.



FIG. 22.—Properties of the optical centre of a lens.

If we know these facts, viz., that rays passing through the optical centre are not deviated, and that rays passing through the principal focus are parallel to the axis after refraction, we can easily construct the image of an object in any given position. Thus, in Fig. 20, if PB is an object, the ray PO through the optical centre O will not be deviated ; the ray PE parallel to the axis will pass through the second principal focus F_2 ; and the ray PF_1 through

the first principal focus will be parallel to the axis after refraction. Hence *pb* must be the image of PB.

The effect of a biconcave lens upon rays of light meeting it



Fig. 23.

is very similar to what would occur if it were replaced by two prisms set apex to apex (Fig. 23).

Here, if the incident rays are parallel to the axis they will be divergent after refraction, and the amount of divergence of the individual rays will be such that if they are produced backwards they will all cross the axis in a single point upon the same side of the lens that they came from. This and the



Fig. 24.—The description of Fig. 20 applies equally to this figure.

corresponding point on the other side of the lens are called the principal foci. The biconcave lens also has an optical centre, situated upon the axis within it and having the same properties as in the case of the convex lens. The image of any object formed by a concave lens can be constructed in exactly the same manner as for a convex lens (Fig. 24). It will be found that in every position of the object the image is always virtual, erect and smaller than the object.

Plano-convex and plano-concave lenses act like biconvex and biconcave respectively, but in them the optical centre is on the curved surface at the point where the axis cuts it. Menisci act as convex or concave lenses according as the convex or the concave surface has the greater curvature. In them the optical centre is outside the lens.

It will have been noticed that the refractive power of a lens varies inversely as the focal distance, *i.e.*, a lens with a short focal distance will bend the rays more than one with a longer focal distance. It is necessary to have some system of numbering lenses so as to indicate their refractive power. The most convenient system for ophthalmic purposes is that which takes a lens with a focal distance of 1 metre as a standard. Such a lens is said to have a refractive power of 1 *dioptre*. A lens with a focal length of half a metre will be twice as strong as one whose focal length is 1 metre: the refractive power of such a lens is therefore 2 dioptres. Similarly, a 3 D (3 dioptre) lens has a focal length of one-third of a metre, or 33 cm.; a 4 D lens, 25 cm.; and so on. It is important to remember that in this system the standard is a metre, not a centimetre or a millimetre; otherwise confusion may arise.

Lenses were formerly numbered according to their focal lengths measured in inches. Since the inch has a different value in different places, the method is unsatisfactory. Prescriptions for spectacles are, however, sometimes met with in this notation. They are easily transformed into the dioptric system by remembering that there are 40 inches (roughly, or 36 Paris inches) in 1 metre. Therefore a 40 inch lens = 1 D; a 20 inch lens = 2 D; a 4 inch lens = $\frac{40}{4}$ = 10 D; and so on: a lens of focal length = 4 Paris inches = $\frac{36}{4}$ = 9 D.

Convex lenses are indicated by a plus sign (+), concave by a minus sign (−) before the number.

Cylindrical lenses are also used in ophthalmology; their nature and use will be considered at a later stage.

We often wish to find out whether a lens is convex or concave, and what its refractive power is. There are several ways of doing this, but the simplest is with the assistance of the trial case. Hold a convex lens up near the eye and look at distant objects through it; then move the lens a little from side to side: the distant objects will seem to move in the opposite direction to that in which the lens is moved. If we repeat the process with a concave lens the objects seem to

move in the same direction as the lens. The reason is to be found in the fact that a convex lens forms an inverted, whilst a concave forms an erect image. If we place two lenses of opposite sign but equal curvature in contact with one another the combination will make a plate with parallel sides: such a plate, as we know, does not practically deflect the rays of light at all. Hence we can determine the strength of a lens by exactly neutralising it with a lens of the opposite sign out of the trial case. Let us take a concrete example, a particular lens which we wish to determine. We hold it up and find that distant objects seem to move in the opposite direction to the lens. We know that it is a convex lens. We then put a weak concave lens in contact with it and repeat the process. We find that with a — 2 D lens objects still seem to move in the opposite direction, though not so much. With a — 3 D lens there is only a trace of movement, and with a — 3·5 D lens there is no movement at all. We conclude that the original lens was + 3·5 D. In performing this test it is important to have the two lenses as closely in contact as possible, and also



Fig. 25.



Fig. 26.

to have their centres in contact. If the centre of one lens is higher than that of the other they will obviously not counteract each other exactly. If they are not in contact the result will be either too high or too low.

When the lenses are in contact the refractive power of the combination (D) is equal to the algebraical sum of the refractive powers of the two lenses (d_1, d_2): *i.e.*, $D = d_1 + d_2$, or $\frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2}$ where F, f_1, f_2, are the respective focal distances (Fig. 25).

Suppose, however, that two convex lenses are separated by a distance c (Fig. 26). The lens A will make parallel rays converge towards a, but after a distance c they meet the lens B : hence the convergence of the rays is not expressed by $\frac{1}{f_1}$, but by $\frac{1}{f_1 - c}$. Therefore the combined effect of the lenses, D, or $\frac{1}{F}$, is now equal to $\frac{1}{f_1 - c} + \frac{1}{f_2}$.

If the second lens (B) is a concave one (Fig. 27) its effect will



Fig. 27.

be one of divergence, so that it must have a negative sign, and D will now be equal to $\frac{1}{f_1 - c} - \frac{1}{f_2}$.

It is to be noted that in the formula

$$\frac{1}{F} = \frac{1}{f_1 - c} + \frac{1}{f_2}$$

F is now the posterior focal length, the incident light impinging upon the lens whose focal length is f, and being directed towards the lens whose focal length is f_2. The following formula gives the equivalent focal length (F_e) of the combination, irrespective of the direction of light :—

$$F_e = \frac{f_1 f_2}{f_1 + f_2 - c}.$$

Examples : (1) $f_1 = 333$ mm. ; $f_2 = 250$ mm. ; $c = 133$ mm.

$$\text{Then } \frac{1}{F} = \frac{1}{333 - 133} + \frac{1}{250} = \frac{1}{111}.$$

That is, the combination of a + 3 D lens with a + 4 D, separated by a distance of 133 mm., will be that of a + 9 D lens, instead of a + 7 D if they had been in contact.

(2) $f_1 = 333$ mm. ; $f_2 = -83$ mm. ; $c = 133$ mm.

$$\text{Then } \frac{1}{F} = \frac{1}{333 - 133} - \frac{1}{83} = -\frac{1}{142}.$$

That is, the combination of a + 3 D lens with a − 12 D, separated by a distance of 133 mm., will be that of a − 7 D lens, instead of − 9 D if they had been in contact.

CHAPTER IV.

ELEMENTARY PHYSIOLOGICAL OPTICS.

The eye as an optical instrument very much resembles an ordinary photographic camera. The latter consists of a dark chamber with an aperture in front containing a strong convex lens, and with a movable back behind. The effect of the lens is exactly like that shown in Fig. 20. PB will represent the object to be photographed; the movable back is adjusted so that it occupies the position of *pb*, in which case a sharp image of the object will be thrown upon the ground glass which forms the back. The ground glass is then replaced by a sensitive plate, and the photograph is taken.

In the eye the retina corresponds with the sensitive plate. Instead of having only one lens in the front aperture, represented by the crystalline lens, there is also a curved plate with parallel sides, the cornea, which acts like another lens, and indeed has a much stronger optical effect than the crystalline lens. The object of this more complicated arrangement is to shorten the focal distance of the system, so that the eye may be shorter and more compact.

From this analogy we see that the eye, from the optical point of view, acts like a strong convex lens. We have already stated that when a lens has the same medium on each side of it the anterior and the posterior focal distances are equal to one another. This is not the case in the eye. Here the medium in front is air, while behind the lens there is the vitreous, which has a higher refractive index, rather more than that of water. Hence the anterior and posterior principal focal distances are not equal, the anterior being about 13 mm. in front of the cornea, and the posterior about 23 mm. behind it.

The cornea has about the same optical density or refractive index as the aqueous, which is also equal to that of the vitreous. The anterior surface of the cornea may be regarded as nearly spherical, the radius of curvature being 8 mm. The centres of curvature of the cornea and the two surfaces of the lens are all on the same straight line, which is called the *optic*

axis. When a ray of light meets the cornea (Fig. 28) the ray will be deflected towards the normal, *i.e.*, towards the radius drawn through the point of incidence. It will pass through the layers of the cornea in the new direction, and will also continue in the same direction through the aqueous, for, as we have said, the refractive index of the aqueous is the same as that of the cornea. When the ray meets the lens, which has a greater refractive index than the aqueous, it will again be deflected in the same sense, *i.e.*, towards the axis upon which the cornea and lens are centred.

We have seen that in the case of a lens, and the same is true of any homocentric optical system, parallel rays meet at the principal focus. Hence, if parallel rays fall upon the cornea, they will be brought to a focus 23 mm. behind it.



Fig. 28.—Emmetropic eye. Parallel rays are brought to a focus on the retina.

Now, the rays which are emitted by a luminous body are divergent. If, however, the object is a long distance away, the individual rays in any small bundle will diverge so little from each other that they may be regarded as practically parallel. This is the case with the small bundles of rays which are able to enter the pupil of the eye. Hence, as in the case of a convex lens (*vide* p. 26), the image formed by the eye of these distant objects will be situated at the principal focus, *i.e.*, 23 mm. behind the cornea. But that is exactly the distance of the retina from the cornea in the normal eye. Hence we see that the normal eye in its condition of rest is so constituted that distant objects form their images upon the retina (Fig. 28).

The optic axis, produced backwards to meet the retina, cuts it almost exactly at the fovea centralis. Hence, any distant object on the prolongation forwards of the optic axis will

have its image at the fovea, which is the best spot for distinct vision.

We notice that, just as with a convex lens, the image is inverted. It is re-inverted psychologically in the brain.

It is easy to find the size of the retinal image which any



Fig. 29.—Nodal point of the eye, visual angle, and relative sizes of object and retinal image.

external object will form, since the eye possesses an optical centre, which, however, is usually called the *nodal point* (N), quite similar to the optical centre of the lens. In the eye this point is situated upon the optic axis near the back of the lens. As in the case of lenses, any ray which passes through this point will not be appreciably deflected. If, therefore, there is



Fig. 30.—Hypermetropic eye. Parallel rays tend towards a focus behind the retina.

an object PB (Fig. 29) in front of the eye, the size of its retinal image *pb* is found by joining the extremities of the object and the nodal point and producing these lines until they meet the retina. The lines will enclose an angle, PNB, which is called the *visual angle;* in other words, the angle subtended by the object at the nodal point is called the visual angle. It is of course equal to the angle *p*N*b*, which is subtended by the retinal image at the nodal point.

In some eyes the retina is not situated in exactly the right place for the images of distant objects to be clearly focussed upon it. It may be too far forward (Fig. 30), or too far back (Fig. 31); in the former case they are said to be hypermetropic, in the latter myopic. If we consider the effect upon parallel rays we shall see that in the hypermetropic eye they have not had space to come to a focus, whereas in the myopic eye they have not only come to a focus, but have commenced to diverge. In each case a blurred image will be formed upon the retina, and vision will be impaired. Such conditions are called errors of refraction or *ametropia* (ἀ, privative, μέτρον, measure; not according to measure). In contradistinction to hypermetropia and myopia the normal condition is called emmetropia.

It has already been stated that in optics the direction of



Fig. 31.—Myopic eye. Parallel rays are brought to a focus in front of the retina.

the rays is reversible. Let us imagine a minute point on the retina to be luminous. It will give out rays which will diverge in all directions. Some of these rays will meet the lens and cornea and pass out of the eye. Now, in the emmetropic eye, those rays which get through the pupil will have to submit to exactly the same optical deviations as the parallel rays falling upon the cornea did when they passed into the eye and came to a focus on the retina. Hence, on the principle of the reversibility of the rays, the rays coming from a point on the retina will be parallel to each other when they leave the eye (Fig. 28).

Suppose, however, that the eye is hypermetropic owing to being too short (Fig. 32). The rays coming from a point on the retina will not have had space to diverge so much from the axis as the corresponding rays of the emmetropic eye before they fall upon the back of the lens. (Compare the effect of

placing an object closer to a convex lens than its principal focus (Fig. 21).) The lens and aqueous and cornea will therefore cause them to converge less than in the emmetropic eye. They will therefore still be divergent when they leave the eye, though of course not so divergent as when they were passing through the vitreous. In fact, their direction will be the same



FIG. 32.—Hypermetropic eye. Rays from a point on the retina are divergent when they emerge from the eye, as if they came from the point, R, behind the eye.

as if they came from a point behind the eye. The nearer the retina is to the lens, the more divergent they will be, and the nearer to the back of the eye will be the point from which they seem to come. This virtual point (R) behind the eye is called the *remote or far point* of the eye. The point on the retina and this point behind the eye are really conjugate foci (Fig. 32).

Suppose now that the eye is myopic owing to being too long



FIG. 33.—Myopic eye. Rays from a point on the retina are convergent when they emerge from the eye, so that they cross at a real point, R, in front of the eye.

(Fig. 33). The rays coming from a point on the retina will have become more divergent from the axis than the corresponding rays of the emmetropic, and still more divergent than those of the hypermetropic eye before they fall on the back of the lens. (Compare the effect of placing an object farther away from a convex lens than its principal focus (Fig. 20).) The refractive media in front will therefore cause them to converge more than in the emmetropic eye. They will

hence be convergent when they leave the eye, and will cross in a point (R) somewhere in front of the eye (Fig. 33). The farther the retina is from the lens, *i.e.*, the higher the degree of myopia, the more convergent they will be, and the nearer to the front of the eye will be the point where they cross. This point is again the conjugate focus to the point on the retina, but in this case it is a real point. It is also called the remote or far point of the eye.

Where then is the far point of the emmetropic eye? We have seen that in each of the other conditions it is where the rays emitted from a point on the retina meet after emerging from the eye. In the emmetropic eye the emergent rays are parallel to each other. But parallel rays meet at infinity; therefore the far point of the emmetropic eye is at infinity.

It is obvious that, in hypermetropia, if we give the rays the requisite amount of convergence before they enter the eye



FIG. 34.—Hypermetropic eye. Parallel incident rays brought to a focus on the retina by means of a suitable convex lens.

they will be brought to a focus upon the retina. We can do this by placing a convex lens in front of the eye (Fig. 34). This is what is done by means of spectacles. The refractive or convergent power of a convex lens is the reciprocal of its focal distance. Hence in hypermetropia of 1 D, a convex lens of 1 D or 1 metre focal distance placed in contact with the cornea will direct parallel rays towards a point 1 metre behind the eye, *i.e.*, to the far point of the eye. Such a lens acting in combination with the refractive force of the eye would bring the rays to a focus on the retina. But lenses cannot be worn in contact with the cornea. If the lens is placed 20 mm. in front of the cornea its focal length will have to be 1,020 mm. instead of 1,000 mm. (*vide* p. 31), but this small difference is negligible, and we are accustomed to measure errors of refraction by the strength of the lens which is required when it is placed in the ordinary position of a spectacle lens (Fig. 34).

Similarly in myopia, if we give the rays the requisite amount of divergence before they enter the eye they will be brought

to a focus upon the retina. We do this by placing a concave lens in front of the eye (Fig. 35). Here we should want a –1 D lens in contact with the cornea to correct a myopia of 1 D, *i.e.*, an eye whose far point is 1 metre in front of the eye. Since the glass has to be worn about 20 mm. in front of the eye it will have to be rather stronger, *i.e.*, it will have to be of a focal distance of 980 mm. instead of 1,000 mm.

There is an advantage in having the correcting glass in ametropia in the position of the anterior focus of the eye, because under these conditions the size of the retinal image is exactly the same as if the eye were emmetropic (Figs. 36, 38). The anterior focus is about 13 mm. in front of the eye, and this is so close that the lashes are apt to rub against the glasses, soiling them and causing discomfort. Hence spectacles are usually placed slightly farther away. We have already discovered one effect of this, viz., that the convex glass in



Fig. 35.—Myopic eye. Parallel incident rays brought to a focus on the retina by means of a suitable concave lens.

hypermetropia has to be rather weaker, and that the concave glass in myopia has to be slightly stronger. It also has an effect upon the size of the retinal image, making it larger in hypermetropia and smaller in myopia (Figs. 37, 39). The increase in size in hypermetropia is advantageous, but the diminution in myopia is a disadvantage, especially in very high degrees. Consequently in the latter the glasses ought to be made to fit as closely to the eyes as possible, the eyelashes being cut short if necessary.

We have seen that in every case the far point and a point on the retina are conjugate foci. Hence an object situated at the far point of any eye will have a sharp image upon the retina (Fig. 33). This may be made clearer perhaps if we consider the myopic eye from another point of view. We have seen that the rays from a point on the retina meet in front of the eye at the far point. We may again use the principle of reversibility of rays. If there is a luminous point at the far point, the rays emitted from it which enter the eye will meet



FIG. 36.



FIG. 37.



FIG. 38.



FIG. 39.

FIGS. 36–39.—Effect of correcting lenses upon the size of the retinal image. In Figs. 36, 38, where the optical centre of the lens, O, coincides with the anterior focal point of the eye, F_1, the size of the retinal image is the same as in emmetropia. When the lens is closer to the eye than the anterior focal distance of the eye the size of the retinal image is diminished (convex lens, Fig. 37) or increased (concave lens, Fig. 39).

on the retina ; in other words, the image of an object at the far point will be upon the retina.

From these considerations we learn how it is that a patient with myopia cannot see clearly things which are a long distance away, whereas he can see things which are near. In common parlance, he is " short-sighted." He can see things at a distance better if he screws up his eyes. This is because he thus makes a narrow slit to look through, and this slit acts like the hole in the cardboard before a candle (*vide* p. 20). The term myopia originated in this peculiarity (μύειν, to shut : ὤψ, the eye or countenance).

The patient with hypermetropia, on the other hand, can see



Fig. 40.—Base of an Ionic column, showing torus. The small section to the right shows the radius of curvature in the vertical direction, the large section that in the horizontal.

neither distant nor near objects clearly with his eyes at rest, since the far point is virtual, and it is impossible to place an object at its situation. We shall see later that he is better off than the myope by virtue of accommodation.

We have already seen that the emmetrope sees only distant objects clearly with his eyes at rest, since the rays from such distant objects are nearly parallel. For practical purposes objects more than 6 metres (20 feet) away from the eye form clear images upon the retina.

The condition of an eye, whether emmetropic, hypermetropic, or myopic, is called its refraction, or more accurately its static refraction, since the term applies to the eye at rest.

We have hitherto considered only such errors of refraction

as are due to axial shortening or lengthening of the eye (*axial ametropia*). It is not difficult to understand that ametropia might be due to other causes. Thus, myopia might be due to the refractive power of the eye being too strong; in this case parallel rays would be brought to a focus in front of the retina even if this were in its normal position. Increase or decrease in the refractive power of the eye might be due to two causes. It might be due to alteration in the refractive indices of the media, or to alteration in the curvatures of the refractive surfaces: ametropia due to these causes is called *index* or *curvature ametropia* respectively. Both are much less common than axial ametropia. Index ametropia is very rare,



FIG. 41.—Sturm's conoid. V V, vertical meridian of refracting surface, more curved than H H, the horizontal meridian. A, B, C, D, E, F, G, sections of conoid. From B to F is the focal interval of Sturm. D shows the circle of least diffusion.

though we shall have a physiological example of it later (*vide* p. 49).

Curvature ametropia has a special importance, not because it gives rise to simple hypermetropia or myopia, but because it is the cause of another very troublesome error of refraction, called *astigmatism*. In most eyes, even if they are ametropic, the areas of the refractive surfaces uncovered by the pupil and used in vision are very nearly spherical. Sometimes, however, they are not. In most of these cases it is the cornea which is at fault, and the error is generally of such a nature that this surface is flatter from side to side than it is from above downwards. Even in these cases the curvatures in the vertical and horizontal meridians are both spherical, but the radius of curvature of the horizontal meridian is longer than that of the vertical. Perhaps the pressure of the lids on the globe tends to squeeze it above and below. A familiar example of such a surface, which is said to have a *toric* curvature, is seen

at the base of an ordinary Ionic column (Fig. 40). Here the centre of curvature of the horizontal section is the same as that of the column, while the centre of curvature of the vertical section is in a line with the surface of the upper part of the column. It is obvious that any other section of this surface than the vertical or horizontal will not be spherical but elliptical.

What will be the effect of such a toric cornea upon the refraction of the eye? Clearly the more curved meridian will have more refractive or convergent power than the less curved: hence if parallel rays fall upon the surface the vertical rays will come to a focus sooner than the horizontal. The rays after refraction will be perfectly symmetrical when referred to the vertical and horizontal planes. They will have two foci. The whole bundle of rays is called Sturm's conoid, and the distance between the two foci is called the focal interval of Sturm. It is difficult to represent this conoid on a plane surface (Fig. 41), but we can see what sections of the bundle or pencil of rays would look like at different distances from the refractive surface (Fig. 41, A—G).



Fig. 42.

At A the section will be a horizontal oval or oblate ellipse, because the vertical rays are converging more rapidly than the horizontal. At B the vertical rays have come to a focus, while the horizontal are still converging: the section will be a horizontal straight line. At C, D, and E the vertical rays are diverging and the horizontal are still converging. At one place in this focal interval there will be a spot (D) where the vertical rays have diverged from the axis exactly as much as the horizontal rays have converged towards it. Here the section is a circle, which is called the *circle of least diffusion*. At F the horizontal rays come to focus while the vertical are diverging: the section will be a vertical straight line. Beyond this point, as at G, both sets of rays are diverging, and the section will always be a vertical oval or prolate ellipse.

What will happen if the retina is situated at either of these points of section? In the first place it is obvious that the retinal image will always be blurred, and it is because the rays never come to a focus in a single point that the condition is called astigmatism (*ἀ*, privative, *στίγμα*, a point). If the

retina cuts the conoid at A, where none of the rays have come to a focus, every meridian will be in the same condition, though in different degree, as in the axial hypermetropic eye: this condition is therefore called *compound hypermetropic astigmatism*. If the retina is at B the vertical meridian will be in the condition of an emmetropic eye, while the horizontal will still be in the condition of a hypermetropic one: this condition is called *simple hypermetropic astigmatism*. At C, D, and E the vertical meridian will be in the condition of a myopic, and the horizontal still in that of a hypermetropic eye: this is called *mixed astigmatism*. At F the vertical meridian is still myopic, whilst the horizontal is in the same condition as in an emmetropic eye: this is *simple myopic astigmatism*. Beyond F, as at G, both meridians are in the condition of an axial myope, the rays having crossed in the vitreous: this is *compound myopic astigmatism*. All these positions of the retina are met with in actual practice, though there is often a combination of axial and curvature defects.



Fig. 43.

It will be readily seen that such a condition cannot be corrected by means of any spherical lens. We must obtain some means of affecting one set of rays more than the other. This means is found in cylindrical lenses.


Fig. 44.—Refraction of parallel rays through a plano-convex cylinder.

Suppose C D E F is a cylinder of glass (Fig. 42): AB is called the axis of the cylinder. If a slice is cut off the cylinder by a plane parallel to the axis, it would form a cylindrical lens. Fig. 43 gives representations of a convex and a concave cylinder. The direction *ab* is called the axis of the cylinder, since it is parallel to the axis of the original cylinder from which the slice may be supposed to have been taken. It is important here to warn the student not to confuse the axis of a spherical and the axis of a cylindrical lens, as they are totally different things. The axis of a cylinder has just been described:

the axis of a spherical lens is the line joining the centres of curvature of the two surfaces.

How will a cylindrical lens affect parallel rays falling upon its surface? In the direction of its axis it is simply a plane lamina with parallel sides, so that it will have no effect upon the rays. In the direction at right angles to its axis it is spherical on one side and plane on the other: it will therefore act exactly like a plano-convex or a plano-concave lens, *i.e.*, it will make the rays either converge or diverge. If a convex cylinder is held between a point of light and a screen, a position can be found for the screen such that a sharp bright line is thrown upon it (Fig. 45): this is the focal line of the cylinder.



FIG. 45.—Refraction of divergent rays from a point of light through a plano-convex cylinder.

It is to be noted that the line is in the direction of the axis of the cylinder. If another convex cylinder of the same strength were held with its axis at right angles to the first, it



FIG. 46.—Effect of accommodation. The dotted lines show the curvature of the anterior surface of the lens and the course of rays with the eye at rest (static refraction). The solid lines show the curvature of the anterior surface of the lens and the course of rays with active accommodation (dynamic refraction).

would obviously form a focal line perpendicular to the first focal line. If the two cylinders are put in contact with their axes at right angles, all the rays after refraction must pass through both lines. The only place where they can go through both lines is where the lines intersect. Hence we see that two cylindrical lenses of equal strength, placed in contact with their axes at right angles, act exactly like a convex spherical lens of the same strength as either of the cylinders.

When the cornea has its directions of greatest and least curvature at right angles to one another, the condition is called *regular astigmatism*. In the commonest form, as we

have said, the vertical meridian is the more curved, the horizontal the less: this condition is generally called regular astigmatism "according to the rule." Sometimes the reverse is found: this is said to be "against the rule." Not infrequently the axes are oblique. Often after ulceration, &c., the surface of the cornea is irregular. This causes the rays of light to be refracted irregularly, so that there is no symmetry at all about them and different groups form foci in various positions. This is called *irregular astigmatism:* it cannot be corrected, and can only occasionally be improved by glasses.

Accommodation.—We have to consider now how it is that a person with normal sight can not only see distant objects, but also near ones. If an object is situated near the eye, *e.g.*, at ordinary reading distance—about 22 cm. or 9 inches—the divergence of the rays which it emits cannot be neglected.



FIG. 47.—Diagram of Helmholtz' theory of accommodation.

Since the converging power of the refractive media of the emmetropic eye is only strong enough to make parallel rays come to a focus on the retina, it is obvious that divergent rays falling upon the cornea will not nearly have come to a focus (Fig. 46). They will indeed be made convergent, but only to such a degree that they would meet somewhere behind the retina. Now if we can make the converging power of the eye stronger, a point may be reached when it is just strong enough to bring them to a focus on the retina. This is what is done by accommodation, and the manner in which the converging power of the eye is increased is by making the crystalline lens stronger.

We have seen that the refractive power of a convex lens depends upon its refractive index and upon the curvature of its surfaces. In accommodation, it is the latter which undergoes change. The curvature of the surfaces of the lens at rest in the eye is approximately spherical, and the radius of curva-

ture of the anterior surface is 10 mm., while that of the posterior surface is 6 mm. In accommodation, the curvature of the posterior surface remains almost the same, but the anterior surface changes so that in strong accommodation its radius of curvature becomes 6 mm. The eye under these conditions, which are called its dynamic refraction, has a much increased converging effect upon the incident rays.

The mechanism by which this change in the curvature of the lens is brought about is as follows (Fig. 47). The lens substance is plastic, so that it tends to conform accurately to the shape of its capsule. As has been described (p. 10), the capsule is anchored to the ciliary body by the suspensory ligament. In the condition of rest, the fibres of the suspensory ligament are stretched. In the ciliary body is the ciliary muscle, which takes its origin from its anterior attachment to the sclerotic at the angle of the anterior chamber. When the muscle contracts, it pulls the posterior part of the ciliary body and the anterior part of the choroid forwards slightly. The effect upon the suspensory ligament is to slacken it, and with it the lens capsule. The posterior surface of the lens is fixed by the support of the jelly-like vitreous, so that the slackening of the capsule makes itself most felt in the anterior part, which becomes bowed forwards.

The generally accepted theory here described is that of von Helmholtz. According to Tscherning the ciliary muscle tightens the suspensory ligament, so that the peripheral parts of the anterior surface of the lens are flattened and the central or pupillary area is increased in curvature.

Our control over the ciliary muscle, though involuntary, is very delicate, so that all distances up to quite close to the eye can be accurately focussed. The nearest point at which small objects can be clearly distinguished is called the *near point*, or *punctum proximum*. It is most accurately determined by gradually bringing a fine thread closer to the eye until it is found impossible to see it perfectly clearly. In practice it is sufficient to use very fine print and to determine the point at which it first becomes impossible to read it. The distance can be measured by a centimetre or inch tape held close beside the eye. For ordinary clinical purposes it is usually sufficient to judge the distance. At this point accommodation is exerted to its maximum, the lens capsule is as slack as it is possible to make it, and an object could only be seen clearly at a nearer point by placing a convex lens in front of the eye.

It has been shown that the far point of the eye varies according to its static refraction, *i.e.*, according to whether it is emmetropic, hypermetropic, or myopic. The near point also varies with the static refraction, but it also varies with the age of the patient, the reason being that the lens becomes less plastic as age advances. We have stated that the lens is a mass of epithelium of which the central part is the oldest (*vide* p. 9). As the lens gets older the central cells become tougher and more compressed, thus forming a relatively hard nucleus. The nucleus is less plastic than the younger cortex, and as age advances more and more of the fibres become converted into nucleus. Consequently the lens tends less and less to respond to changes in tension of the capsule. Thus, a child of ten, the earliest age at which it is possible to obtain satisfactory measurements, is able to see a small object clearly when it is only 7 cm., or less than 3 inches, from the eye. A person of thirty years of age cannot see clearly at less than 14 cm., or about $5\frac{1}{2}$ inches, from the eye.

Now, we have pointed out that the refractive power of a lens in dioptres is the reciprocal of its focal distance measured in metres (*vide* p. 29). The same method is applied to measure the static and dynamic refractive powers of the eye. Thus, the static refractive power of a myopic eye whose far point is 1 metre in front of the eye is said to be 1 dioptre; this is usually expressed by saying that the eye has 1 D of myopia. Similarly, if a hypermetropic eye has its far point half a metre behind the eye it is said to have 2 D of hypermetropia. By this method the emmetropic eye, which has its far point at infinity, has no refractive power when it is at rest, since $\frac{1}{\infty} = 0$.

Applying the same method to the dynamic refractive power, the child of ten, whose near point is 7 cm. from his eye, has a refractive power of $\frac{100}{7} = 14$ D, and a man of thirty, whose near point is 14 cm. from his eye, has a refractive power of $\frac{100}{14} = 7$ D.

By this means we can obtain a general rule for indicating the amount or *amplitude of accommodation*, not only of emmetropic but also of hypermetropic or myopic eyes. This is given by the formula $A = P - R$, which states that the amplitude of accommodation is equal to the refractive power of

the eye when fully accommodated (*i.e.*, the reciprocal of the distance of the near point in metres) less the refractive power of the eye at rest (*i.e.*, the reciprocal of the distance of the far point in metres).

A few examples will make this clearer. Thus, the emmetropic child of ten has an amplitude of accommodation of $\frac{100}{7} - \frac{1}{\infty}$ $= 14 - 0 = 14$ D. What is the amplitude of accommodation of an emmetrope whose near point is 12·5 cm. from his eye? Here $A = \frac{1000}{125} - \frac{1}{\infty} = 8$ D. From statistics which have been collected we can deduce that this man was about twenty-six years old (*vide* p. 50). Now let us take a case of myopia, *e.g.*, a myope of 2 D whose near point is 8 cm. in front of his eye. His amplitude of accommodation will be $A = \frac{100}{8} - 2$ $= 10{\cdot}5$ D. What is the amplitude of accommodation of a hypermetrope of 3 D whose near point is 12·5 cm. from his eye? Here the far point is behind the eye and distances measured in this direction must have the opposite sign to those measured in front of the eye. Hence $A = \frac{1000}{125}$ $- (-3) = 8 + 3 = 11$ D.

The numbers given by these calculations for the amplitude of accommodation give the strength of the convex lens which would have to be placed in contact with the cornea in order that the near point might be brought to the required distance without using the accommodation. Several interesting facts come to light from the calculations. Thus a hypermetrope of 3 D has to exert 11 D of accommodation in order that he may see clearly at 12·5 cm., whilst an emmetrope has to exert only 8 D of accommodation to bring about the same result. We see, then, that the hypermetrope has to exert an amount of accommodation equivalent to the amount of his hypermetropia in order to focus parallel rays upon his retina, *i.e.*, he has to use this amount of accommodation in order to see distant objects clearly. Again, in the case of the myope of 2 D, his far point is half a metre, or 50 cm., from his eye; he can see clearly at that distance without accommodating, but he has to exert 10·5 D of accommodation in order that he may see clearly at 8 cm. from his eye. This patient, then, has to exert nearly as much accommodation to alter his points of clear

vision from 50 cm. to 8 cm., *i.e.*, through 42 cm., that a hypermetrope of 3 D has to employ in order to move his point of distinct vision from infinity up to 12·5 cm. We see, therefore, that the *range of accommodation*, *i.e.*, the distance between the far point and the near point, is not always the same for a given amplitude.

The effect of age upon the static and dynamic refraction is given in Fig. 49, which is the result of a large number of statistics and gives the average results. From this table we see that even the far point alters in advanced age. After about fifty the eye tends to become hypermetropic, so that at eighty it has about 2·5 D of hypermetropia. This has nothing to do with accommodation, and hence nothing to do with loss of plasticity in the lens. It is, however, due to changes going on in the lens, viz., an alteration in its refractive index so that it has a weaker converging power.

The refractive indices of the successive layers of the lens increase from the periphery towards the nucleus. The effect is twofold: it tends to correct aberration by increasing the convergence of the central rays, and the total refractive index of the whole lens is increased, being greater than the refractive index of the nucleus. For it may be looked upon as a central bi-convex lens encapsuled in two menisci (Fig. 48). The menisci act as concave lenses because the curvature of the nucleus is greater than that of the periphery of the lens. Hence they tend to counteract the effect of the central lens, but not so much as if their refractive index was the same. In old age the index of the peripheral layers increases, so that the total refractive index of the lens becomes less, and the eye becomes hypermetropic.

FIG. 48.

If we turn our attention to the curve of the near point we see that the amplitude of accommodation gradually diminishes throughout life. Now, we are accustomed to hold books for reading or work for sewing, &c., at about 10 inches, or 22 cm., from the eyes. In order to be able to see clearly at 25 cm. from the eye we have to exert $\frac{100}{25} = 4$ D of accommodation. If we look at the table we shall see that an emmetrope has only 4 D of accommodation left at about forty-one years of age. He will still be able to see clearly at 25 cm., but not closer. If he is about forty-six he will have only 3 D of accommodation left. He will not now be able to see clearly

JUN 28 1923

at 25 cm., but he will have to hold his book farther off, viz., at $\frac{100}{3} = 33$ cm. If he is still older he will have to hold the book yet farther off, and he will probably have to use very large print or he will not see clearly at all. This is the condition which is called *presbyopia* (πρέσβυς, old).

It will be seen that a patient never requires more than + 4 D to correct his presbyopia only, since that is the amount of accommodation required to place the far point of the



FIG. 49.—Chart of static and maximum dynamic refraction at various ages. (Donders.) Abscissæ, ages; ordinates, accommodation in dioptres.

resting emmetropic eye at reading distance. A convenient rule to remember is that a presbyope requires + 1 D for every five years after forty; *i.e.*, at forty-five, + 1 D; at fifty, + 2 D; at fifty-five, + 3 D; at sixty and later, + 4 D. The rule errs in giving rather too liberal a correction; a slightly smaller correction is often more comfortable.

It is a common error among students to think that presbyopia is a condition which commences at about forty-five years of age in emmetropes, and earlier in hypermetropes. Of course this is not so: the condition which has been increasing throughout life first becomes troublesome when the near point

of the eye has receded so far that it is beyond comfortable reading distance.

There are two other phenomena which occur with accommodation, one affecting the iris, the other the direction of the eyes. In order that we may see a near object we must look at it; hence in order that we may see it at the same time with both eyes they must each turn inwards or converge. The amount of convergence, like the amount of accommodation, depends upon the distance of the object. It is therefore easy to understand that there is a close relationship between accommodation and convergence. We shall have more to say upon this subject when we consider the various forms of squint.

When we accommodate for a near object the pupil becomes smaller, or contracts. Experiment has shown that this movement of the iris is associated with the accompanying act of convergence rather than with accommodation *per se*. It is probably of the nature of an associated movement, or, as I have termed it, synkinesis (σύν, with, κίνησις, movement).

Contraction of the pupil during accommodation is not for the purpose of diminishing aberration, since this is already diminished by the act of accommodation. It has the effect of compensating for the relative increase of light entering the eye from near objects, but is greater than is necessary to produce this result.

It is not uncommon for the refraction of the two eyes to be different: this condition is called *anisometropia* (ἄ, privative; ἴσος, equal; μέτρον, measure). It might be anticipated that this could be corrected to some degree or entirely by unequal accommodation in the two eyes. Thus, if one were emmetropic and the other hypermetropic both eyes would be able to distinguish distant objects clearly if the hypermetropic one alone accommodated the requisite amount to correct its hypermetropia. It has been conclusively proved, however, that this does not occur. When these cases are not corrected by the proper glasses clear vision is wholly uniocular.

Although astigmatism is chiefly due to faulty curvature of the cornea, in some cases there is also lenticular astigmatism. This is not generally due to unequal curvature of the surfaces, but to slight tilting of the lens, so that the incident rays fall upon it obliquely. If we look through a tilted glass lens at printed matter we shall see that the letters become distorted and elongated in one direction; this is a form of astigmatism. The astigmatism of the crystalline lens is generally of such a

nature that it tends to counteract the corneal astigmatism, though sometimes it adds to the effect. As in anisometropia, it might be thought that astigmatism could be corrected by accommodation. If, for instance, the ciliary muscle acted only at the sides and not at all above or below, the anterior surface of the lens would become more curved in the horizontal than in the vertical meridian. This would counteract the effect of the ordinary form of corneal astigmatism. It has been proved, however, that this also does not occur. When the ciliary muscle acts, it acts equally all round the circle ; and



Fig 50.—Spherical aberration.

when one ciliary muscle acts, the one in the opposite eye acts simultaneously and equally under ordinary conditions.

The Pupils.

The iris acts like the diaphragm of any ordinary optical system, such as a photographic camera or a microscope. In discussing the effects of spherical mirrors in reflecting, and of spherical surfaces in refracting the rays of light, we said that in each case they were all brought to a focus in a single point. This is really only an approximation, which is sufficiently accurate for rays close to the axis. In a convex spherical lens, for instance, parallel rays near the axis meet at the principal focus (*vide* p. 25). The rays farther away from the axis, however, are refracted too much, so that they cut the axis nearer the lens than the principal focus (Fig. 50). This causes a blurring of the edges of the image, which is said to be due to *spherical aberration*. The diaphragm cuts off these peripheral rays, and thus prevents the blurring. In the eye the surfaces are not even spherical near the periphery, and are often not so in the centre, so that much more aberration is liable to occur. The iris reduces the effects of the evil to a minimum.

There is also another form of aberration due to the imperfect

refraction at spherical surfaces. White light is made up of all the colours of the spectrum. The component rays are refracted differently, the violet most, the red least. Hence there is a tendency for the white light to be split up into its components, in which case the image will have a coloured edge. This phenomenon is called *chromatic aberration*.

When light enters the eye and falls upon the retina the pupil contracts. We have already seen that the pupils also contract when the optic axes converge in accommodation. On the other hand they dilate if the skin of any part of the body is pinched or any sensory nerve is stimulated to the extent of causing pain. These responses to stimuli of various kinds are very rapid and delicate, and are easily observed. When they are altered by disease the changes which occur afford very valuable information as to the condition of the nerve tracts involved. The tracts are rather complicated, but it is essential that they should be understood.

Under normal conditions, with equal illumination—a point too frequently neglected—the pupils are equal on the two sides. It is very rare to meet with unequal pupils in a normal person ; such cases do apparently occur, but every possible pathological cause must be eliminated before we admit that the condition is an idiosyncrasy.

On the other hand, the size of the pupils varies much in different people under the same conditions of illumination, &c. In old people it is smaller than in the young, sometimes to so great an extent that the pupils are almost " pin-point." They are often smaller in hypermetropes, and larger in myopes, than in emmetropes ; they are said to be smaller in blue eyes than in brown. The causes of these differences are conjectural, and need not detain us. The two facts of prime importance have been mentioned, viz., that there are two reflexes, that to light and that to sensory stimulation, which act in opposite directions. The normal size of the pupil may be looked upon as essentially the resultant of these two forces.

The motor innervation of the pupil is as follows (Fig. 51). The sphincter pupillæ is innervated by the third cranial nerve. The pupil-constrictor fibres start in the anterior part of the third nucleus in the floor of the aqueduct of Sylvius. They pass out of the mid-brain and run in the main trunk of the third nerve as far as the orbit. Here the fibres pass into the branch which supplies the inferior oblique muscle, leaving it by the short root of the ciliary ganglion. From the ciliary ganglion they pass by the short ciliary nerves to the eye,



FIG. 51.—Diagram of the efferent pupillary paths. Dotted lines, pupillo-constrictor: *III nuc.*, nucleus of third nerve; *c.g.*, ciliary ganglion; *s.c.*, short ciliary nerves. Solid lines, pupillo-dilatator: *Dil.*, hypothetical dilatator centre in the medulla; *c.c-sp.i.*, Budge's centrum cilio-spinale inferius; *D I, D II, D III*, first, second, and third dorsal nerves (see text); *r.c.*, ramus communicans; *Stellate g.*, stellate ganglion; *ann. V.*, annulus of Vieussens; *i.c.g.*, inferior cervical ganglion; *cerv. symp.*, cervical sympathetic; *s.c.g.*, superior cervical ganglion; *Gass. g.*, Gasserian ganglion; *V1, V2, V3*, first, second, and third divisions of the fifth nerve; *V nasal*, nasal branch of the ophthalmic (first) division of the fifth nerve; *l.c.*, long ciliary nerves

piercing the sclerotic around the optic nerve, being here in company with the short ciliary arteries (*vide* p. 11). The nerve fibres pass forwards in the choroid and ciliary body to the iris.

The dilatator pupillæ is supplied by the cervical sympathetic nerve. The dilatator tract probably commences in the mid-brain not far from the constrictor tract. It passes downwards through the medulla oblongata into the lateral columns of the cord. The fibres leave the cord by the ventral roots of the first three dorsal and probably the last two cervical nerves, enter the rami communicantes, and run to the first thoracic or stellate ganglion. From here they pass by the anterior limb of the annulus of Vieussens into the cervical sympathetic. In this nerve they run up the neck to the superior cervical ganglion, whence they pass with the carotid plexus into the skull. They run over the anterior part of the Gasserian ganglion and pass into the first or ophthalmic division of the fifth nerve, following the nasal branch, which they leave finally to enter the long ciliary nerves, thus avoiding the ciliary ganglion. The long ciliary nerves enter the eye on each side of the optic nerve, accompanying the long ciliary arteries. Like them, they run forwards between the choroid and sclerotic, enter the ciliary body and thus reach the iris.



FIG. 52.—Diagram of the afferent and efferent pupillary paths for light stimuli. Afferent paths from left sides of retinæ, thick solid lines; afferent paths from right sides of retinæ, thick dotted lines; efferent paths of left eye, thin solid lines; efferent paths of right eye, thin dotted lines. *Ir.*, iris; *R*, retina; *II.*, optic nerve; *o.t.r.*, optic tract; *Coll. Supr.*, colliculus superior or anterior corpus quadrigeminum; *IIIrd nuc.*, nucleus of third nerve; *e.g.b.*, external geniculate body; *c.g.*, ciliary ganglion; *s.c.n.*, short ciliary nerves.

These complicated paths will be seen at a glance in the accompanying diagrams.

We have now to consider the nervous mechanism of the reflexes. The light reflex is carried out entirely through the constrictor centre, so we may put the dilatator tract aside for the moment. The afferent fibres are contained in the optic nerve, arising from all parts of the retina (Fig. 52). It is unknown whether they belong to the large or the small fibres of the optic nerve. It is certain that they undergo partial decussation in the chiasma, like the visual fibres, and that they enter the optic tracts. It is also certain that, unlike the visual fibres, they do not enter the lateral geniculate body, but leave the tract to pass by an unknown path to the third nucleus.

The constrictor centre possesses "tone," *i.e.*, it is perpetually sending out impulses which keep the pupil slightly contracted. If light falls upon the retina of one eye its pupil contracts—*the direct reaction to light;* but the pupil of the opposite eye also contracts simultaneously—*the consensual reaction to light.* This consensual reaction should always be tested, since it gives useful information which cannot be obtained from the direct reaction. Thus, if there is a block on one optic nerve so that there is no direct reaction to light, but the consensual reaction from light thrown upon the other eye is unimpaired, we know that the block, whatever it may be, does not affect the efferent constrictor tract, *i.e.*, the trunk of the third nerve, the branch to the inferior oblique, and the short ciliary nerves are intact. The consensual reaction is probably carried out by means of fibres which unite the two constrictor centres in the third nucleus.

That the afferent pupil-constrictor fibres undergo partial decussation in the optic chiasma is proved by Wernicke's *hemianopic pupil reaction.* This reaction is pathognomonic of disease of one optic tract. It will be seen from Fig. 52 that such a lesion will cut off the afferent impulses from corresponding halves of each retina, *i.e.*, from the temporal half of one and the nasal half of the other. If light is thrown upon these parts of the retinæ the pupils do not contract, but if it is thrown on the other halves of the retinæ the pupils respond.

The sensory reflex is more complicated than the light reflex, for both the dilatator and the constrictor centres play a part in its production under normal conditions. It has been shown that sensory stimulation causes first a rapid dilatation of the pupil due to augmentation of the dilatator tone through the cervical sympathetic, and then a second dilatation, rapid in

onset but slow in disappearance, due to inhibition of the constrictor tone. There are other reflexes and synkineses, *e.g.*, emotional, which need not detain us.

Minute examination of the pupil when the intensity of the light entering the eye is altered, shows that the pupil contracts and then oscillates rapidly, finally settling down into a condition of contraction which is slightly less than the summit of the first wave. In its sudden response, the pupil as it were oversteps the mark, oversteps it again in the opposite direction, and so on. Two different types of exaggeration of this oscillation are met with under abnormal conditions. One is the condition in which the oscillations are very large and easily seen, and which are to a large extent independent of the light falling upon the eye. This is called *hippus ;* its origin is obscure, but it undoubtedly depends upon the rhythmic activity of the nervous centres, and is not a peripheral phenomenon. More important is the lack of sustained contraction under the continued influence of light. Here the pupil contracts sluggishly when the intensity of the light is increased, but while the light is still kept constant it slowly dilates, often with superposed sluggish oscillations. This is a pathological phenomenon dependent upon diminished conductivity in the afferent path of the light reflex, *i.e.*, usually in the optic nerve (*see* Retrobulbar Neuritis).

Drugs are so frequently employed in ophthalmic practice for the purpose of dilating or constricting the pupils or paralysing the accommodation, that it is important to know exactly how they act. Pupil-dilating drugs are called mydriatics ; pupil-constricting, miotics ; drugs which paralyse the ciliary muscle, cycloplegics. All drugs which dilate the pupil also paralyse the accommodation in greater or less degree ; many attempts have been made to discover a drug which will effect the former purpose without the latter, but without success. Similarly, all miotics stimulate the ciliary muscle to contract, so that the eye assumes a condition of partial or complete accommodation.

The vexed question whether these drugs act upon the respective muscle fibres or upon the nerve endings contained in them need not be discussed here. The strongest mydriatic which we possess is *atropin ;* it paralyses the sphincter iridis and ciliary muscle completely, and is said also to stimulate the dilatator iridis. It has so potent an action that it abolishes the tone of the ciliary muscle. Thus, an emmetropic eye placed fully under the influence of atropin becomes hyper-

metropic to the extent of about 1 D ; this must be taken into account in correcting errors of refraction. Atropin solution (*e.g.*, 1 per cent.) instilled into the conjunctival sac is absorbed through the cornea into the anterior chamber, where it acts locally upon the intrinsic muscles. It takes a considerable time to cause complete paralysis, hence it is usual to order it for use at home three times a day for at least three days. The effects do not pass off for about ten days. One drop of 0·5 per cent. atropin sulphate solution causes wide dilatation of the pupil in thirty to forty minutes, and complete paralysis of accommodation in about two hours ; the effects do not pass off entirely till from three to seven days. Duboisin, hyoscin or scopolamin, and daturin act similarly to atropin.

Homatropin acts more quickly than atropin, and the effects pass off more quickly. Its full effect is obtained by an oily solution (in Ol. Ricini, 1 per cent.) in three-quarters of an hour, especially if combined with cocain (2 per cent.), which acts chiefly by increasing the permeability of the cornea. The effects pass off completely in forty-eight hours, or much more quickly if a drop of eserin (1 per cent.) is instilled. The mixture of homatropin and cocain, which is commonly employed for estimating refraction, does not paralyse the intrinsic muscles so fully as atropin, the tone of the ciliary muscle not being abolished so thoroughly. Homatropin probably acts upon the iris through the sphincter only.

Cocain, besides its anæsthetic effect through the endings of the fifth nerve in the cornea, iris, &c., also stimulates the sympathetic nerve endings in the dilatator iridis. It does not paralyse the sphincter, so that the dilatation of the pupil is only moderate, and the pupil continues to react to light even after prolonged application. Cocain is a useful drug in confirming the diagnosis of paralysis of the sympathetic nerve : if this nerve is paralysed cocain fails to dilate the pupil. The effect is not due to degeneration of the nerve endings, as I have found that cocain fails to act very soon after section of the sympathetic in the neck in animals.

Eserin, or physostigmin, the most powerful miotic we possess, acts by stimulating the third nerve endings in the sphincter and in the ciliary muscle. It is therefore an antagonist of atropin, but it is unable to overcome the dilatation produced by atropin. On the other hand, eserin readily overcomes the dilatation produced by homatropin and cocain. These facts are of very great importance and must be carefully borne in mind. Comparably with cocain eserin fails to

produce constriction of the pupil after section of the third nerve.

Eserin, unlike the common mydriatics, causes some smarting and injection of the ciliary vessels when instilled into the conjunctival sac. What is more unpleasant is the "dragging" sensation in the eye which patients complain of when it is acting. It may be so irritating as to cause vomiting, but this only occurs in very sensitive persons or when the drug is pushed. Owing to these symptoms it should not be instilled more frequently nor in stronger doses than requisite to ensure the desired result. A 0·5 per cent. solution or one considerably weaker is often adequate.

Eserin begins to contract the pupil and cause spasm of accommodation in about five minutes; its maximum effect is reached in twenty to forty-five minutes. The effect on accommodation lasts only an hour or two, that on the pupil two to three days.

Pilocarpin acts similarly to eserin, but is less potent.

In irritative miosis, due to stimulation of the third nerve, light, accommodation, and eserin will cause greater constriction, atropin dilatation. In paralytic miosis, due to paralysis of the sympathetic, light, accommodation, and eserin will cause constriction, atropin little or no dilatation.

In spastic mydriasis light, accommodation and eserin will cause constriction. In paralytic mydriasis there is no reaction to light or accommodation, and eserin acts very feebly.

Visual Perceptions.

When light falls upon the retina it acts as a stimulus to a sensory nerve ending. As contact of the skin with a foreign substance causes the sensation of touch, so stimulation of the retina causes visual perceptions. The changes which go on as the result of a suitable stimulus in an ordinary tactile end organ, the physiological impulses in the afferent sensory nerves, and the psychological interpretation of these impulses in the brain which we call tactile sensation, are all relatively simple. In the visual nervous mechanism they are much more complex and highly differentiated.

We may first very briefly consider the changes which occur in the end organ itself. Light falling upon the retina causes at least four effects. (1) The pigment in the hexagonal cells of the retinal epithelium migrates from the bodies of the cells into the processes which lie between the rods and cones. (2) The

cones become shorter. Both these effects are slow ; both give more response to the violet end of the spectrum than to the red end ; both are consensual as well as direct—light on one retina causes the effect on both, though the other be kept in darkness. (3) The visual purple, a substance which is found only in the rods, is bleached, so that a sort of photograph or optogram of the luminous object is formed. (4) Changes of electrical potential are set up in the retina.

We are more concerned, however, with the sensations which result from stimulation of the retina with light. These are of three kinds, which are called the Light Sense, the Colour Sense, and the Sense of Form. Each of these may become disordered, so we must examine what they really mean.

The *Light Sense* is the faculty which permits us to perceive light, not only as such, but in its gradations of intensity. By utilising shadows cast by the retinal blood vessels upon the rods and cones it can be proved that the neuro-epithelium is the actual sentient layer of the retina (Purkinje's experiment). It is in this layer that the clear images of objects in the outer world are focussed. The rays stimulate the rods and cones and give us the sensation of light. Hence rays falling upon the optic disc give rise to no visual sensation, and this is therefore called the *blind spot* (Mariotte).

If the light which is falling upon the retina is gradually reduced in intensity there comes a point when it is no longer perceived : this is called the *light minimum*. It varies very greatly according to the amount of light which has been falling upon the retina before the observation is made. We are all aware that if we go from bright sunshine into a dimly lit room we cannot perceive the objects in the room until some time has elapsed : the eyes have to become " adapted " to the amount of illumination. Hence observations on the light minimum are only comparable one with the other when the eye is in the condition of dark adaptation. Since this involves keeping light from the eye for at least twenty to thirty minutes the investigation is tedious, and perhaps this is the chief reason why it is much neglected clinically. The light minimum for the fovea is considerably higher than for paracentral and peripheral parts of the retina, and retinal adaptation affects the macula relatively little.

Having ascertained the light minimum, if the light is gradually increased in intensity there are points at which we can clearly distinguish a difference in the amount of illumination. We can do this best if we have two illuminated areas of equal

size to compare, as is done in special instruments for the purpose, called photometers. We can then find out how much brighter one area must be in order that we may distinguish a difference in illumination. This is called the *light difference*. It is found to vary with the amount of illumination—the greater the amount of illumination the greater will have to be the difference in order that we may be conscious of any difference at all. Indeed, light difference tends to follow a



FIG. 53.—Acuity of form sense in different parts of the retina. (Dor.) *F*, fovea centralis. Abscissæ, degrees towards the periphery of the retina; ordinates, relative acuity, that at the fovea being unity.

well-known law which is approximately valid for all sensory impressions (Weber's Law).

Neither of these functions of the light sense is much used in ophthalmology. There is no doubt that the light minimum is increased in diseases which impair the precipient elements, viz., the rods and cones (*see* Retinitis pigmentosa). It is said that disease of the conducting elements, the nerve fibres, causes increase in the light difference.

The rods appear to be much more sensitive to low illumination than the cones, so that in the dusk we see with our rods (*scotopic vision*). Nocturnal animals, like the bat, have few or no cones.

The *Form Sense*, which is next in importance, is the faculty which enables us to perceive the shape of objects in the outer world. Here the cones play the predominant part, and where they are most massed together and most highly differentiated,

viz., at the macula, there the form sense is most acute. It falls off very rapidly towards the periphery, as is shown in Fig. 53, and it is noticeable that the curve agrees fairly well with the diminution in the number of cones. We are accustomed to speak of acuity in distinguishing the shapes of objects as *acuity of vision*, and we mean by that the greatest acuity which it is possible to obtain. The acuity of vision, therefore, applies to central vision, or the vision of objects whose images are formed at the fovea and its immediate neighbourhood, the macula lutea.

The form sense is not a purely retinal function, but in the perception of composite forms—such as letters—is largely psychological. The purely physiological elements which enter into the complex form sense are (1) the light sense, (2) the sense of position, (3) the sense of discrimination. The sense of position depends upon the light sense and upon the conditions of contrast between the object and its background. A physiological process, called spatial induction, causes a lowering of sensibility of the area surrounding a stimulated area, so that the demarcation between the two areas is increased. The sense of position is measured by the *minimum visibile*. The sense of discrimination is the power to distinguish two visible objects as separate, and is measured by the *minimum separabile*. Form sense is measured by the *minimum legible* or *cognoscibile*.

In determining the acuity of vision, we utilise the visual angle (*vide* p. 34). We naturally choose as our basis the *minimum visual angle*, *i.e.*, the angle which two luminous points must subtend at the nodal point of the eye in order that they may be perceived as separate and distinct. Now, in order that we may get separate impressions from two points close together on the retina it is necessary that two cones shall be stimulated, and that there shall be a cone between these two which is not stimulated. If we know the diameter of a cone we can calculate the minimum angle which must be subtended at the nodal point in order that these requirements may be fulfilled. This angle, as we have already seen, is equal to the angle subtended on the other side of the nodal point by the two luminous points. As a matter of fact these calculations agree fairly well with the results of observations. It might be thought that the observation was a very easy one, but there are several complications. It is found that there is a certain amount of spread of the stimulus from one cone to surrounding ones, due chiefly to natural irregular astigmatism of the refractive media, especially the lens. This causes bright

objects on a dark background to appear a little larger than they really are : the phenomenon is called irradiation. It is not altogether a disadvantage, for it helps us very much in seeing actual points of intense light, like the stars, millions of miles away. In clinical work it would be a disadvantage to have an unduly high standard of visual acuity, because nearly everybody would be abnormal according to the standard. It is found that a minimum visual angle of 1 minute (1′) or one-sixtieth of a degree gives a very good average, and it is upon this basis that test types are constructed (*vide* p. 125).

The *Colour Sense* is that faculty whereby we are enabled to distinguish different colours and different colour tones. The exact investigation of the colour sense is one of great complexity, for the different colours of the spectrum differ in luminosity, so that this disturbing factor has to be added to those dependent upon the physiological condition of the retina, *e.g.*, its state of adaptation, and so on.

Appreciation of colours occurs only with lights of moderate or high intensity and some degree of light adaptation of the retina (*photopic vision*). If a spectrum of low intensity is viewed with the dark adapted eye it appears as a grey band, differing in brightness in different parts (*scotopic vision*). The brightest part is at about 530 $\mu\mu$, corresponding to the green of the photopic spectrum. The brightest part of the photopic spectrum is at about 580 $\mu\mu$, near the sodium line in the yellow. Scotopic vision is probably a function of the rods, colour vision of the cones.

If three colours sufficiently far apart in the spectrum are chosen, all the other colours, though not in the same degree of saturation, can be formed by their combination in suitable proportions, and white light can also be formed in the same manner. Hence normal colour vision is called trichromatic. There are reasons for choosing red, green and blue for these " primary " colours. Now, we know that physiological impulses are in some sense a reflex or image of the physical stimuli which give rise to them, *e.g.*, sound waves cause physiological impulses, which are perceived as sound, and so on. If, therefore, we imagine three slightly different kinds of impulse set up by the stimulation of the retina by red, green and blue light respectively, their combinations in suitable proportions would enable us to perceive the whole gamut of the spectrum, including white light. This is the basis of the Young-Helmholtz theory of colour vision. According to Hering's theory, chemical changes in three different types of " visual substance " situated in the retina cause the sensations of colour (including

white and black). The three substances are white-black, red-green, and blue-yellow. If anabolic or building-up changes (Assimilation) are set up in these substances, the sensations of white, red and blue are caused respectively. If katabolic or breaking-down changes (Dissimilation) are set up, black, green, and yellow result.

Perhaps no subject affords a better field for conjectures than the theory of colour vision. It is sufficient here for us to emphasise the importance of the three primary colours, upon which stress is laid by the Young-Helmholtz theory, and the intimate relations which exist between red and green, blue and yellow, and white and black, which are the foundation of the Hering theory. It is necessary to realise so much because the colour sense is defective in a fairly large proportion of people. This congenital abnormality is called *colour blindness* (*q.v.*). In it the importance of the three primary colours and the intimate relation of the pairs of colours are forcibly brought out.

The three types of visual perception are not confined to the minute area of central or macular vision. All are present in greater or less degree in more peripheral parts of the retina. In disease the earliest and most delicate traces of failing function are often to be found in the peripheral parts, central vision remaining perfectly normal. Hence the great importance of knowing the normal limits of the light perceptive and the colour perceptive areas of the retina.

The *field of vision* is the projection of these percipient areas of the retina on the outer world. When we stand upon the seashore and look at, or " fix," a ship on the horizon it forms a retinal image at the fovea. We are accustomed, then, to consider that any object in the outer world which forms the image at the fovea is situated somewhere upon the *line of vision, i.e.*, the line passing through the fovea and the nodal point of the eye. The foveal image is " projected " outwards along this line. Whilst still fixing the ship we are conscious of seeing, less clearly, innumerable objects for miles around. From our knowledge of the refractive mechanism of the eye we know that these objects must form their retinal images upon peripheral parts of the retina. Regarded from the side of the eye the image upon any point of the peripheral part of the retina is " projected " outwards along the line joining the point with the nodal point. The field of vision, then, is the projection outwards of all the points upon the retina which can give rise to visual perceptions. We will postpone the consideration of its properties to a later stage (*vide* p. 133).

CHAPTER V.

THE NEUROLOGY OF VISION.

In the preceding chapter we have considered the process of vision up to the point at which the retinal receptive elements, the rods and cones, have become stimulated. As with other



Fig. 54.—Diagram of the base of the brain, showing superficial origin of the I, II, III, IV, V, and VI cranial nerves: *inf.*, infundibulum; *o.tr.*, optic tract; *ped.c.*, cerebral peduncle; *i.g.b.*, internal geniculate body; *e.g.b.*, external geniculate body; *pv.*, pulvinar of optic thalamus; *M.O.*, medulla oblongata.

sensory nerves, stimulation of the end organ causes the development of nervous impulses which travel up the afferent tracts of the central nervous system to the brain. The comparison of the afferent tracts of common sensation with those

of vision throws so much light upon the latter that it is worthy of a moment's consideration.



Fig. 55.



Fig. 56.

Figs. 55 and 56.—Fig. 55 represents a diagram of the neurons of the most direct path of common sensation. Fig. 56 represents the neurons of the afferent visual path. I, II, III, neurons of the first, second, and third orders respectively; *n.c.*, nucleus cuneatus; *n.g.*, nucleus gracilis; *s.c.q.*, superior corpus quadrigeminum; *n.lent.*, nucleus lenticularis; *o.th.*, optic thalamus. I in Fig. 56, rod and cone bipolars in the retina. *Ch.*, chiasma; *e.g.b.*, external geniculate body; *pv.*, pulvinar of optic thalamus.

The sensory impulse of common sensation, *e.g.*, in the leg, is carried by a nerve fibre along the sensory nerve and the dorsal spinal root to the cord: it travels up in the posterior columns

of the cord to the nucleus gracilis or the nucleus cuneatus as the case may be (Fig. 55). The whole of this course is along the processes of a single cell or neuron, which has been called the neuron of the first order. The impulse is taken up in the nucleus gracilis or cuneatus by a second cell, and is carried along the nucleo-thalamic tract or mesial fillet to the opposite



FIG. 57.—Diagram of the afferent visual paths from the retinæ, with the corresponding fields of vision. (After von Monakow.) *Ch.*, chiasma; *pv.*, pulvinar of optic thalamus; *e.g.b.*, external geniculate body; *s.c.q.*, superior corpus quadrigeminum.

optic thalamus; other fibres, especially those derived from the nucleus cuneatus, pass to the superior colliculus or corpus quadrigeminum. The cells in the nuclei gracilis and cuneatus are the neurons of the second order. A third cell, the neuron of the third order, situated in the thalamus or colliculus, carries on the impulse to the cortex cerebri. Here the nervous impulse is transformed into a psychic impulse, a change which is not and probably never can be understood.

Let us compare with this the visual afferent tracts (Fig. 56).

The end organ is the neural epithelium of rods and cones. The first true conducting nerve cell or neuron of the first order is the bipolar cell of the inner nuclear layer with its axon in the inner reticular layer. This microscopic cell corresponds morphologically with a dorsal root ganglion cell and its long processes stretching, in some cases, from the tip of the toe to the top of the spinal cord. The neurons of the second order are the ganglion cells in the retina whose processes pass into the nerve fibre layer and along the optic nerve to one of three terminations : most (80 per cent.) end in the lateral or external



Fig. 58.—Distribution of the papillo-macular fibres in the optic nerve. The shaded areas in A′, B′, C′, represent the papillo-macular bundle in sections corresponding with A, B, and C.

geniculate body, others pass to the optic thalamus, and a few go to the superior colliculus. At these three sites a new cell, the neuron of the third order, takes up the transmission of the impulse, travelling by way of the optic radiations of Gratiolet to the cortex of the occipital lobe, which is the so-called *visual centre.*

We see, then, the morphological identity of the two systems, in spite of the great anatomical differences which specialisation has brought about. We may emphasise again the fact that the peripheral optic nerve proper is a bipolar cell in the inner nuclear and inner reticular layers of the retina, while the so-called optic nerve is a part of the central nervous

system homologous with the posterior columns of the spinal cord.

We must now investigate more minutely the individual parts of the visual system. The results which are about to be described are derived from three sources—embryology, experiments upon animals, and clinical observation as controlled by post-mortem findings.

Let us first trace the fibres from the various parts of the retina (Fig. 57). In general it may be said that the fibres from peripheral parts enter the middle of the nerve, while the fibres from parts near the nerve enter the peripheral parts of the nerve: they probably maintain this relative position as far back as the chiasma. There is, however, one disturbing factor, viz., the fibres from the macular region. This part is specially well supplied. The fibres pass into the outer part of the nerve, where they are spread over an area which is triangular in section, with the apex towards the centre of the nerve (Fig. 58). These *papillo-macular fibres* soon become more centrally situated, so that in the posterior part of the nerve they are all in the centre. Tracing them still farther backwards some decussate in the chiasma, while others enter the optic tract of the same side. They spread out widely in the two tracts and in the lateral geniculate bodies. The axons of their corresponding neurons of the third order are also widely distributed in the optic radiations and end in the lips of the calcarine fissure of the occipital lobe, probably along its whole length. We see then that each macular region is represented in each occipital lobe, so that no lesion confined to one occipital lobe will abolish central vision in either retina.

According to Gordon Holmes and Lister, from observations on patients with gunshot wounds of the occipital lobes, the maculæ are represented at the most posterior part of the visual cortex, probably on the margins and on the lateral surfaces of the occipital poles; each macula is represented only in the opposite occipital lobe (Figs. 61, 62).

The fibres from peripheral regions of the retina, on the other hand, form two distinct groups, corresponding with the temporal and nasal halves of the retina. The limitation is very exact, as if a vertical line divided the retina into two halves at the level of the fovea; since the foveal fibres have a different distribution the line makes a little curve round it so as to avoid it (Fig. 57). The fibres from the temporal half of the retina

enter the chiasma but do not decussate; they pass into the optic tract of the same side. Thence they run to the lateral geniculate body, and a few to the thalamus and superior colliculus. The fibres from the nasal half of the retina enter the chiasma, decussate, and pass into the optic tract of the opposite side having a similar distribution on the other side. The corresponding neurons of the third order pass by the optic radia-



FIG. 59.—A diagram to represent the distribution of the different cortical areas in a coronal section through the left hemisphere one centimetre behind the fossa parieto-occipitalis. *LV.*, posterior cornu of lateral ventricle; *s.calc.*, posterior calcarine sulcus; *s.lim.sup.ar.str.*, sulcus cunei (sulcus limitans superior areæ striatæ); *s.lim.inf.ar.str.*, sulcus lingualis (sulcus limitans inferior areæ striatæ). Note how the stripe of Gennari (the thick black line) ends abruptly at the bottom of the sulcus cunei and the sulcus lingualis. (After Elliot Smith.)

tions to the corresponding occipital lobes. We see, therefore, that a lesion of one occipital lobe or optic tract will cause blindness of the temporal half of the retina on the same side and of the nasal half of the retina on the opposite side. Projecting this outwards, such a lesion will cause loss of vision in the opposite half of the binocular field of vision, a condition which is known as *hemianopia* (hemiopia, hemianopsia). We may recall the fact that the afferent pupil-constrictor fibres have a similar semi-decussation in the chiasma.

The visual fibres in the optic radiations maintain a posterior position in common with the sensory fibres from the rest of the body, being thus situated behind the motor fibres in the internal capsule. They are close to the posterior cornu of the lateral ventricle, so that they are liable to pressure here when the ventricle is distended.



FIG. 60.—A diagram to illustrate the distribution of the cortical areas on the mesial surface of the occipital region of the left hemisphere. The two arrows indicate the plane of the section shown in Fig. 59. The area striata is represented by dots; the area parastriata by crosses; the area peristriata by circles. (After Elliot Smith.) Compare the above with Gordon Holmes's diagram (Figs. 61, 62), arrived at on clinical grounds.

The occipital cortex in and about the calcarine fissure differs from the cortex elsewhere in the possession of a white line, the line of Gennari, interpolated in the grey matter. This area, which is the primary visual or visuo-sensory area (Figs. 59–62), is the cortical projection of the corresponding halves of both retinæ. In this projection the part above the calcarine fissure represents the upper corresponding quadrants, the part below the lower corresponding quadrants of both retinæ.

It may be mentioned that there are efferent fibres in the optic nerves. They are axons of cells in the lateral geniculate body. Their function is unknown, but they are probably



Fig. 61.—A diagram of the probable representation of the different portions of the visual fields in the calcarine cortex. Fig. 61 is a drawing of the mesial surface of the left occipital lobe with the lips of the calcarine fissure separated so that its walls and floor are visible. The markings of the various portions of the visual cortex which is thus exposed correspond with those shown on the chart (Fig. 62) of the right half of the field of vision. (Gordon Holmes.)

concerned in the movements of the pigment epithelium and of the cones; possibly also in the chemical changes in the visual purple.



FIG. 62.

SECTION II.

THE EXAMINATION OF THE EYE.

CHAPTER VI.

EXTERNAL EXAMINATION.

Ophthalmic patients may be roughly divided into two groups: those who present manifest objective signs of disease, and those who, presenting no outward and visible signs, have abnormal subjective symptoms. The division is convenient both in theory and in practice. The second group, in all but "functional" cases, have latent objective signs which it is our duty to discover. In the first group the manifest signs may mask even more serious conditions which it is our duty to bring to light. Not infrequently we are confronted with obscure cases which demand the most careful systematic examination in order that nothing may be overlooked. Though it is not always possible, or even necessary, to go through the lengthy routine of an exhaustive systematic examination, yet the details of such a routine must be firmly engrafted in our minds, ready for instant application, if we wish to avoid mistakes in diagnosis.

We shall first describe the methods of examination of the parts which can be observed with the unaided eye, though we shall often ensure accuracy by artificial assistance. Next, we shall describe the methods which must be employed to examine the deeper parts of the eye. Finally, we shall map out a routine of systematic examination.

In the patients who belong to our first group we shall at once be confronted with visible signs of disease. We shall not consider in this place gross signs such as marked protrusion or

proptosis, great deviations of the optic axes from the normal parallelism, and so on. These will be more conveniently investigated in their special relations. We shall confine ourselves to conditions affecting the globe itself.

The Conjunctiva.—In the normal position of the lids only that part of the bulbar conjunctiva which is exposed in the palpebral aperture, together with parts of the intermarginal strip along the edges of the lids, is visible. In order thoroughly to investigate the whole conjunctival sac it is necessary to expose the palpebral conjunctiva and the fornices.

The lower fornix is easily exposed by drawing down the lower lid while the patient looks towards the ceiling.

The upper palpebral conjunctiva is exposed by everting the upper lid.

Eversion of the upper lid requires some practice. (1) The best, and often the easiest, method is as follows : Stand facing the patient. Place the right index finger horizontally along the patient's left upper lid while he looks towards his feet. Draw the skin of the lid outwards : this causes the inner part of the edge of the lid to come forwards, while at the same time the pressure of the finger affords a fixed point around which the lid can revolve in a vertical direction. Insinuate the right thumb under the projecting edge of the lid, and roll the lid upwards towards the index finger. The right lid is everted in the same manner, using the left hand.

This method is very easy when the eyes are prominent, and it causes a minimum of discomfort to the patient. When the eyes are deeply set in the orbit, as is often the case in old people whose orbital fat has become to a great extent absorbed, more pressure is needed and a little pain is caused. In such cases the following method may be adopted ; the tyro will generally find it easier :

(2) Place a probe or thin pencil horizontally along the skin of the upper lid at the level of the upper border of the tarsus, the patient looking towards his feet. Seize the eyelashes between the left index and thumb, and draw the lid away from the globe, using the probe as a fixed point. Rotate the lid in a vertical direction round the probe, which is then withdrawn.

In many cases we wish to evert the upper lid when standing behind the patient, who may be lying on a couch. In this case the following is the best method :

(3) Place the left index finger vertically upon the lid while the patient is looking towards his feet. Seize the lashes with

the right index and thumb, and rotate the lid around the tip of the left index.

In babies a special arrangement of the patient facilitates thorough examination of the conjunctival sac and eye :

(4) The surgeon sits facing a nurse, who holds the child on her lap. The baby's head is placed between the surgeon's knees ; its body is on the nurse's lap. She holds the child's hands against its body, thus keeping them out of the way, and at the same time steadying the child. If, as is often the case, there is blepharospasm, eversion of the lids is extremely easy : indeed, it becomes troublesome when we wish to examine the cornea. Here the spasm of the orbicularis fixes the lids against the globe, and the slightest attempt to draw the lids apart causes both to become everted. When this does not occur, method (3) must be adopted.

Having everted the upper lid we can examine the palpebral conjunctiva, but we are still unable to see the upper fornix. This can usually be effected in adults by the following manœuvre : (1) With the lid still everted by the first method it is fixed in that position by the left thumb placed upon its margin at about the middle. The right thumb or finger is placed in the middle of the lower lid. Firm, steady pressure is then made through the lower lid upon the globe in a direction straight backwards, as if to push the globe into the orbit. In the meantime firm pressure is also exerted backwards upon the upper lid with the left thumb. The fornix will generally start forwards suddenly, but only if the patient keeps looking well down towards his feet all the time.

This method, though unpleasant, is not painful. The only other method of exploring the upper fornix is more effectual, but also painful. The eye should therefore be well cocained.

(2) The upper lid is everted in the usual manner. A retractor is then inserted under the everted lid into the fornix. The margin of the lid being fixed as in (1), the lid is everted a second time, so that the fornix is fully exposed. Sometimes it is necessary to grasp the everted lid with forceps and thus evert it a second time.

By these manœuvres the conjunctival sac can be thoroughly explored. Special attention must be paid to the favourite sites for foreign bodies and manifestations of disease, *e.g.*, foreign bodies often lodge on the palpebral conjunctiva about 2 mm. from the margin at about the middle of the lid ; trachoma follicles are most marked in the upper fornix ; scarring from old trachoma is most marked in the palpebral conjunctiva, &c.

Adhesions between palpebral and ocular conjunctiva and obliteration of either fornix cannot fail to be noticed. Inflammatory conditions can be thoroughly examined; accurate diagnosis often depends upon minute investigation. Patches of granulation tissue which bleed easily may mark the site of embedded foreign bodies, or, if on the palpebral conjunctiva, the site of a chalazion.

The ocular conjunctiva can be almost completely examined without everting the lids if the eye is moved up and down while the lids are kept apart. The redness which is observed in irritative and inflammatory conditions varies in its distri-



FIG. 63.—Conjunctival congestion—engorgement of the posterior conjunctival arteries and veins. (After Guthrie.)



FIG. 64.—Congestion of the perforating branches of the anterior ciliary arteries. (Dalrymple.) The dusky spots at the seats of perforation are often seen in dark-complexioned persons.

bution and nature according to the cause. Here we must remember the groups of vessels with which we have to deal (*vide* p. 11). Three groups may be distinguished, though most of them are too small to be recognised in health: (1) the proper vessels of the conjunctiva or posterior conjunctival vessels (Plate II.); (2) the anterior conjunctival vessels, which supply the area adjacent to the limbus or corneal margin and send minute loops into the edge of the cornea itself (*vide* p. 13, Fig. 63); (3) the anterior ciliary vessels, lying in the subconjunctival or episcleral tissue. In the last group the perforating branches of the arteries are seen in health as several comparatively large tortuous vessels which suddenly cease about 2 to 4 mm. from the corneal margin. They have very numerous small episcleral branches which are invisible in health, but

when dilated form a pink zone of fine, straight, very closely set vessels around the cornea. The corresponding perforating veins are very small, but more numerous than the arteries; their episcleral branches form a closely-meshed network.

Congestion of the individual groups of vessels affords important evidence as to the seat of the mischief. The conjunctival vessels can be distinguished from the anterior ciliary by the following points: (1) they are a brighter brick-red, the ciliary vessels being seen through the conjunctiva, which imparts a purple tinge; (2) if the conjunctiva is moved to and fro over the sclerotic by the finger placed on the lower lid, the conjunctival vessels also move while the ciliary remain stationary; (3) the individual vessels and the network they form can be seen in the conjunctival system, whereas the ciliary



Fig. 65.—Ciliary congestion—engorgement of episcleral twigs of the anterior ciliary arteries. (After Dalrymple.)



Fig. 66.—Congestion of the anterior ciliary veins, episcleral venous plexus. (After Dalrymple.)

form for the most part a diffuse reddish-violet blush in which the separate vessels are indistinguishable; (4) if the blood is pressed out of the vessels the ciliary fill up at once on removing the pressure, since they anastomose very freely, while the conjunctival fill slowly.

In general, congestion of the conjunctival vessels, leaving a relatively white zone around the cornea, accompanied by mucous or muco-purulent secretion, is indicative of conjunctivitis. If there is much irritation and so-called photophobia, with some blepharospasm and very watery—lacrymal—secretion, we suspect the presence of a foreign body on the cornea or under the lid, so that it rubs against the cornea: the condition may be due to misplaced lashes (trichiasis). Phlyctenular ophthalmia may produce a similar picture. In such a case there is also usually congestion of the anterior conjunctival vessels. Careful examination shows that the vessels in

the circumcorneal zone are bright red, and that the corneal loops are also dilated and visible. Any irritation of the cornea—ulcers, abrasions, &c.—causes this conjunctival congestion of the circumcorneal zone and corneal vessels. Though conjunctival they do not move with the membrane. A definite leash of dilated vessels confined to the conjunctiva or encroaching upon the cornea is usually indicative of phlyctenular disease.

Pink circumcorneal congestion is also met with in inflammation of the iris. Here the anterior ciliary vessels are also involved.

Circumcorneal congestion of a peculiar lilac tint, more deeply seated and often patchy, is associated with cyclitis and deep scleritis. This is the condition which is known as ciliary congestion *par excellence*.

Dusky congestion at the limbus, composed of a fine reticulum —the episcleral venous plexus—often points to glaucoma, but may accompany other diseases, especially in old people.

These conditions run into one another very frequently, so that they then cease to have special diagnostic importance.

Lacrymal Apparatus.—Conjunctival congestion of one eye only, or signs of irritation such as watering, should lead us to suspect the efficiency of the lacrymal apparatus. Simple epiphora or flow of tears on to the cheek may be due to blocking of one or both puncta or to their malposition, or to blockage elsewhere in the canaliculi or nasal duct. Displacement of the lower punctum may be easily overlooked. The puncta are not visible normally without slightly everting the lids. Displacement is often very slight, due to slackness of the lid causing a little rolling out or ectropion, especially in old people. In spasm of the orbicularis the lid may be rolled in too much (entropion); this may also cause epiphora. Sometimes with more conjunctival inflammation, but often without, there is distension and chronic inflammation of the lacrymal sac. In all such cases we carefully observe the exact position of the puncta, whether they are in apposition to the bulbar conjunctiva as they should be, and also whether there is any regurgitation from the lacrymal sac when it is pressed upon. The lacrymal sac is situated in the lacrymal fossa between the inner canthus and the nose: the fundus of the sac extends slightly above the level of the inner palpebral ligament, which is on a line with the canthus. Pressure inwards and backwards in this position will press upon the sac. If there is any obstruction to the flow of the contents into the nose by the

nasal duct, as is usually the case when the sac is inflamed, the contents will generally regurgitate into the conjunctival sac by way of the canaliculi, and will be seen pouring from the puncta. We note whether the contents are tears, mucus, or muco-pus.

Without describing special methods of ascertaining if the lacrymal passages are patent (Chap. XXXII.), a simple test may be mentioned. A drop of fluorescein solution is placed in the conjunctival sac and the patient is told to blow his nose. If the passages are free, the handkerchief will be stained with the solution.

The Sclerotic.—Inspection of the sclerotic around the cornea may reveal raised nodules of episcleritis with their localised areas of vascular congestion (*vide* p. 236). Deep scleritis may be shown by dusky ciliary congestion and opacification of the deeper layers of the cornea at the periphery (sclerosing keratitis, *vide* p. 238).

Definite blue coloration of the circumcorneal sclerotic, except in young children, is pathological. It is most frequently due to ciliary staphyloma, a giving way of the sclera owing to inherent weakness (injury, scleritis, &c.) or to increased intraocular pressure (glaucoma). Discoloration may be due to pigmentation. Slight duskiness around the spots where the anterior ciliary vessels perforate is not uncommon in people with dark complexions. Otherwise pigmentation in this neighbourhood, either in the conjunctiva or sclerotic, should be regarded with suspicion as indicative of melanotic sarcoma. Definite nodules of deeply pigmented tissue in the situation of the perforating vessels is very significant of sarcoma of the ciliary body.

If there is bulging of the sclerotic an attempt should be made to transilluminate it in the dark-room. If it is a true ectasia (staphyloma) light will pass through into the eye. The light should be concentrated upon the spot by a strong lens (*vide* p. 82).

The abrupt or very gradual curvature of the sclerotic as it passes back from the cornea may indicate high hypermetropia or myopia respectively.

The Cornea.—A little experience will enable us to recognise at a glance if the cornea is smaller than usual. A small cornea with a shallow anterior chamber is very suggestive of glaucoma.

The cornea should be bright and transparent. We first examine its surface. Any loss of substance, such as an abra-

sion, may easily be overlooked without special care. The condition of irritation and lacrymation will often put us on the track, but these features may be so slight as to pass unnoticed. The following methods should be adopted :

(1) Place the patient facing the window. Stand in front and direct the patient to follow the index finger, which is held horizontal and moved slowly up and down. The finger is then



FIG. 67.—Focal or oblique illumination. The diagram shows how the focus of the light may be altered by slightly shifting the position of the concentrating lens.

held vertical and moved from side to side. While these manœuvres are being carried out the image of the window, or corneal reflex as it is called, is carefully watched. If the surface of the cornea is normal there will be no distortion of the reflex as it passes over it. If there is an abrasion the image will be distorted here, and will be less clearly defined. This method should always be resorted to first, as it is good practice in observation and needs no artificial aid, which may not be available under some circumstances.

(2) If the first method gives uncertain results or fails, instil

a drop of fluorescein solution (2 per cent.). This is best done by telling the patient to look towards his feet; the conjunctiva above the cornea is then lightly touched with the tip of a glass rod which has been dipped in the solution. A pad of cotton wool is immediately placed upon the closed lids so as to mop up the excess and the tears which tend to flow over the face. If this detail is not attended to the face becomes stained unnecessarily. It is a good plan to wash out the excess of fluorescein with a drop of cocain solution, but it is not essential. Any spot on the surface of the cornea which is denuded of epithelium will appear green.

A bundle of dilated conjunctival vessels near the limbus will often mark the site of a foreign body upon the cornea.

Opacities of the cornea may be so faint that they require very minute investigation, and the same is true of the details —depth, &c.—of gross opacities. We can study them best by focal or oblique illumination.

Focal or oblique illumination is carried out as follows (Fig. 67): the patient is placed, preferably in the dark room, with a light about two feet in front but slightly to one side. The light is concentrated upon the cornea by a strong convex lens (the large lens in the ophthalmoscope case). The rays of light are brought to a focus by the lens. The cornea or other superficial structures can thus be examined under the intense light of the converging rays. The position of the minute image of the light formed by the lens can be moved over the surface of the cornea by slight lateral movements of the lens without altering the position of the light. Similarly the light may be focussed upon the iris or crystalline lens by moving the lens slightly nearer to the eye.

Having thus brilliantly illuminated the part of the cornea which we wish to investigate, we may magnify the spot by looking through a very strong convex lens or corneal loupe held in the other hand. The management of the two lenses requires a little practice, but is easily mastered. A few words of explanation about the corneal loupe will help us to employ it to best advantage.

When we magnify a small object with a strong convex lens we place it within the focal distance of the lens and view it through the lens. We know that under these conditions the lens forms an enlarged image upon the same side as the object, but farther away (*vide* p. 30, Fig. 21). In order that we may see the image to best advantage we must see as much of it as possible, and we must see it under the largest possible visual

angle. The first requisite demands that the observer's eye shall be as close to the lens as possible. The second requisite depends upon the relative distances of the object and the eye from the lens; in practice these are found by slight movements of the lens.

In employing focal illumination, then, first focus the light upon the required spot. Then place the corneal loupe near the spot and look through it. Slowly advance the loupe towards the cornea until the spot comes into focus. Then get one's eye as close to the loupe as possible. In this manner we shall see all there is to be seen.

By moving the light and the loupe slowly over the whole surface of the cornea we can thoroughly explore it. By advancing the convex lens we can illuminate successively the back of the cornea, the iris and anterior part of the lens, and finally the deeper parts of the lens. By simultaneously advancing the position of the loupe towards the cornea we can successively bring these structures into accurate focus and examine them under considerable magnification. We cannot get beyond the back of the lens with a high power loupe, as it works at too short a focal distance. Moreover, in order to examine the deeper parts of the lens we must have the light almost in front of the patient, otherwise they will not be illuminated.

With a binocular loupe a stereoscopic effect is obtained, and the depth of opacities can be determined with great accuracy, but the degree of magnification is less. Special methods of examination with the corneal microscope, Gullstrand's slit lamp, contact illumination, &c., are useful in difficult cases.

Focal illumination without the assistance of a loupe or a dark room is often of great advantage through the good illumination which it affords. One soon gets into the habit of concentrating the light from the window upon the eye with the convex lens with a view to improving the optical conditions.

The cornea is often affected secondarily to the conjunctiva, as in phlyctenular ophthalmia. In such conditions the eye is most irritable and resistant to examination in bright light. The slightest attempt to separate the lids is accompanied by violent blepharospasm, especially in children. Yet it is in such cases that it is of the greatest importance to know and watch the condition of the cornea. For aught we know it may be ulcerated, and may even be upon the point of becoming perforated. Any roughness, or even an amount of pressure

which is quite justifiable in other cases, may suffice to cause the perforation which it should be our chief effort to avoid.

Fig. 68. — Bent wire lid retractor.

In such a case the lids must be separated by retractors. We may use simple bent wire retractors (Fig. 68) or Desmarres' retractor (Fig. 69). In babies the position recommended in method (4), p. 76, is employed: older children are placed upon a couch. The retractors are inserted gently into the palpebral aperture, first the upper, then the lower, the curved ends being insinuated between the lids and the globe. Traction is then made upon the retractors, not only in opposite directions up and down, but also away from the globe, so that the lids are lifted off the globe at the same time that they are separated. In these cases the globe tends to roll forcibly upwards when light falls upon it, which makes it very difficult to see the cornea even with the use of retractors. The difficulty may be overcome by pressing the end of the lower retractor well into the lower fornix, which drags the eye downwards by pulling on the ocular conjunctiva.

Fig. 69. — Desmarres' lid retractor.

When we have satisfied ourselves that there is little or no fear of perforation we may dispense with retractors. It has been pointed out that in children an attempt to separate the lids when there is much blepharospasm is usually followed by eversion of both lids, the cornea still remaining hidden. The way to overcome this difficulty is to place the two thumbs close to the edges of the lids and to press gently but firmly upon the globe as the lids are drawn apart. In this manner they are separated without becoming everted, but we must be extremely careful not to exert undue pressure and not to touch the cornea with the thumb nails.

In many diseases new vessels are formed in the cornea. An exact knowledge of their position, whether superficial or deep,



FIG. 70.—Diagrams of superficial corneal vessels. A, general vascularisation; B, a single leash of vessels, as in fascicular ulcer (*vide* p. 210); C, localised vascularisation, as in trachomatous pannus (*vide* p. 170).

and of their distribution, whether localised, general, peripheral, above, and so on, will often settle a disputed point in diagnosis.

Superficial vessels (Fig. 70) in the cornea are distinguished from deep (Fig. 71) by the following features: (1) superficial vessels can be traced over the limbus into the conjunctiva, while deep ones seem to come to an abrupt end at the limbus; (2) superficial vessels are bright red and well defined, while deep ones are ill defined, greyish red, or cause only a diffuse red blush; (3) superficial vessels branch in an arborescent fashion, dichotomously, while deep ones run more or less parallel to each other in a general radial direction, and branch at very acute angles, like a besom; (4) superficial vessels may raise the epithelium over them so that the surface of the cornea is uneven, while with deep ones the cornea, though hazy, is smooth. The peculiar course of deep vessels is probably determined by the lamellar structure of the substantia propria.

Fig. 71.—Diagram of deep corneal vessels, as in interstitial keratitis (*vide* p. 222).

The sensibility of the cornea may be tested by touching it in various spots with a wisp of cotton-wool twisted to a fine point and comparing the effect with the opposite side. Normally there is a brisk reflex closure of the lids. The sensibility is often diminished in corneal affections, but the change may be of some diagnostic significance in certain cases, *e.g.*, herpes (*vide* p. 220).

The Anterior Chamber.—The anterior chamber is shallow in extreme youth and in old age: at other periods of life it is about 2·5 mm. deep normally. It must be remembered that we estimate the depth of the anterior chamber by the position of the iris, and that we view the iris through the cornea, which is a strongly refracting convex surface. The effect of this is to magnify the iris and pupil, and to make it appear farther forwards than it really is. The same applies to anything in the anterior chamber, *e.g.*, the point of a knife in operations.

Good binocular vision enables us to estimate the depth of the anterior chamber when we are looking at it from in front.

The observation should be confirmed by taking a profile view.

The anterior chamber is abnormally shallow in glaucoma. It is often abnormally deep in irido-cyclitis. It is frequently unequal in depth in different parts. For example, it may be deeper at the periphery than in the centre in irido-cyclitis ; on the other hand, when the iris is bowed forwards (iris bombé) it is funnel-shaped, the centre being deep, the periphery very shallow.

After considering the depth, attention must be paid to the contents. In some wounds and ulcers of the cornea, and rarely without them, there is pus in the anterior chamber. It forms a layer at the bottom, the surface of the pus being level (hypopyon). A similar layer of blood may occur after contusions or spontaneously (hyphæma). The aqueous may be hazy, a condition not always easy to distinguish from haziness of the cornea. Such cases lead us to examine very carefully the back of the cornea with the loupe under focal illumination to see if there are any precipitates ("keratitis punctata") upon it, or we may see flocculent specks in the aqueous. All these conditions are of great diagnostic and prognostic importance.

The Iris.—We pay attention first to the *colour* of the iris and the clearness of its pattern. The two irides or parts of the same iris may be of different colour, both conditions being known as heterochromia iridis. A grey iris, with ill-defined pattern, suggests atrophy from cyclitis, glaucoma, &c. Dark brown spots in the iris, not raised above the surface, are common. Care must be taken to distinguish them from small nodules of the same colour or white (sarcoma, tubercle, gumma). "Muddiness" of the iris is the expression used for indistinctness of the pattern, caused by inflammatory exudates. A muddy iris, with small irregular pupil and sluggish reaction to light, is indicative of iritis.

The *position* of the iris must be noted, especially the plane in which it lies (*vide* pp. 248, 258, 268). Special attention should be paid to any adhesions (synechiæ), anterior—to the cornea, or posterior—to the lens capsule. *Tremulousness* of the iris (iridodonesis) is seen when the eyes are moved rapidly if the iris is not properly supported by the lens, *e.g.*, in absence, shrinkage, or dislocation of the lens, slackness of the suspensory ligament, &c. It is best seen in a dark room with oblique illumination.

The Pupils.—A point which should be examined at an early

stage in every routine examination of the eyes is the condition of the pupils. This is the more important since the routine examination frequently demands the use of a mydriatic, and if the pupils have not previously been noted it may be necessary to require the attendance of the patient on another occasion when the effect of the mydriatic has passed off.

The examination of the pupils requires careful attention to details if trustworthy results are to be obtained. It is best carried out as follows :

Place the patient facing the light, which should not be too bright ; see that the two pupils are equally illuminated. Note the size, shape and contour of each pupil. Cover both eyes with the palms of the hands, preferably without touching the face. Tell the patient to look straight at you. Remove one hand and watch the pupil. Replace this hand and remove the other, watching the other pupil. Note down and compare the results (*direct reaction to light*). Remove one hand so that this eye is exposed to light (it should be shaded from intense light). Watch this pupil as the hand is removed from the other eye. Repeat the process whilst watching the other pupil (*consensual reaction to light*).

Now tell the patient to look quite across the room, as far off as possible. Suddenly hold up the index finger vertically at about six inches from the patient's nose, and tell the patient to look at it. Watch the pupils while he accommodates for the finger (*reaction to accommodation*).

When the reaction to light is feeble and the pupils are already small, it is difficult to be certain of the results in bright diffuse daylight, the corneal reflexes adding to the difficulty. In such cases the patient should be taken into the dark room and the light concentrated upon one pupil by focal illumination. By a slight lateral movement of the convex lens (*vide* p. 82) the focus of light can be moved on or off the pupil, the pupillary movements being watched the while. If there is no movement under these conditions we may conclude that the reaction to light is absent.

It is better not to use the ophthalmoscope mirror in this procedure, as the patient is very likely to look at it, and a reaction to accommodation may be mistaken for one to light. Note very carefully if the constriction of the pupil to light is well maintained (*vide* p. 363).

The same method will elicit the *hemianopic pupil reaction* (Wernicke) in the rare cases (lesion of one optic tract) in which it is present. To test for it, the light is placed in front, but

rather to one side of the patient. The light is focussed with the lens upon the opposite side of the retina, and the pupil watched. The light is then moved to the other side and is now focussed on the other side of the retina. If the reaction is present the pupil will react briskly when one half of the retina is illuminated, but very slightly or not at all when the other half is illuminated. It usually reacts slightly even in the latter case, owing to the impossibility of preventing diffusion of light on to the sensitive half of the retina, and for this reason the test is rarely unequivocal.

When the pupils are small to start with ("spinal miosis"), do not react to light, but react to accommodation, the condition is known as the *Argyll Robertson pupil*. It occurs especially in para-syphilitic disease, most commonly and in its most characteristic form in tabes, and frequently in general paralysis of the insane; but it is also found in other syphilitic diseases of the central nervous system.

If the above directions are carried out we shall have reliable information as to the shape and relative size of the pupils and their reactions. A few of the commoner conditions may be enumerated here.

Very large pupils will suggest that a mydriatic has been used. It is not uncommon for it to have been used inadvertently. We not infrequently see a patient with the right pupil widely dilated complaining of dimness of vision. Inquiry will often elicit the fact that he has been using a liniment for rheumatism. The explanation is that the liniment contained belladonna, and that after using it with his right hand he rubbed his right eye with the soiled fingers. Often patients use ointment or drops prescribed for other patients. We must always be on our guard against such traps. These pupils are usually quite immobile, and the patient complains of dimness of vision, especially in near work.

The pupil is also large and immobile in complete atrophy of the optic nerve: this may be due to absolute glaucoma. In acute glaucoma it is usually large, immobile, and oval with the long axis vertical; the condition is generally unilateral. If only one eye is blind from disease of the optic nerve this pupil is rather larger than its fellow as a rule, but the consensual reaction to light on the sound eye is much increased, granted of course that the third nerve is intact. Dilatation of the pupils with retained mobility is found sometimes in myopia and in conditions of impaired nerve tone, *e.g.*, anæmia; it is also found in cases of disseminated sclerosis with optic atrophy,

which rarely if ever leads to complete loss of sight in this disease. Dilatation as a reflex to painful impressions has already been mentioned. Unilateral dilatation may result from irritation of the cervical sympathetic in the presence of glands in the neck, pneumonia, phthisis, chronic pleurisy, cervical ribs, thoracic aneurysm, &c., but it is rare from this cause : it may also be due to syringomyelia, acute anterior poliomyelitis and meningitis affecting the lower cervical and upper dorsal part of the spinal cord and to pressure on the sympathetic fibres leaving the cord in the lower cervical and upper dorsal ventral roots. Many of these causes lead eventually to constriction of the corresponding pupil from paralysis of the sympathetic. Temporary dilatation of one pupil is not very uncommon. Unilateral dilatation with immobility may result from a blow on the eye (Chap. XXI.).

The pupils are small in babies and in old people. When the pupils are small, as under the influence of bright light, the relation to the centre of the cornea can be best seen : the centre of the pupil is usually a little to the nasal side of the centre of the cornea. Small pupils are rarely perfectly round. A small immobile pupil should make us suspect old iritis with posterior synechiæ, and should lead to investigation with a mydriatic —homatropin for diagnostic purposes—to see if the pupil dilates regularly. Bilateral small immobile pupils make us suspect disease of the central nervous system (*e.g.*, pontine hæmorrhage) ; further examination may show that the immobility is confined to reaction to light (*vide* p. 89). A small sluggish pupil, with muddiness of the iris, is associated with iritis, which may be primary or secondary to corneal trouble.

Very small immobile pupils suggest the use of drugs, either locally, *e.g.*, eserin, or through the general system, *e.g.*, morphia.

The chief causes of inequality of the pupils have already been mentioned incidentally. As indicative of central nervous disease it is found in general paralysis and tabes.

We must note carefully the nature of the contraction when the pupil reacts to light, and especially if the constriction is well maintained (*vide* p. 363).

Direct reaction to light does not eliminate the possibility of the patient being blind, *e.g.*, in uræmia and post-basic meningitis.

The Lens.—The lens cannot be thoroughly examined without the assistance of the ophthalmoscope. By inspection, aided if necessary by focal illumination, we note any opacities

in the pupillary area. The pupil may be blocked with iritic exudates (inflammatory pupillary membrane, blocked pupil).

Opacities in the lens itself are seen by oblique illumination as grey, white or yellowish patches. According to their distribution and nature we diagnose the various forms of cataract, but our observations must always be confirmed and controlled by ophthalmoscopic examination. The following example will show how easily one may be led astray. When the light is concentrated by focal illumination upon the pupil of a young person's eye the lens substance seems almost perfectly clear; at most we see a faint bluish haze. If we examine the lens of an old person in the same manner the haze is much more pronounced; the lens substance in fact looks slightly milky. We might conclude that the patient has cataract. Examination with the ophthalmoscope will, however, show a perfectly clear red reflex. The explanation is that the lens substance becomes more optically dense, *i.e.*, the refractive index increases, as the person grows older (*vide* p. 49). Now the higher the refractive index of a substance the greater will be the reflection from its surface. The milkiness which we see is due to rays of light which are reflected from the lens and enter our eyes. The more rays reflected the more will the lens appear milky.

If, however, the white appearance is very pronounced, and especially if it is strictly localised to certain parts of the lens substance, we may safely diagnose cataract. A spot in the centre of the pupil, looking as if it were on the surface of the lens, may be a pupillary exudate or an anterior polar cataract. Triangular spokes of opacity with their apices towards the centre are indicative of senile cataract. A very white appearance over the whole pupillary area suggests a total cataract; if it is yellow and the iris is tremulous we suspect a shrunken calcareous lens.

The Tension.—Last in the external examination, but by no means of least importance, we test the tension of the eye. It is done in the same manner as testing for fluctuation in other parts of the body.

Stand facing the patient, who is told to keep looking towards his feet. Place the index fingers of both hands side by side upon the upper lid, steadying them by the other fingers lightly applied to the brow. Keep one finger quite still, pressing upon the globe through the lid. Now attempt gently to indent the globe with the other finger, pressing directly downwards, concentrating the attention meanwhile on the impression

which is conveyed to the stationary finger. Repeat the process on the other eye.

The student should practise this manœuvre on a number of healthy eyes. He will thus obtain a mental estimate of what is to be considered normal tension—Tn. In absolute glaucoma the eye is usually stony hard; this condition is generally indicated by the convention T + 3, a misuse of numbers, but one in common use. The gradations of increased tension from Tn to T + 3 are usually indicated by the conventions T full (T +), T + 1, T + 2. Similarly the gradations of diminished tension are represented by T minus (T —). T — 1, T — 2, T — 3.



Fig. 72.

Instruments known as *tonometers* have been devised for measuring the tension of the intact eye. They are far less reliable than the manometer, which, however, cannot be used on the human eye. The best tonometer for clinical use is that invented by Schiötz (Fig. 72). With it the depth of the indentation in the cornea made by a weighted stylet is measured by a lever which travels over a scale. There are four weights (5·5, 7·5, 10 and 15 gms.) and the greatest accuracy is attained with the weight which gives a deflection of the lever of 2 to 4 mm. The instrument is calibrated so that the equivalents of the readings in millimetres of mercury can be read off a chart. It is doubtful whether the readings are accurate when transformed into pressures in millimetres of mercury, but the tonometer is certainly useful for comparative measurements, *e.g.*, between the two eyes or between successive measurements on the same eye.

CHAPTER VII.

OPHTHALMOSCOPIC EXAMINATION.

The internal parts of the eye beyond the lens cannot be seen without the assistance of the ophthalmoscope. A little consideration of the optical conditions of the eye will show the reason.

Under ordinary circumstances the pupil looks black, and no red reflex, much less a clear image, is obtained from the fundus. If, as in Fig. 73, there is a source of light, L, in front of the eye, and the eye is focussed upon it or accommodated for it, the light and a spot upon the retina are conjugate foci ; *i.e.*, the image of the spot of light is a spot on the retina.



Fig. 73.

Reversing the direction of the rays, all rays from the illuminated spot of the retina are brought to a focus at the source of light. Therefore no rays will enter an observing eye unless it is situated actually at the source of light. The problem solved by Babbage (1848) and rediscovered by von Helmholtz when he invented the ophthalmoscope (1851) was practically that of making the observing eye at the same time the source of illumination of the observed fundus.

If the eye is not focussed for the source of light the conditions are different, and some slight luminosity of the pupil may be seen. This is one cause of luminosity in the pupils of the hypermetropic eyes of young children and most carnivora. Extreme hypermetropia is also the cause of the so-called amaurotic cat's eye, which is due to detachment of the retina, glioma of the retina, &c. In these cases the retina is pushed forwards and the fundus at this spot becomes highly hypermetropic, the reflex from the pupil being often the first

symptom noticed. The same principle applies to the reflex from the eye after the lens has been removed by extraction of cataract.

In hypermetropia the conjugate focus of the source of light, L, is a point, *l*, behind the retina (Fig. 74). Hence the emergent rays from the illuminated area of the fundus are divergent, as if coming from *l*. Therefore an observing eye situated anywhere within the area l_1 l_2 of the cone of emergent rays will catch some of them, and the pupil of the observed eye will appear feebly illuminated. Under these circumstances it is not necessary for the observing eye to occupy the exact position of the source of light, but only a spot in its immediate neighbourhood. On the same principle, the extremely hypermetropic retina in glioma retinæ, &c., can be seen well by focal illumination.

In high myopia the emergent rays are strongly convergent,



Fig. 74.

and become divergent after coming to a focus at the remote point (Fig. 33). Beyond this point some of the divergent rays may enter an observing eye suitably situated and the observed pupil appears illuminated.

The luminosity of albinos' eyes is due to light entering the eye, not only through the pupil, but also through the iris and sclerotic. That this is the true explanation is shown by the fact that the pupil looks black if it is observed through a small hole in an opaque screen. A small amount of light passes through the sclerotic in the normal eye.

It will help us to understand the principles of the ophthalmoscope if we say a few words about its historical development. The ophthalmoscope was invented by Babbage in 1848, but its importance was not recognised, and it was rediscovered by von Helmholtz in 1851. The original ophthalmoscope of von Helmholtz was merely a plane plate of glass (Fig. 75). A source of light was placed beside the observed eye and the glass plate obliquely in front of it, so that a portion

of the light was reflected from the surface of the plate into the eye. On looking through the transparent plate an observer could now receive some of the rays from the fundus into his own eye, and thus obtain an image of the illuminated fundus. Since but a small proportion of the light received upon the plate is reflected at its surface the illumination is feeble. Nevertheless, the principle is worth bearing in mind as a ready means of getting a view of a fundus in the absence of a more satisfactory ophthalmoscope. Moreover, an error of refraction in the observed eye may be obviated by using the corre-



FIG. 75.—Diagram of von Helmholtz' ophthalmoscope. O_1, observed eye; O_2, observer's eye; L, source of light; *l*, image of L formed by the plane mirror—immediate source of light; M E H, relative positions of retina in myopia, emmetropia, and hypermetropia respectively, showing the relative sizes of the areas of retina illuminated in each case.

sponding spectacle glass of the patient as the ophthalmoscopic mirror.

Von Helmholtz next increased the amount of light reflected by superposing three plane plates. The back of the glass was next converted into a more powerful mirror by silvering it, leaving a small portion unsilvered or leaving a hole in the mirror, through which the observer might look. The illumination was still feeble, since the rays reflected by a plane mirror are divergent (*vide* p. 21). Ruete therefore (1852) introduced the perforated concave mirror which still holds the field. The final modification was the addition of a battery of small lenses of various strengths, which might be brought into position behind the aperture. The multitudinous forms of

"refraction ophthalmoscopes" are merely various mechanical contrivances for doing this most conveniently.*

There are two chief methods of ophthalmoscopic examination, the direct method (v. Helmholtz, 1851) and the indirect method (Ruete, 1852). The ophthalmoscope is provided with two mirrors, one a small one, slightly tilted, for the direct method, the other a large one, not tilted, for the indirect method. Both are concave, the former with a focal distance of 30 cm., the latter of 10 cm. It is an advantage to have also two plane mirrors corresponding with these; with such a four-mirror ophthalmoscope the surgeon is fully equipped for every detail of ophthalmoscopy and retinoscopy.

The importance of system in his methods is so often impressed upon the student that he is liable to underrate it through sheer reiteration. It will perhaps suffice to say here that in using the ophthalmoscope he will inevitably come to grief unless he pursues his examination on a well ordered plan. The order of examination should be as follows:

(1) Preliminary examination with the mirror alone at a distance of about 1 metre from the patient;

(2) Examination with the mirror alone at a distance of about 20 cm. (reading distance) from the patient; this is sometimes called the distant direct method;

(3) Examination by the indirect method;

(4) Examination by the direct method.

The following facts will impress upon the student the reasonableness of this procedure. By (1) we obtain knowledge of the nature of the refraction of the eye under examination: this will prevent many little difficulties when we come to closer quarters. By (2) we see any gross changes, especially opacities in the refractive media; these may be made at once evident by this method, whereas they may be very puzzling if first observed by (3) or (4). In addition, we shall see the details of any very hypermetropic part of the fundus, such as a detached retina or glioma of the retina; these also are by

* The student is advised to procure a good ophthalmoscope at the outset of his clinical work in the medical wards. The cheaper forms are not only waste of money, but are a perpetual source of annoyance. The modification of Couper's ophthalmoscope, generally known as Morton's, is most strongly recommended. In recent years various self-luminous ophthalmoscopes have been devised. In most of these half the aperture of the ophthalmoscope is used for illuminating the fundus from a small electric bulb contained in the instrument, the other half being used for observation. The lamp is run off a small dry battery which may be placed in the handle of the ophthalmoscope. These instruments are particularly useful for examining bed-ridden patients.

no means difficult to miss by (3) and (4). By (3) we get a general view of the fundus—the largest possible area under moderate magnification ; it is exactly comparable to microscopic examination with a low power. By (4) we examine details under a higher magnification ; it is exactly comparable to microscopic examination with a high power.

The student should commence by taking a patient whose pupils have been dilated with atropin, *e.g.*, a boy of twelve or fourteen who has come for the correction of his refraction. The atropin will have had the additional advantage of having paralysed the patient's accommodation. The observer should know his own refraction.

The patient is taken into the dark room and seated beside the light. The light is placed to the side which is to be examined, but well behind the level of the patient's face ; the eye should be as much as possible in darkness. The observer sits facing the patient, about a metre from him. He reflects the light from the large ophthalmoscope mirror into the eye, meanwhile looking through the sight-hole. This requires a little practice, but is quickly mastered. When the light falls on the eye he notices a red reflex from the pupil. There ought to be no black spots in the pupillary area, but either a uniform red reflex or obscure details of the fundus. By tilting the mirror to and fro in various directions he can obtain an approximate idea of the refraction of the eye (*vide* p. 101).

The observer now stands up and approaches the patient until his eye, still with the large mirror, is about 9 inches from the eye under observation. He can now see the cornea and iris clearly, and can confirm any points which he has made out previously by the external examination.

He then sits down again at about a metre from the patient. Still keeping the light upon the eye with the large mirror, with his left hand he holds the large convex lens, which he will find in the ophthalmoscope case, close in front of the eye. He will be wise not to hold the lens absolutely vertical, but to tilt it very slightly. He will probably see only the magnified iris through the lens. He now watches the red reflex from the pupil, and slowly withdraws the lens from the eye towards himself. At a certain point he will see an inverted image of the fundus quite clearly. The indirect method requires also some practice, but the amount required may be much diminished and much greater accuracy attained if the optical conditions under which the examination is made are thoroughly understood. These will be explained immediately.

Having obtained a good general view of the fundus, the observer again approaches the patient. He now uses the small tilted mirror of the ophthalmoscope. The mirror is tilted so that it faces towards the light. Looking through the sight-hole he first gets the light upon the eye; this is best effected from a short distance away. When the light is well on the pupil and the observer can see the red reflex he approaches slowly nearer and nearer, watching that the light does not leave the pupil, until his brow is almost or quite touching the patient's brow. If now both the patient and the observer are emmetropic, the inexperienced observer will probably see the details of the fundus only indistinctly. He should then turn up with his index finger applied to the milled disc on the back of the ophthalmoscope successively stronger concave glasses (usually marked with white numerals). He will then probably see the fundus quite clearly. The image is erect, *i.e.*, the opposite of that by the indirect method. Here again practice is needed, and a knowledge of the optical conditions is quite essential.

We will now consider the chief features which are to be learnt in each stage of the examination, and how they are to be learnt.

I. Preliminary Examination with the Mirror at 1 metre.—We will suppose that the observer is emmetropic, or that his refraction has been corrected, and that the accommodation of the observed eye is at rest or paralysed. In examining the right eye the patient is told to look at the observer's right little finger which is held up; this is easily done while holding the ophthalmoscope. In examining the left eye the patient is told to look at the observer's left ear. In this manner the optic disc, which lies a little to the nasal side of the posterior pole of the eye, is brought into the observer's line of vision. When the optic disc is opposite the pupil we shall notice from a distance of 1 metre that the red reflex becomes much paler or even whitish.

If the eye is highly hypermetropic or myopic we shall see some details of the fundus, *e.g.*, a few vessels running across the reflex. The explanation is easy from what we have already learnt.

Consider first the hypermetropic eye. If we think of two spots on the retina, say at opposite edges of the disc, the rays reflected from these points will form two bundles of divergent rays when they leave the eye, just as if they came from the corresponding virtual remote points behind the eye (Fig. 76).

The greater the distance from the eye, the greater will be the area over which these divergent rays will spread, so that at 1 metre some of the peripheral rays of each pencil will enter the observer's eye. By a slight effort of accommodation the observer will be able to bring these divergent rays to a focus on his retina, so that he will obtain a clear image of each point, and necessarily also of the intervening region. It is exactly as if the eye were taken away and the two points were situated at the remote points. Hence the image will be erect.

If the observer now shifts a little to one side, the observed eye remaining stationary, more rays will enter his eye from the neighbourhood of the opposite point, and less from the neighbourhood of the point on the same side as that to which his movement is directed. Although the points remain



FIG. 76.—Examination with the mirror at 1 metre. O_1, observed eye, which is hypermetropic ; O_2, observer's eye, emmetropic, but accommodated for the divergent rays from O_1.

stationary, more of the fundus on the opposite side and less of the fundus on the same side will be seen. Hence the points will seem to move in the same direction as his own movement. The observer mentally regards the very sharp outline of the pupil as a fixed object of comparison, and as more of the fundus on the opposite side comes into view, whilst a corresponding amount on the same sdie disappears, this is mentally interpreted as a movement of the image in the same direction.

If, therefore, when the light is reflected into the eye at a metre distance we see vessels in the pupillary reflex, and if they appear to move in the same direction when the head is moved slightly to one side, we conclude that the eye is hypermetropic.

Consider now the myopic eye (Fig. 77). Here the emitted rays from the two points will be strongly convergent in each case, and a real inverted image of the points and intervening

area will be formed at the remote point of the eye, *i.e.*, between the observer and the observed eye. The rays will diverge from this image, and the effect will be exactly the same as



FIG. 77.—Examination with the mirror at 1 metre. O_1, observed eye, which is highly myopic; O_2, observer's eye, emmetropic, but accommodated for the divergent rays from the far point of O_1.

if there were an actual inverted object in this position. If th myopia is sufficiently high, the image will be beyond the observer's near point, so that he will be able to accommodate for it. If he moves to one side he will see more of the observed



FIG. 78.—Examination with the mirror at 1 metre. O_1, observed eye, which is emmetropic; O_2, observer's eye: none of the rays from the widely distant points on the fundus of O_1 enter O_2. If the points are close together the rays of the two bundles will be nearly parallel, and would form a clear image on the retina of O_2 if the accommodation of O_2 were almost completely in abeyance.

fundus on the same side and correspondingly less on the opposite side, so that the fundus will appear to have moved in the opposite direction.

If, therefore, when the light is reflected into the eye at a metre distance we see vessels in the pupillary reflex, and if

they appear to move in the opposite direction when the head is moved slightly to one side, we conclude that the eye is myopic.

What will happen in emmetropia (Fig. 78) or low myopia, for the effect will be similar ? Here the rays passing out of the eye from the two points will be parallel or very slightly convergent, and their direction will be that of their axes, which is the continuation of the lines joining the points with the nodal point of the eye. As these axes constantly diverge from one another, the observer at a distance of 1 metre cannot receive portions of both pencils of rays upon his own pupil, consequently he cannot obtain a clear image of the whole intermediate region between the spots. He may get a clear image from two spots very close together, but only if his accommodation is almost completely suspended, so that nearly parallel rays are brought to a focus upon his retina.

The same reasoning applies to low hypermetropia, for here the remote point of the eye is so far behind the retina that the rays diverge very little when they leave the eye, so that they are almost parallel.

If, therefore, when the light is reflected into the eye at a distance of a metre we see only a red reflex in the pupil, without any details, we conclude that the eye is either emmetropic or has only a low degree of ametropia.

A still simpler means of discovering the condition of the refraction is as follows. Still throwing the light into the eye with the large concave mirror we tilt it gently in various directions. We shall see a shadow move across the pupil ; if the shadow is very dark there is considerable error of refraction. If it moves in the opposite direction to that in which we move the (concave) mirror the eye is hypermetropic ; if in the same direction it is myopic. This method is used for correcting refraction, and we shall consider it in detail later (*see* Retinoscopy).

II. Preliminary Examination with the Mirror at the convenient distance for near vision (22 cm.).—At this distance the observer will be most suitably situated for distinct unaided vision, and he will be able to examine the superficial parts of the eye more accurately. If he is presbyopic he will naturally have to correct his presbyopia, and he may have to use a convex lens if he is strongly hypermetropic. If he is very myopic he will have to approach closer.

The advantages of a preliminary examination in this manner are (1) the recognition of opacities in the refractive media ;

(2) the recognition of a detached retina or other substance not far behind the lens ; (3) the confirmation of the results found by the external examination.

(1) *The diagnosis of opacities in the refractive media.*—If the eye is normal there will be a red reflex from the pupil. If there is any opaque body in the course of the rays reflected from the fundus it will stop these rays and will therefore appear black. The whole field may be black, as when the lens is entirely opaque, or when there is blood in the vitreous. In the latter case oblique illumination will show the red blood behind the transparent lens if it is sufficiently far forward in the vitreous : the blood looks red in this case because of the light reflected from its surface.

Opacities vary in shape, size, and position. We are particularly concerned to discover their position, as this frequently gives the key to their nature.

The first point to determine is whether the opacity is movable. This is done by telling the patient to move his eye in different directions—towards the ceiling, towards the floor, to the right, to the left—and then to look straight forward. A floating opacity will then continue to move after the eye is brought to rest. It must therefore be either in the aqueous or vitreous. In the former position it can be seen and diagnosed by other methods. If it is in the vitreous and is freely movable we also learn that the vitreous is fluid, which is not its normal consistency. If the opacity moves only with the eye it may be in the cornea, lens, or vitreous, which, under these circumstances, will have its usual viscous consistency.

The next point is to determine its exact position. This is effected in the preliminary examination with the mirror alone by *parallactic displacement.*

In Fig. 79, if 4 is the centre of rotation of the eye, and if there are opacities at 1, 2, 3, 4, 5, then, when the eye is rotated a small amount, the opacities, 1, 2 and 3, in front of the centre of rotation will move in the direction of rotation, and 5, behind the centre, will move in the opposite direction, while 4, at the centre, will not move. It is obvious that the amount of movement will be greater the farther the opacity is from the centre of rotation. Now, we have no means of defining the centre of rotation by ophthalmoscopic examination, but all the movements will be referred to the edge of the pupil for comparison (*vide* p. 99). If the observer is situated at A, all the opacities will appear as a single spot in the centre of the pupillary reflex. If he shifts his position to B, or if the

eye is rotated a corresponding amount in the opposite direction, the opacity 2 will remain in the centre of the pupil, whilst 1 will appear to move towards one edge of the pupil, and 3, 4 and 5 towards the opposite edge, 5 being lost entirely behind the iris.

Hence we deduce the rule that if the eye is moved slightly in a given direction, opacities in the pupillary plane will appear stationary ; those in front of that plane will move in the same direction, and those behind will appear to move in the opposite direction, the amplitude of apparent movement being a rough indication of their distance from the pupillary plane.

There is another guide which we may make use of, viz., the corneal reflex. This is the image of the mirror formed by



Fig. 79.

the cornea. With the ordinary concave mirror it is a virtual image (*vide* p. 22) situated about 4 mm. behind the anterior surface, *i.e.*, a short distance behind the anterior surface of the lens (behind 2 in Fig. 79). The centre of curvature of the cornea is situated 8 mm. behind its anterior surface, *i.e.*, less than 1 mm. behind 3 (Fig. 79). The corneal reflex will always cover this latter spot, the centre of curvature of the cornea, no matter what the position of the eye. Hence an opacity situated here will always be covered by the corneal reflex ; opacities in front of the centre of curvature move in the same sense with regard to the reflex as the eye moves ; and opacities behind it move in the opposite direction to the movement of the eye. Therefore, in Fig. 79, in the first position of the eye, the opacities 1, 2, 3, 4, 5, will all appear in the centre of the corneal reflex (A′) : in the second position

they will appear as in B′ ; so that an opacity at the posterior pole of the lens will scarcely leave the edge of the reflex, whereas an anterior polar opacity will move much farther from it.

One peculiar apparent opacity is seen by the mirror alone, and this method affords the surest means of discovering the defect. This is the edge of a dislocated lens, or the notch in the edge of the lens in congenital coloboma of the lens. When the edge of the lens crosses the pupillary area it is seen as an intensely black crescent, sharply defined peripherally but merging centrally into the clear red reflex. The reason of this appearance is that the whole of the light which falls upon the extreme edge of the lens is totally reflected ; none of it leaves the eye, so that none can enter the observer's eye.

We not infrequently meet with very fine opacities, especially in the vitreous. If we use a concave mirror and a bright light we shall probably fail to see them, the reason being that these very delicate opacities are partially transparent, so that if the light is very bright some passes through them and contrast is reduced. Contrast is further reduced by reflection of light from their surfaces. They are, as it were, drowned in light. In order that we may be sure of not missing fine opacities the best method to adopt is to use a plane mirror. The rays reflected from a plane mirror are divergent (*vide* p. 21) ; hence less light enters the eye. If we have no plane mirror available the light should be reduced, but this is not so satisfactory. We may increase our chances of seeing the specks if we place a convex lens behind the mirror, which will have the effect of magnifying them (Fig. 21).

(2) *The recognition of detached retina, &c.*—A detached retina is situated much in front of its normal position ; it is therefore in the position of the fundus of a very hypermetropic eye. When light is thrown in from a mirror at reading distance light will probably be reflected from the surface in such a way that some rays will enter the observer's eye. A difference of reflex in different directions is noticed, red in some, grey or black in others. More minute investigation will reveal a whitish or greyish uneven surface upon which there are almost black wavy lines ; these are the retinal vessels. Particular stress is laid upon this point, because the appearances of a detached retina by the indirect and direct methods may be very puzzling to the beginner. If the precaution is always taken of commencing the examination with the mirror alone little difficulty is likely to arise.

Besides a detached retina we shall also be able to see anything else in a similar position, *e.g.*, a tumour pushing the retina forwards, or a tumour of the retina itself (glioma), and so on.

None of these will be seen unless they are pushed forwards very considerably : hence we must not asseverate the absence of a detached retina, &c., if we fail to see it by this method.

(3) *Confirmation of the results found by the external examination.* —We are able by this method not only to confirm the results previously arrived at by external examination, but also to supplement them by important subsidiary information. Thus we are able to map out the limits of opacities in the lens much more accurately, since they now appear black on a red background, and as has already been shown we can determine their exact position with much greater precision.

We may have noticed a black spot in the iris in a case with the history of a foreign body having gone into the eye. It is probable that the foreign body has passed through the iris, and that the black spot is a hole. The examination with the mirror often at once settles the question, for if there is a hole we shall be able to find some position in which a red reflex can be seen through the hole. The absence of a red reflex does not prove the absence of a hole, for the lens may be opaque behind the hole.

The following is a somewhat similar example. We have noticed a black patch at the ciliary margin of the iris, convex in outline towards the pupillary margin. It may be a melanotic sarcoma of the ciliary body growing forwards and implicating the iris : or it may be a separation of the iris from its ciliary attachment (iridodialysis). In the latter case it will be possible to obtain a reflex through it by the mirror, whereas in the former it will be opaque.

We have said that by this method opacities in the refractive media appear black. Superficial opacities, however, such as those in the cornea and near the anterior surface of the lens, can be seen in their natural colours by approaching still nearer to the eye. Under these conditions more light is reflected from the surface of the opacities and some of it enters the observer's eye. It will be objected rightly that now we shall be within our near distance and consequently shall not be able to see anything clearly. This is true, but it can be obviated by assisting our accommodation by putting up gradually stronger convex glasses behind the ophthalmoscope mirror as we approach the eye. This has the additional advantage of magnifying the opacity. If we approach very

close to the eye and place a + 20 D lens behind the mirror we shall see the opacities highly magnified. This glass will be acting very much like an ordinary magnifying glass, so we shall have to focus it in much the same manner. We therefore start a little distance from the cornea and watch carefully as we get nearer and nearer ; there will come a point when the opacity is very clearly defined.

Suppose now that under the same circumstances, with the + 20 D lens in the position for seeing the cornea, we wish to examine an opacity near the surface of the lens it can be done in two ways. We may continue to approach still nearer until it comes into focus ; or we can use a weaker lens, retaining our



Fig. 80.—Indirect method. Illumination of the fundus, showing the course of rays from the source of light to the mirror, through the lens, and through the eye ; also the area of the field of illumination.

original position. The weaker lens is most easily produced by moving up stronger and stronger concave lenses in front of the + 20, until the opacity is accurately focussed. This is made possible in most ophthalmoscopes by having a + 20 D lens set in a disc which lies behind the sight-hole : the lens can be turned into position when required, and does not interfere with the use of the other lenses at the same time. The opacity in the crystalline lens will of course not be quite so highly magnified by the second as by the first method.

III. The Indirect Method. - The indirect method of examination with the ophthalmoscope consists essentially in making the eye, whatever be its refraction, highly myopic by placing a strong convex lens in front of it (Figs. 80—82). The effect of this will be to form a real inverted image of the

fundus between the observer and the convex lens, as will be easily understood from the accompanying diagrams. If the eye is already myopic the convergent rays which come from any point on its fundus will be made still more convergent by the lens, and the inverted image which is always formed in myopia will be brought close to the lens. If the eye is emmetropic the parallel rays emitted will be made strongly convergent, and where they cross the inverted image will be formed. If the eye is hypermetropic the rays will still be made convergent, for the lens used is so strong that the divergence in hypermetropia is never strong enough to prevent it.

It will be seen that with the same lens the inverted image is formed at different distances beyond it according to the refraction of the eye. If the lens is kept at a constant distance from the eye, *e.g.*, its own focal distance, the emmetropic image will be formed at the focal distance of the lens beyond it: the myopic will be nearer to the lens, the hypermetropic farther from it (Fig. 82).



FIG. 81.—Indirect method. Emergent rays from the fundus, showing the formation of the image. In the figure the lens is situated at the anterior focal plane of the eye; the rays which are parallel inside the eye, therefore, pass through the optical centre of the lens. The rays which pass through the nodal point of the eye are rendered convergent by the lens. The points where these two systems of rays cross gives the position of the image, which is seen to be inverted.

In all cases the image is magnified, the amount of magnification depending upon the refraction of the eye, the strength of the lens, and its distance from the eye. With a + 13 D the fundus of an emmetropic eye is magnified about five times.

One of the greatest difficulties in using the indirect method is the group of reflexes formed by the eye and the surfaces of the lens. We have seen that the cornea forms a reflex of the mirror when it is used alone. This reflex, when seen through the convex lens, is magnified, so that it may cover the pupil and prevent anything behind being seen. But the surface of the lens towards the observer acts like another convex mirror and forms another reflex situated behind the lens. Similarly the surface of the lens near the patient acts like a concave mirror and forms a reflex on the observer's side of the lens. These reflexes are very troublesome, but they may be got out of the way by a little manœuvring. It has been said that the

two lens reflexes, which are the most troublesome, are images of the mirror formed on opposite sides of the lens. If we tilt the lens a little it will be found that these reflexes move in opposite directions, and we can look quite comfortably between them. We must be careful not to tilt the lens more than is necessary, because if we look obliquely through a tilted lens objects appear distorted: in fact we produce one type of astigmatism. The distorted image of the disc produced in this manner may be attributed to astigmatism in the eye, when none is really present.

Another difficulty which the beginner usually experiences is due to getting too close to the patient. If he understands the position of the image which he is looking for, as described above, he will discover why he can see no sharp image when



FIG. 82.—Indirect method. Position of the image according to the refraction of the eye. In this figure the lens is situated at its own focal distance from the cornea. In emmetropia the parallel emergent rays, therefore, cross at the principal focus of the lens, E. In myopia the convergent emergent rays cross nearer to the lens than its principal focus, viz., at M; in hypermetropia the divergent emergent rays cross farther from the lens than its principal focus, viz., at H.

he is too close to the patient. Most people using the indirect method think that they are looking at the pupil. As a matter of fact, when the fundus is seen clearly, they are not accommodating for the pupil, but for the real image of the fundus, which is in the air somewhere between the lens and the observer. Now we can only see an object clearly with the unaided eye if it is at a convenient distance away. Consequently, if the observer gets so close to the patient that he is less than the distance of his near point from the aerial image he cannot see it clearly.

If we like to do so we can get over this difficulty and still remain closer to the patient. In order to do so, the observer must help his accommodation by putting up a convex lens in front of his eye. If we put up a + 1 D or + 2 D behind the ophthalmoscope mirror, we shall not only see the image clearly

at a shorter range, but we shall also magnify it, an additional advantage.

As regards the position of the convex lens before the patient's eye, there is a considerable range over which we can see the fundus quite well, but some positions are better than others. In practice we find the best position by putting the lens close to the eye to start with, steadying it by the little finger applied to the patient's brow, and gradually bringing it farther away from the eye until the best position is obtained.

Theoretically, from the point of view of the maximum field of fundus seen, the best place for the lens is its own focal distance from the patient's pupil. But this is the very worst place from the point of view of the corneal reflex. The latter is situated at about the level of the iris (4 mm. behind the cornea, *vide* p. 103). If the convex lens is at its focal distance from it, the rays from this image will be made parallel by the lens, *i.e.*, the reflex will fill the whole area of the lens, and we shall see nothing else. Hence the best position for practical purposes is either nearer to or farther from the eye than this position. We shall see later that a convenient distance is when the lens is at its focal distance from the anterior focus of the eye. Here, slight tilting of the lens, besides shifting the lens reflexes out of the way, will also move the corneal reflex and the image of the fundus in opposite directions, and so get the corneal reflex out of the way.

We can tell by the indirect method whether the eye is emmetropic or ametropic by observing the effect of shifting the lens on the size of the image of the fundus. We use the disc as the best guide, getting it into the field by telling the patient to look in the appropriate direction, *i.e.*, towards the raised right little finger when examining the right eye, towards the left ear when examining the left.

Place the lens close to the eye and gradually bring it farther away. If the image of the disc does not alter in size the eye is emmetropic; if it gets smaller the eye is hypermetropic; if it gets larger the eye is myopic.

If we understand why this is so, we shall be able to remember what happens in each case. Imagine two points, *a* and *b*, upon the fundus, *e.g.*, upon opposite edges of the disc (Fig. 83). If they are illuminated, one of the many divergent rays emitted by each must be parallel to the axis. These two rays, when they pass out of the eye, will cross at the anterior focus of the eye, whatever its refraction may be, as long as the error is one of

undue length or shortness (*axial ametropia*). They will cross at the anterior focus because they are parallel to each other before refraction. Now, suppose that the convex lens used in the indirect method is situated at its focal distance from the anterior focus of the eye. These two rays, after they have crossed, will meet the convex lens, and since they come from the focus of the lens they will be parallel to each other after refraction. As we have already seen (p. 107), under such circumstances the image of the fundus is formed in emmetropia at the focal distance of the convex lens (F^1) from it (E): in myopia the image will be nearer the lens (M), in hypermetropia farther away (H). The two rays which we have been considering must represent the images of the two points on the fundus. Hence the distance between them after refraction by the eye and by the lens will give the size of the image of the portion of fundus between *a* and *b*. We see therefore that when the lens is at its own focal distance from the anterior focus of the eye the magnification of the image



Fig. 83.

is the same in emmetropia, in axial myopia, and in axial hypermetropia.

If the lens is nearer the eye than the above distance (Fig. 84), the rays under consideration will diverge after refraction by the convex lens. Hence, if the lens is less than its own focal distance from the anterior focus of the eye, the magnification is greatest in axial hypermetropia, least in axial myopia, and intermediate in emmetropia. Conversely, if the lens is farther from the eye, the rays under consideration will be convergent after refraction by it. Hence, if the lens is more than its own focal distance from the anterior focus of the eye, the magnification will be greatest in axial myopia, least in axial hypermetropia, and intermediate in emmetropia.

In *curvature ametropia*, such as we meet with in astigmatism, the results are not quite the same, but they are easily deduced if we remember that there are now two anterior foci to the eye, one for each meridian. There are also two nodal points. When the lens is at its focal distance from the cornea the magnification is the same in emmetropia and any ametropia of curvature.

Under these circumstances the disc appears circular. If the lens is nearer the eye the image is elliptical, with its long axis in the less refractive meridian, *i.e.*, generally horizontal. If the lens is farther from the eye the long axis is in the more refractive meridian, *i.e.*, generally vertical. As mentioned before, it is essential that the lens should be held almost vertical, as any inclination makes it itself astigmatic. If the disc is really oval, as in high myopia, the axis of the ellipse will of course remain unaltered.

Ametropia of index of refraction occurs in old age. Aphakia, the condition when the lens has been removed, may be considered an extreme form of index ametropia. Here the position of the nodal point remains invariable, and if the convex lens is at its focal distance from this point the image is the same size in emmetropia and ametropia of index of refraction. If the lens is moved



FIG. 84.

closer to the eye the image increases in hypermetropia and diminishes in myopia, whilst it remains the same in emmetropia. Since the image is formed at a considerable distance beyond the focus of the lens in the high hypermetropia of aphakia it is convenient to use a stronger lens, *e.g.*, + 18 D.

Differences of level of two points near each other on the fundus are made very evident by parallactic displacement in the indirect method. Thus, in Fig. 85, if there are two spots, *a* and *b*, at different levels in the fundus, *e.g.*, on the edge of the disc and at the bottom of a glaucoma cup, when the lens is shifted slightly so that its optical centre moves from o_1 to o_2, the images of *a* and *b* will move from a_1 to a_2, and b_1 to b_2. It is of historical interest that this displacement was at one time wrongly interpreted, so that a glaucomatous cupping of the disc was diagnosed as a swelling.

IV. The Direct Method.—In the direct method the observer

approaches as close as possible to the patient's eye (Fig. 86). If the eye is hypermetropic the emergent rays will be divergent, as if coming from the virtual remote point behind the eye. Owing to the short distance between the eyes a large pencil will fall upon the observer's pupil, and may be brought to a focus upon his retina if he makes a suitable effort of accommodation. If he is presbyopic, or if his accommodation is relaxed, he will only obtain a clear image by placing a convex lens behind the sight-hole of the mirror (Fig. 87, H).

If the observed eye is emmetropic the emergent rays will be parallel, and consequently can only form a clear image upon the observer's retina if his accommodation is absolutely relaxed (Fig. 87, E)—unless, indeed, he counteracts the amount of his accommodation by a corresponding concave lens in front of his eye (*vide* p. 113).

If the observed eye is myopic the emitted rays are conver-



Fig. 85.—Indirect method. Parallactic displacement.

gent. If the myopia is moderate the real image of the fundus at the far point of the eye will be behind the observer's head, *i.e.*, he will catch the convergent rays before they have come to a focus. These convergent rays, entering his emmetropic eye, are brought to a focus in his vitreous ; hence he cannot possibly obtain a clear image unless he counteracts the convergence by an equivalent concave lens behind the mirror (Fig. 84, M). If the observed eye is very highly myopic its punctum remotum will be situated somewhere in the space between the eye itself and the observer's retina, and it may be in such a position that it is impossible to obtain a clear image with any correction. For example, the remote point may be just behind the sight-hole of the mirror. Here it is too close to be accommodated for, and no correcting glass situated at the same position will have any effect upon the rays, for they will nearly all pass through the optical centre of the lens. The practical outcome of this discussion is to get as close to the eye as possible.

Much stress is generally laid upon the necessity and the difficulty of relaxing one's accommodation in examination by the direct method. It is difficult to relax the accommodation entirely when the eye is apparently close to the object looked at. The observer should try to think that he is looking at a very distant object, but even then, as soon as he directs his attention to details of the picture, he is almost certain to accommodate. It is best for the beginner not to worry himself about this point: if he cannot see an emmetropic fundus clearly let him put up minus lenses until he does. After he has acquired facility in seeing anything at all it will be soon enough for him to grapple with this difficulty.

The image by the direct method is always erect. Thus, in



FIG. 86.—Direct method. Illumination of the fundus, showing the course of rays from the source of light to the mirror and through the eye; also the area of the field of illumination. Compare with Fig. 80.

hypermetropia it is exactly as if the observer were looking at an actual object situated at the remote point of the eye, *i.e.*, some distance behind the eye. In myopia the converging rays are caught before they cross; they are made suitably less convergent by the correcting lens behind the mirror, so that again an erect image is seen. In emmetropia the emergent rays are parallel and are, therefore, also caught before they cross—at infinity; hence, again, an erect image is seen.

The image is always magnified, and it is magnified more than by the indirect method. In emmetropia the fundus is seen magnified about fifteen times. In hypermetropia it is magnified less, and in myopia more than in emmetropia.

The field of ophthalmoscopic vision by the direct method, *i.e.*, the area of the fundus which can be seen, varies with the distance of the observer from the eye and with the refraction

of the eye. It increases as the eye is approached—another reason for getting as close to the eye as possible. It is greatest in hypermetropia, least in myopia, and intermediate in emmetropia. Thus, we see the largest area, least magnified, in hypermetropia, and we see the least area, most magnified, in myopia.

In astigmatism the magnification is greatest in the more myopic meridian, and least in the more hypermetropic. In



FIG. 87.—Direct method. Emergent rays from the fundus of the observed eye, O_1, showing the formation of the retinal image on the retina of the observer's eye, O_2. In emmetropia, E, the emergent parallel rays are brought to a focus on the retina of O_2 if the accommodation of this eye is absolutely at rest. In hypermetropia, H, the emergent divergent rays are brought to a focus on the retina of O_2, either by means of accommodation or by placing a convex lens in front of O_2. In myopia, M, the emergent convergent rays can only be brought to a focus on the retina of O_2 by placing a concave lens in front of O_2.

the usual form of astigmatism the image of the disc is an ellipse with the long axis vertical—the opposite of the usual image by the indirect method, with the lens near the eye (*vide* p. 117). It is obvious that there can be no clear image of the whole field by the direct method in astigmatism. *Only lines perpendicular to the meridian which is corrected are seen clearly.* Lines in any meridian other than the two principal ones cannot be seen clearly by any spherical correcting glass, but only by a cylindrical lens or combination of a spherical and a cylindrical.

From what has been said it is obvious that the correcting lens behind the sight-hole of the mirror also represents the spectacle glass which will be required to correct the refraction if it is placed in the same position. This, of course, is only true if the observer is emmetropic, or has his refraction corrected, and if his accommodation is quite at rest. Since it is difficult to relax the accommodation completely, this method of determining any error of refraction should only be used as a rough estimate, unless the observer is an expert. Some surgeons attain great skill in measuring the refraction by the direct method. They can even correct astigmatism. In this case they correct one meridian first, which gives the lens correcting the meridian at right angles to it ; the process is then repeated in the other meridian.

If there is a difference in level between two points on the fundus, *e.g.*, the edge of the disc and the bottom of a glaucoma cup, it is made manifest by the direct method also by parallactic displacement. If we focus the edge of the disc and then move slightly to one side the edge of the disc will appear to move over the bottom of the cup, *i.e.*, it moves in the opposite direction. An object farther forward, therefore, moves in the opposite direction to the movement of the observer's head.

The difference in level can be accurately measured. In the example given, the bottom of the disc will be relatively myopic to the edge, since it is farther away from the back of the lens. If the eye is emmetropic and the edge of the disc can be seen clearly without the assistance of any correcting lens we shall require a concave lens to see the vessels at the bottom of the cup clearly. It can be proved that if the correcting lens is at the anterior focus of the eye a difference of 3 D is equivalent to 1 mm. difference of level. We must get as close as possible to the eye when measuring differences of level, because only then are the conditions of accuracy fulfilled. If, with the observer's accommodation at rest, he sees the edge of the disc clearly with no correcting lens, but requires — 3 D to see the bottom clearly, he knows that the bottom is 1 mm. behind the edge. He will of course see the bottom clearly with a higher concave lens if he counteracts the excess by accommodating : hence he must be careful to choose the lowest minus lens.

Similarly projections forwards can be measured. Here the observer chooses the highest convex lens with which he can see some well-defined point on the top of the eminence. The

same rule that 3 D is equivalent to 1 mm. holds good. Suppose, for example, that he is measuring the swelling of the disc in a case of optic neuritis. He first finds the highest convex glass with which he can see clearly a retinal vessel a little distance away from the disc. He then finds the highest convex glass with which he can see a vessel or a small hæmorrhage as near the top of the swollen disc as he can judge. The difference between the two lenses will give the height of the swelling.

It will be seen that the difficulty of relaxing the accommodation enters into this estimation. The student need not, however, be dismayed. Very fairly accurate results can be obtained without relaxing the accommodation, for if he is accommodating the same amount when he measures the top of the swelling that he is when he measures the level of the surrounding fundus the difference between the two observations will be the same as if he was not accommodating at all. He can ensure this fairly well by always choosing the highest convex lens ; it is safest to choose the lens which just makes the object looked at appear a little blurred. Of course the eye may be myopic ; if for " convex lens " in the above description " relatively convex lens " be substituted (— 1 D being relatively convex to — 4 D) the principle is the same.

An object in the vitreous, *e.g.*, a large opacity, is in the same condition as the fundus of a hypermetropic eye. If the eye is emmetropic, so that the fundus is visible without any correcting lens, the opacity can be examined either by accommodating for it or by putting up convex lenses until it is clearly focussed. If it is close behind the lens accommodation will have to be assisted by a convex lens in any case, unless the observer withdraws farther from the eye. It will be seen, therefore, that by putting up convex lenses from 0 to + 20 D we can thoroughly explore the emmetropic eye from the fundus to the surface of the cornea.

Examined in this manner the appearance of opacities in the vitreous or lens will vary with the amount of light stopped by them, *i.e.*, by their density, and with the amount of light reflected from their surfaces. If they are very dense they will appear black against the background of the red reflex. If they are semi-transparent they will appear red or whitish according to the relative amounts of light transmitted from the fundus and reflected from the surface. A detached retina may therefore look red or white, according to its degree of transparence. If much light is reflected from the surface details may be seen upon it ; otherwise it appears uniformly black.

FIG. 1.



FIG. 2.

PLATE III. (*To face p.* 117.)

THE NORMAL FUNDUS.

Fig. 1.—Appearance of a typical disc, macula, and vessels ; slightly marked physiological cup, showing stippling of lamina cribrosa.

Fig. 2.—A "tigroid" fundus, showing appearances due to excessive choroidal pigmentation or defective pigment in the retinal epithelium.

CHAPTER VIII.

THE FUNDUS OCULI.

When the fundus is observed by the indirect method it is seen to be of a bright red colour. This is due chiefly to the blood circulating in the choroid. In people of dark complexion no choroidal blood vessels are seen on account of the retinal pigment epithelium, which, while dense enough to blur any details, is not sufficiently so to prevent the colour of the blood manifesting itself.

The Optic Disc.—The first object to be sought is the optic disc or papilla (Plate III., Fig. 1). As already mentioned, it is done by making the patient look slightly towards the nasal side. The reflex then suddenly changes from bright red to white, and if the optical conditions are properly arranged in accordance with the directions given in the last chapter the disc will be clearly seen.

The disc is pale pink in colour, the tint showing considerable variations within the limits of normality. It is nearly circular, but seldom perfectly so ; it is about 1·5 mm. in diameter, but of course is seen magnified. The oval appearance due to astigmatism must be borne in mind (*vide* p. 114). The edges are usually quite sharp, but sometimes a little irregular. Not uncommonly, especially in old people, there is a narrow white ring around the pink disc, the *scleral ring ;* this is due to the choroid and the pigment epithelium of the retina not extending quite up to the margin of the disc so that the sclerotic is seen through the retina. Sometimes there is a ring of black pigment around the margin of the disc, due to the retinal pigment epithelium being heaped up here. More commonly parts of the circumference have black patches, but they are not continuous. These features are of no importance from the pathological point of view.

The disc itself is seldom uniformly pink. The central part is usually paler and may be quite white, and this lighter area may extend nearly to the temporal edge of the disc ; it rarely extends quite to the edge. The temporal side is therefore normally paler than the nasal. The central vessels emerge

from the middle of this white area, and careful examination with the direct method will show that the area is a funnel-shaped depression, the *physiological cup*. This cup varies very much in different eyes. When it is very deep the central part may be seen to be speckled with grey spots; these are the meshes of the lamina cribrosa through which the nerve fibres are passing. Sometimes there is scarcely any physiological cup; the disc is then more uniformly pink, and the central vessels may have already divided before they come to the surface. The true nature of the physiological cup is best understood by comparing the ophthalmoscopic picture with a microscopic section vertically through the nerve head.

The colour of the disc is due to the white fibres of the lamina cribrosa seen through the vascularised nerve fibres. Where the nerve fibres are thinnest, *i.e.*, in the cup, the white lamina shines through brightest. The grey spots in the lamina, when they are seen, are due to the non-medullated nerve fibres reflecting less light than the white connective tissue fibres.

The Retinal Vessels.—The retinal vessels are derived from the central artery and vein, which usually divide into two branches at or near the surface of the disc. These branches are above and below, and form a superior and an inferior trunk (Plate I.). Each trunk usually divides into two, one of which sweeps up or down towards the temporal side, the other sweeping up or down towards the nasal side. These branches are called the superior and inferior temporal and nasal arteries and veins. They divide dichotomously into innumerable branches.

The arteries are distinguished from the veins in being lighter red and narrower. The veins have a purplish tint and are often more convoluted; less frequently the arteries are tortuous. What is seen is the blood column, not the actual vessel wall, which is transparent. Each, but especially the arteries, may have a bright silvery streak running longitudinally down the centre, due to reflection of light from the convex cylindrical surface.

The mode of branching of the vessels is subject to great variation, though it is derived from the fundamental type described. The variations are generally of no practical importance. The primary division of the superior and inferior trunks usually takes place on or very near the disc. The nasal branches run much more radially than the temporal, which make a very decided sweep to avoid the macula.

The Macula lutea is situated about 3 mm. or 2 disc-diameters (2 p.d.) to the temporal side of the edge of the disc, and is a little below the level of the horizontal diameter. It is very difficult to see without a mydriatic, for the bright light on this most sensitive spot causes maximal constriction of the pupil: the corneal reflex then usually obliterates all view. It may generally be seen by using very dim illumination.

The macula varies in appearance according to illumination, refraction, complexion, &c. In general, it is a small circular area of a deeper red than the surrounding fundus, sometimes looking almost black. There is nearly always a *foveal reflex*, due to reflection of light from the walls of the foveal depression. This is most frequently seen as a silvery ring of light, hiding everything behind it: it may be circular or oval, according to the incidence of the light and the refraction of the eye. Often there is an intensely bright spot at or close to the fovea, also due to reflection. The deeper red of the macula is due to the thinness of the retina, so that the specially vascular choriocapillaris of this region is seen more clearly: shadows thrown by the edges of the foveal depression may contribute a share to the deepening of the colour.



Fig. 88.—A cilio-retinal artery.

The macular region is supplied by twigs from the superior and inferior temporal arteries, and by small branches coming straight from the disc. There are no retinal blood vessels actually at the fovea (Fig. 5), and none can be seen ophthalmoscopically for a little distance around. Occasionally there are small arteries (cilio-retinal) derived from the ciliary system. They start near the edge of the disc, run inwards, and then bend sharply outwards towards the macula (Fig. 88).

The General Fundus. The appearance of the general fundus varies enormously within healthy limits. It is especially determined by the complexion of the patient, which may be taken as an index of pigmentation in different parts of the body. In people who are neither very dark nor very light the spaces between the retinal vessels show a uniform redness, occasionally with a very delicate punctate stippling, especially

towards the periphery. In albinos the choroidal vessels are seen clearly, the spaces between them being white, due to the sclerotic shining through. In very dark people the fundus is a darker red, and indications of the choroidal vessels are often seen as indefinite brighter red streaks. Sometimes the pigment between the choroidal vessels is particularly dense, or the pigment is deficient in the retinal pigment epithelium, while the choroid is deeply pigmented : the choroidal vessels are then seen separated by deeply pigmented polygonal areas (*tigroid* or *tesselated fundus*) (Plate III., Fig. 2).

There is no difficulty in distinguishing the choroidal from the retinal vessels when both are visible (*cf.* Plate X., Fig. 2). The former are broader and ribbon-like, without any central reflex streak : they anastomose freely, unlike the retinal vessels, which do not anastomose at all. Moreover in certain parts, their anatomical distribution is very characteristic (*vide* p. 11).

All the details of the fundus will be seen much better by the direct method, unless the eye is very myopic, when the magnification is so great and the area seen is so small that it is difficult to find any given spot.

Under normal conditions no pulsation can be seen in the retinal arteries. The retinal veins, however, may often be seen to pulsate at or near the edge of the disc, or indeed wherever they take a very sharp bend. This is usually due to transmitted pressure. The blood pressure is lowest in the veins near the disc, and there is a certain amount of obstruction to the flow of blood as the vessels pass through the narrow neck at the lamina cribrosa. With each arterial pulsation the intraocular pressure is suddenly raised slightly, so that the pressure on the outside of the walls of the veins is increased. This causes a sudden increased obstruction to the outflow of blood from the eye ; the wall of the vein becomes slightly compressed, recovering itself during the arterial diastole. Hence pulsation is observed, and it will be seen best where the intravenous pressure is least, viz., nearest the heart, *i.e.*, at the disc, and where there is any additional obstruction, viz., near the lamina cribrosa and at any sharp bend. The venous pulsation can be increased or made manifest if absent by slight pressure on the globe, which has the effect of increasing the intraocular pressure. This normal venous pulse is seen without the artificial aid of pressure on the globe in 70 to 80 per cent. of people. It will be noticed that it is diastolic ; it has therefore been called the *negative venous pulse*.

Two other forms of venous pulse occur in pathological conditions.

The *positive venous pulse* is presystolic, continuing into the systolic phase: it is due to tricuspid regurgitation, and is permitted by the normal insufficiency or absence of valves in the jugular veins. The *transmitted centripetal venous pulse* is an accentuation of the normal tendency of the pulse wave to progress through the capillaries into the veins, owing to the intraocular tension. It is due to venous congestion, with or without increased vis a tergo.

Visible arterial pulsation is always pathological. The blood pressure in the ophthalmic artery is only a few mm. Hg. below that of the carotid in animals. Considering the differences of blood supply it would be unwise to apply the result directly to man, but there is no doubt that the pressure in the central artery is far above the intraocular pressure. It would not be surprising, therefore, if the pulse wave were transmitted and could be seen. There are two reasons which militate against this: (1) the intraocular pressure damps the pulsation, and the increase in pressure which accompanies each pulsation is spread over the whole volume of the contents of the globe, and is transmitted to the plastic sclerotic; (2) such pulsations as survive this damping effect are too slight to be observed in such small vessels by ordinary ophthalmoscopic examination.

Two types of arterial pulsation occur pathologically: (1) a true pulse wave, accompanied by locomotion of the vessels; (2) an intermittent flow of blood, or pressure pulse. In the latter the arteries fill only with the heart beats, being empty between them; it is only visible on the disc, and may be produced in a normal eye by external pressure upon the globe by a finger applied to the lid. This type of pulsation is a pure pressure phenomenon, and is caused by any considerable increase of intraocular tension with normal or lowered blood pressure, *e.g.*, in glaucoma, or by any considerable diminution of blood pressure with normal intraocular pressure, *e.g.*, in syncope, orbital tumours, &c. The true arterial pulse occurs in cases of aortic regurgitation or aneurysm, in exophthalmic goître, &c.; it is not confined to the disc. It is equally a pressure phenomenon, but the differences of pressure are smaller.

Capillary pulsation is seen only in aortic regurgitation as a systolic reddening and diastolic paling of the disc.

The order of examination of the details of the fundus should

be systematic. Applying the indirect method we obtain a general view. The patient is instructed to fix the gaze in such a direction that the disc is brought into view. It will occupy about the centre of the field, and a considerable area around will be visible. Any gross abnormality is detected at once. The shape and colour of the disc, the arrangement of the vessels, the colour of the choroidal reflex (its uniformity or tesselation), gross abnormalities (white or pigmented spots, &c.), are readily noted. The patient is then directed to look up to the ceiling, to the right, to the left, and down to the ground; in the latter position the lid is gently raised by a finger of the hand which is holding the large lens, as otherwise it will cover the pupillary area. In this manner the periphery of the fundus is brought into view. Even when the central parts of the fundus are uniformly tinted the periphery often displays traces of the choroidal vessels, associated with greater pigmentary stippling or a diminution of pigment. Only minute investigation with the direct method can show whether this is normal or pathological. The characteristic type and distribution of the pigmentation of retinitis pigmentosa is best demonstrated in this manner.

Having thus obtained a good general idea the systematic examination is repeated by the direct method, paying special attention to points which the indirect method has left uncertain. The details of the disc—physiological or pathological cupping, blurring of the edges or swelling, abnormalities of the edges in the form of crescents and so on—are inspected. Attention is then turned to the vessels. Abnormalities in arrangement or distribution, the presence of cilio-retinal vessels, &c., are noted. The details of the individual vessels—their relative size, irregularity of contour or varicosity, visibility of the walls as shown by the presence of white lines along the edges, abnormalities of the reflex streak, &c.—are investigated. The vessels are traced towards the periphery and the smaller vessels inspected. Changes often occur near the vessels, such as small hæmorrhages, white spots of exudate, &c.; these are carefully looked for.

Next, the macula is examined: this should never be omitted. It may be brought into view by telling the patient to look into the light: with unintelligent patients it is best to say nothing, but fix the temporal edge of the disc and pass horizontally outwards for a distance of about two papilla diameters (a convenient unit in ophthalmoscopic topography), when the macula will be found. If the patient is not under a mydriatic

or the pupil movements are not abolished by disease (optic atrophy, &c.) the light should be lowered so that the constriction of the pupil may be reduced to a minimum. The corneal reflex is always troublesome, but has to be dodged. Any abnormality at or near the macula is of the utmost importance. Black or white spots are often very difficult to distinguish from shadows or reflexes : if either can be made out by any means (*e.g.*, the use of a concave lens) to have a definite sharp contour, and if they do not seem to shift in the slightest degree when a minute movement is made with the mirror, it may be concluded that they are pathological entities.

Finally, the periphery of the fundus is investigated. It is important to know how far out we can see by the direct method. It can be shown theoretically and proved practically that the fundus can usually be seen only a short distance beyond the equator of the eye. There is therefore a zone behind the ora serrata which is invisible. The periphery, even in an emmetropic eye, is usually best seen with a low convex lens, owing to the obliquity of the axis of the rays as they pass through the crystalline lens.

CHAPTER IX.

FUNCTIONAL EXAMINATION.

In the second great group (*vide* p. 74) of ophthalmic patients there are no manifest objective signs of disease, and we are dependent at the outset upon the subjective symptoms of which the patient complains. In these cases, after a rapid, but careful, external examination to eliminate any objective sign which may have escaped observation, it is usually most convenient to proceed at once to the functional or subjective examination. Whether this produces evidence of abnormality or not, it is imperative to proceed then to the systematic internal examination with the ophthalmoscope.

On the other hand, in the first group of ophthalmic patients, in spite of external signs of disease which may suffice to account for the symptoms, it is the surgeon's duty to eliminate as far as possible all other evidence of abnormality. Ophthalmoscopic examination may be impossible at the first visit, and subjective functional testing may be so vitiated by the obvious disease as to afford no useful information. In such cases functional and internal examination should be postponed until a future visit, but in no case should it be entirely omitted. It should be borne in mind that, rightly or wrongly, the surgeon is liable to be held responsible for the discovery of any disorder of the eye which manifests itself at or about the time of his examination.

The functional examination of the eye proper consists in testing the acuity of the forms of visual perception which have been already mentioned, viz., the light sense, the colour sense, and the sense of form. They are usually tested in the reverse order. Each eye must be tested separately throughout. In the preliminary examination it is sufficient to test first the acuity of central vision for distant and for near points, then roughly the field of vision, and finally the reactions of the pupils, afterwards proceeding forthwith to the ophthalmoscopic examination. It may then be necessary to revise the rough test of the field of vision by taking it with greater precision, to supplement the tests with white light by those with coloured

objects, both in the central and peripheral parts of the field, and to estimate, more or less accurately, the light minimum and the light difference. In the routine examination it is well to test roughly the condition of the extrinsic muscles before proceeding to the ophthalmoscopic examination (Section V.).

The Acuity of Vision.—The acuity of distant central vision is now almost invariably tested by means of Snellen's Test Types (Fig. 89). These are constructed upon the fundamental principle that the average minimum visual angle is 1 minute.

The types consist of a series of letters diminishing in size. The breadth of the lines of which the letters are composed is such that the edges subtend an angle of 1 minute at the nodal point of the eye. Each letter is of such a shape that it can be placed in a square, the sides of which are five times the breadth of the constituent lines (Fig. 89). Hence the whole letter will subtend an angle of 5 minutes at the nodal point of the eye (Fig. 90).



FIG. 89.—Snellen's Distant Test Types (reduced). The lines, from above downwards, should be read at 60 metres, 36 metres, 24 metres, 18 metres, 12 metres, 9 metres, and 6 metres respectively, *i.e.*, at these distances the letters subtend a visual angle of 5′.

In order that these conditions may be fulfilled it is obvious that such a letter to be used as a test a long distance from the eye must be larger, and the constituent lines must be broader than in the case of a letter to be used nearer the eye. In Snellen's types the largest letter will subtend 5 minutes at the nodal

point if it is 60 metres from the eye. Those in the second line will subtend 5 minutes if they are 36 metres from the eye ; those of the consecutive lines 24 metres, 18 metres, 12 metres, 9 metres, and 6 metres. Sometimes smaller letters corresponding to 5 metres, 4 metres, 3·5 metres are used.

A person with average acuity of vision ought therefore to be able to read the top letter at 60 metres, the second line at 36 metres, the third at 24 metres, and so on. Now, it would be very inconvenient to have to alter the distance between the patient and the letters to this large extent. A numerical convention which gives a fair comparative estimate of the value of different acuities of vision has therefore been adopted.

The patient is kept at a fixed distance from the types. This distance should never be less than 5 metres, or preferably 6 metres. At such a distance the divergence of the rays in the small bundle which enters the pupil is so slight that it



Fig. 90.

can be neglected, the rays being considered parallel. If the distance were 3 metres, for example, an appreciable amount of accommodation would have to be exerted by an emmetropic eye in order to bring the rays to a focus upon the retina : hence the estimate of distant vision would be fallacious.

A normal patient 6 metres from the types ought to be able to read every letter from the top to the end of the 6 metre line ; many people can read more in a good light. Suppose the patient can only read the 18 metre line. His distant vision is obviously defective. The numerical convention which is used to record this is a fraction in which the numerator is the distance at which he is from the types, and the denominator is the distance at which a person with normal vision ought to be able to read the last line which he succeeds in reading. The patient under consideration will therefore have his distant vision recorded thus : $V = \frac{6}{18}$. The normal patient's vision will be $V = \frac{6}{6}$.

These fractions give an indication that the normal patient's vision is unity, whilst the other patient's is one-third as good. The fraction should not be reduced in this manner, because it is only an accurate numerical estimate under special conditions. It should be used merely as a convention, just as numbers are used to indicate variations in tension, though this is far less defensible. If the fraction is reduced much valuable information is lost. In its original form it indicates the actual types used and the actual distance away from the types: it therefore eliminates doubt as to the accuracy of the application of the test.

The amount of illumination on the test card has a considerable influence on normal visual acuity. It has been found that the acuity rises rapidly as the illumination is increased from zero up to 2-ft. candles: above 2–3-ft. candles the rise is very slight. The illumination of the test card should, therefore, never be allowed to fall below 3-ft. candles, and it would be advantageous if the standard illumination recommended by the Council of British Ophthalmologists were universally adopted.

If the patient cannot read the largest letter he is told to walk slowly towards the types. At a certain distance he may be able to see the top letter. He should then be moved back a little, since he may not have understood exactly where to look. In this manner the farthest point at which he can distinguish the top letter is determined. If this is 3 metres, the vision is recorded thus—V = 3/60. Perhaps he is unable to see the top letter even close to it. In that case he is asked to count the extended fingers of the surgeon's hand, held up at about 1 metre against a dark background: the distance is varied to obtain about the maximum. This is recorded thus—V = fingers at 1 metre. If he cannot count fingers he is told to look at the light, either artificial or the window; the surgeon's hand is then moved between the eye and the light. If he can distinguish the movements of the hand it is recorded as V = hand movements. If he is unable to distinguish hand movements he is taken into the dark room and the light is alternately switched on and off, or light is concentrated on his eye with a convex lens or with the ophthalmoscope mirror, and he is asked to say when the light is on the eye and when it is off. If he succeeds in doing this, V = p. l. (perception of light). If he fails to see the light at all the vision is recorded as V = no p. l.

It does not follow that a patient who reads 6/6 is emme-

tropic : he cannot be myopic unless he is screwing up his eyes, and in any case he cannot be very myopic. He may, however, be hypermetropic, for by an effort of accommodation he can bring the practically parallel rays emitted by the letters to a focus on his retina. We wish, then, to find out if he is accommodating. It is done by finding out the highest convex lens, placed before his eye in a testing spectacle frame, with which he can still read 6/6.

Directly a convex lens is placed before the eye in a patient who has good accommodative power it tends to make objects look blurred. Hence it is best to start with a convex lens which will quite definitely blur the types, and then gradually counteract it by placing concave lenses of increasing strength in contact with it. A + 4 D lens is put in front of the eye. Probably the patient now reads only a few lines. With the + 4 D still in position a — 0·5 D is put in front of it ; the patient perhaps reads another line. The — 0·5 D is replaced by a — 1 D ; he still fails to read 6/6. The — 1 D is replaced by — 1·5 D ; we will suppose that he now reads 6/6. This proves that he has average normal vision with a + 2·5 D lens. With the assistance, therefore, of a 2·5 D convex lens he can relax his accommodation the corresponding amount. It by no means follows that this represents the total amount of his hypermetropia. As has already been pointed out (*vide* p. 48), the younger the person the greater the capacity to accommodate. Young people, therefore, have great facility in accommodating. If they are hypermetropic this fund has been drawn upon for a long period, so that a condition of partial contraction of the ciliary muscle becomes normal to them. Sometimes even there is excessive contraction of the muscle, so that they become artificially myopic, a condition which is called " spasm of accommodation " ; it is probably diagnosed more often than it occurs.

The younger the patient, therefore, the easier it will be for him to accommodate, and the more difficult to relax his accommodation completely. That part which he can relax when convex lenses are used as described above is called his *manifest hypermetropia* (Hm.). The part which he is unable to relax, which can only be determined by paralysing the ciliary muscle, is called his *latent hypermetropia* (Hl.). The sum of the manifest and latent hypermetropia is called the *total hypermetropia*. In extreme youth nearly all the hypermetropia is latent : the lens is so resilient that it is impossible to prevent it responding to the slightest stimulus. As the lens becomes less plastic

more and more of the hypermetropia becomes manifest, until, finally, when accommodation disappears entirely, all the hypermetropia is manifest. The older the patient, therefore, the nearer the manifest hypermetropia represents the total amount.

The vision of the patient in the above example is usually recorded thus : $V = \frac{6}{6}$, Hm. + 2·5.

With intelligent patients who do their best to read as many letters as possible without continual encouragement from the surgeon, the manifest hypermetropia is obtained with less trouble by simply putting up convex lenses of gradually increasing strength until the last line which can be read becomes blurred.

An older patient than the one considered in the previous example will very likely read more with a convex glass than without it. Thus a patient of fifty-five may perhaps read only 6/12, whilst with a + 2 D he reads 6/6. This man has a manifest hypermetropia of 2 D. Since he is fifty-five years of age he has only 1 D of accommodation left (*vide* p. 50). When he reads with the unaided eye he uses up this 1 D in getting as far as 6/12 ; he is unable to accommodate any more, so the lower letters are too blurred to read. He may manage to read 6/6 with the assistance of a + 1 D, since this, with the remnant of his accommodation, will fully correct his hypermetropia. On seeking the highest convex glass we find he can read 6/6 as well or better with + 2 D. This, therefore, represents his manifest hypermetropia. Such a case is recorded—$V = \frac{6}{12}$, Hm. + 2 = $\frac{6}{6}$.

Apart from counteracting a convex lens as described above the student should never use concave lenses in testing the distant vision, unless the patient is under a mydriatic. An emmetrope or even a hypermetrope, if neither is presbyopic, will read 6/6 quite well with weak concave lenses in front of the eye ; he simply accommodates the amount which is requisite to counteract the lens. Hence, unless the patient is under the influence of a mydriatic, we learn nothing from the procedure ; it is therefore redundant.

Having tested the distant vision and determined the amount, if any, of the manifest hypermetropia, the near vision should next be tested. For this purpose test types for near vision are used (Fig. 91). Snellen's are constructed on exactly the same principle as the distant ones, and are therefore most

scientifically accurate. For actual reading, however, people can often read types which are smaller than Snellen's smallest if the lines forming the letters are a little broader. This is chiefly due to the fact that an idea of the shape of letters will often permit them to be accurately guessed when they are not clearly visible. This is, indeed, an objection to the use of letters as test objects at all. Jaeger's near test types, which are very frequently used, are simply the ordinary printers' founts of type, from the smallest upwards (nonpareil, minion, &c.). They are sufficiently accurate for practical purposes.

The patient is told to hold the test card. The position where he holds it of his own accord will often impart useful information. If he is old and holds it a long distance away, he is most likely to be an emmetropic or hypermetropic presbyope. If he holds it closer than the ordinary reading distance and reads the smallest type fluently, he is probably myopic, whatever his age may be, though children often get into the habit of holding books unnecessarily close.

Take first the example of an emmetrope. We find that he reads 6/6, that he has no manifest hypermetropia, and that he reads Jaeger 1, holding it of his own accord at ordinary reading distance (22 cm. or 9″). This is recorded $-V = 6/6$, no Hm., J. 1. If no distance is stated in recording the near vision it is understood to be the normal distance.

Take now a patient who reads only 6/60, has no manifest hypermetropia, but reads Jaeger 1 fluently, only, however, when the card is held much closer than normal to the eye. In this case the distance at which the card is held should be guessed or measured. Let us suppose that it is about 5 inches. This is recorded—$V = 6/60$, no Hm., J. 1 at 5″.

Take now a patient of fifty-five who reads 6/6, and has no manifest hypermetropia. We give him the near types, and he holds the card a long distance off, but even so cannot read Jaeger 1. In this case it is waste of time to discover exactly which type he can read, and to measure the distance at which he can read it. We know that as he is fifty-five he has only 1 D of accommodation remaining (*vide* p. 50). What we wish to know is whether he can read Jaeger 1 at ordinary reading distance if we correct his presbyopic defect. We therefore at once put up a + 3 D lens before the eye, tell him to hold the types closer, and ask him if he can read the smallest. He will probably do so easily. This is recorded—$V = 6/6$, no Hm., $\bar{c} + 3 =$ J. 1.

Finally, take a patient of fifty who reads 6/12, but with

J. 1 (Sn. 0.5). 50 cm.

As she spoke, Moses came slowly on foot, and sweating under the deal box, which he had strapt round his shoulders like a pedlar. "Welcome, welcome, Moses! well, my boy, what have you brought us from the fair?"—"I have brought you myself" cried Moses, with a sly look, and resting the box on the dresser. "Ay, Moses," cried my wife, "that we

J. 2 (Sn. 0.6). 60 cm.

five shillings and twopence is no bad day's work. Come, let us have it then."—"I have brought back no money," cried Moses again. "I have laid it all out in a bargain, and here it is," pulling out a bundle from his breast: "here they are; a gross of green spectacles, with silver rims and

J. 4 (Sn. 0.8). 80 cm.

mother," cried the boy, "why won't you listen to reason? I had them a dead bargain, or I should not have brought them. The silver rims alone will sell for double the money."—"A fig for the silver rims," cried my wife, in a passion: "I dare

J. 6 (Sn. 1). 1 m.

the rims, for they are not worth sixpence; for I perceive they are only copper varnished over."—"What!" cried my wife, "not silver! the rims not silver?"—"No," cried I, "no more silver

J. 8 (Sn. 1.25). 1.25 m.

with copper rims and shagreen cases? A murrain take such trumpery! The blockhead has been imposed upon, and should have known his company better."—"There,

J. 10 (Sn. 1.5). 1.5 m.

the idiot!" returned she, "to bring me such stuff: if I had them I would throw them in the fire."—"There again you are wrong, my dear," cried I;

J. 12 (Sn. 1.75). 1.75 m.

By this time the unfortunate Moses was undeceived. He now saw that he had

J. 14 (Sn. 2.25). 2.25 m.

asked the circumstances of his deception. He sold the horse, it

FIG. 91.—TEST-TYPES FOR NEAR VISION.

Jaeger Test-Types, with approximate Snellen equivalents, and the most remote distances at which each should be read with average normal vision.

+ 2 D reads 6/6. He will hold the near types a long distance away as in the last example; if we investigate the question he will not be able to read nearly as well even a long distance away as the last patient. He has only 2 D of his accommodation remaining, but he also has 2 D of hypermetropia. We cannot therefore expect him to read Jaeger 1 at ordinary reading distance unless we not only correct his presbyopia but also his hypermetropia. We therefore at once put up a + 4 D, and find that he reads Jaeger 1 at ordinary distance quite well. This is recorded—V = 6/12, Hm. + 2 = 6/6, $\bar{c}$ + 4 = J. 1.

The ordinary rule of presbyopic loss of accommodation, viz., 1 D for each five years after forty, is rather a liberal allowance, and we often find that patients are more comfortable with slightly less (*vide* p. 50). Only in very exceptional cases should more be ordered.

An indication of the *range of accommodation* is given by the knowledge of the manifest hypermetropia, combined with the ability to read the small types at ordinary reading distance. Strictly, the accommodation should be more carefully tested in each case, but it is often neglected.

The method adopted to find the near point of the eye has already been mentioned (*vide* p. 46). For practical purposes it is sufficient to use the smallest Jaeger or Snellen near type and approach it nearer and nearer to the eye until it can no longer be read. The last point at which it can be read gives the near point. The distance of the near point from the eye is then measured with a tape. This distance is transformed, if necessary, into millimetres (25 mm. = 1 inch), and the range of accommodation is deduced from the formula A = P R (*vide* p. 47). Of course, the full range of accommodation in a hypermetrope cannot be accurately arrived at unless the total hypermetropia is known; this may require the use of a mydriatic. Practically, however, we are chiefly concerned in discovering paralysis or paresis of accommodation, such as may occur after diphtheria or previous use of a mydriatic. In these cases the knowledge of the distance of the near point is sufficient.

The next step—one which is far too often neglected—is in every case to test the *pupil reactions* and record them. If the visual tests have shown deficiency it may be necessary to use a mydriatic, in which case it will be impossible to test the pupil reactions at a later stage of the same visit; hence the importance of recording them at once.

We should also test the field of vision roughly.

The Field of Vision.—There are several methods of testing the field of vision.

(1) A rough, but very useful, method which should be applied in every case, at any rate if there is the slightest suspicion of defect, is as follows :—

The surgeon stands facing the patient at a distance of 18 inches to 2 feet. The patient covers his left eye with the palm of his hand. He is told to look straight into the surgeon's left eye. The surgeon closes his right eye. He then moves his hand in from the periphery towards the common line of vision of the patient's right and his own left eye, keeping his hand in the plane half-way between the patient and himself. Directly he sees it himself the patient ought to say that he also sees it. The movements of the hand are repeated in various parts of the field—above, below, to the right, to the left, and so on.

This method is extremely simple, rapidly applied, and an excellent test. It will be seen that the surgeon tests the range of the patient's field by that of his own, which may be considered normal ; moreover, he is continually watching the patient's eye, so that he can at once observe any deflection from the point of fixation.

The gross defects in the field which are most likely to escape recognition are homonymous and bitemporal hemianopia, the latter usually due to acromegaly or tumour of the pituitary body (Chap. XIX.). They may be roughly tested for by telling the patient to look straight at the surgeon, situated as before, both eyes being open. The surgeon holds up both hands, one in each temporal field, and the patient is told to touch the surgeon's hand. If he asks " Which one ? " he has not bitemporal hemianopia, since he sees both hands. If he promptly points to one hand he should be asked if he sees the other ; if he does not, he probably has homonymous hemianopia.

If any defect is indicated by these methods or is suspected from other features of the case it must be accurately mapped out and recorded with the perimeter.

(2) *The Perimeter*:—The perimeter consists essentially of an arc, marked on the back in degrees of a circle, capable of being revolved round a pivot which the patient fixes with the eye under examination (Fig. 92). The chart, which has concentric circles marked upon it, corresponding with the degrees on the arc, is under the surgeon's control at the back of the

perimeter. In self-registering perimeters, which are almost invariably used, the readings are recorded by perforations with a sharp point.

The details of taking a perimetric chart can only be taught by actual demonstration. It will suffice to emphasise here



Fig. 92.—McHardy's perimeter.

the chief procedures to be followed in order that accuracy may be attained.

The patient is seated with his back to the light. His chin rests upon the chin-rest; the face is vertical and not tilted to one side; one eye is covered. The other eye, situated at the centre of the arc, fixes the white dot around which the arc revolves.

The field should first be taken with a white object 10 mm. in diameter. At least eight meridians must be investigated,

preferably sixteen. If the field is very small a 5 mm. square should be substituted, and the process repeated. In each meridian the object should be carried quite up to the fixation point, as there may be areas inside the limits of the field which are blind (*absolute scotomata*). These should be mapped out with the same accuracy as the extreme limits of the field. If the scotomata are small the limits may be determined with an intelligent patient best with a very small object, *e.g.*, 2 mm. square. The size of the test object and the distance from the perimeter may be conveniently recorded by a convention similar to the mode of recording visual acuity, *e.g.*, 10/300, both distances being expressed in millimetres (A. H. H. Sinclair). With small objects *relative scotomata* can be found which are not demonstrable with large objects.

Having mapped out the field for white the process should be repeated with similar, but coloured, objects. The limit of the field for a colour is the point at which, passing from the periphery to the centre, the colour first becomes evident. Peripheral to this limit the object may still be perceptible but appears grey. The exact limit is difficult to determine, for most colours change in hue and saturation as the object passes from the fixation point towards the periphery. Red or green should be used first, then blue or yellow. Under ordinary circumstances, the blue field is largest, slightly smaller than the white : then follow the yellow, red and green, in the order named (Fig. 93). There is a particular purplish-red and a particular bluish-green (490 $\mu\mu$) which have the same field ; and similarly a particular blue (460 $\mu\mu$) and a particular yellow (570 $\mu\mu$). These pairs of colours are complementary, *i.e.*, a mixture of the red and the green, or of the blue and the yellow, produces white.

The extent of the normal field, with a 10 mm. square, under good illumination, is shown in the accompanying chart (Fig. 93). The peculiar shape is essentially due to the shape of the sensitive area of the retina as projected outwards, but is often modified when the field is taken in the ordinary manner by interference caused by the nose and the brows ; this complication can be eliminated if, when the field of the right eye is being taken, the head is turned somewhat to the left, and *vice versâ*. It is seen that the field for white extends upwards 45°, outwards rather more than 90°, downwards 70°, and inwards 60°. The size varies with the illumination, the size of the test object, the contrast of the test object with the background, and the state of adaptation of the eye. The field for

blue and yellow is roughly 10° less in each direction, that for red and green another 10° less. The limits of the colour fields vary not only with the intensity of the light, but also with the saturation of the colour, and above all the size of the object. If these are sufficiently great colours may be recognised almost, if not quite, at the periphery.

Even the ordinary perimetric observation is a relatively rough test and purely subjective. It must always be regarded



FIG. 93.—Perimeter chart of right eye (Landolt). T, temporal side; N, nasal side; W, for white object; B, for blue; R, for red; G, for green.

with suspicion in dull or neurotic patients. The most variable factor is the illumination, and sufficient attention is not usually paid to this point. With good illumination an object subtending a visual angle of $\frac{1}{2}°$ will give the full formal field for white. The ordinary 10 mm. object at the distance generally used, *viz.*, 30 cm., *i.e.*, 10/300, corresponds with a visual angle of 2°. Deductions made from variations in the colour fields are particularly unreliable (*vide* p. 135).

Special care must be taken to investigate the central part of the field for red and green, since conditions are not uncommon, *e.g.*, tobacco amblyopia and retrobulbar neuritis, in

which these colours are not recognised by central vision (*central relative scotomata*). The 5 mm. square should be placed over the point of fixation and the colour changed: blue and yellow will be recognised as such, but not red and green.

(3) For more accurate investigation of details another method must be employed, but it is applicable only to the central and paracentral areas. It consists in placing the patient 2 metres from the centre of a large black screen, 2 metres or more broad (Bjerrum's screen). The patient fixes a spot in the centre of the screen and smaller circular discs of ivory, 1 mm. to 10 mm. in diameter, attached to a long black



FIG. 94.—Diagram of the right eye, showing the relationship of the retina to the degrees of the perimetric arc, and the relative value of the latter when projected on a tangent scale. *n*, nodal point. F, point of fixation.

rod are brought in from the periphery on a level with the screen. At this distance a 3 mm. object subtends a visual angle of about 5 minutes. It will be noticed that, the angles being projected on to a flat surface, tangents are recorded, not angles themselves as with the arc. Hence only a small area can be investigated, and the distortion must be taken into account. Some points of diagnostic importance which cannot be elicited by the perimeter can be brought out by this method. Various *scotometers* have been devised on the principle of Bjerrum's screen.

If the charts of the two eyes are superposed there will be a large central area which is common to both eyes: this is the *field of binocular vision*.

The Light Sense may be roughly tested by means of Bjerrum's test types, which consist of Snellen's test types printed grey on a grey background of different intensity; or it may be tested against the surgeon's as a control by using the ordinary Snellen's test types and gradually lowering the illumination. No simple means of accurately testing the light sense has yet been devised, most of the clinical photometers, such as Förster's, being very unsatisfactory.

The Colour Sense requires elaborate apparatus for its scientific investigation. The methods used will be discussed later (p. 380).

SECTION III.

DISEASES OF THE EYE.

CHAPTER X.

DISEASES OF THE CONJUNCTIVA.

The conjunctiva shows very considerable variations in appearance at different ages and in people who follow various employments. The peculiarities of colour, vascularity, laxity, &c., which are consistent with health can be learnt only by repeated observation.

It is necessary for a scientific appreciation of pathological conditions to be cognisant of the normal structure of a part. The conjunctiva is divided into two portions, palpebral and bulbar: the folds uniting these parts are the fornices. The palpebral conjunctiva is said to commence at the anterior margin of the edge of the lid, but from this point to the posterior margin of the edge (the intermarginal strip) and for about 2 mm. beyond (to the sulcus subtarsalis) there is a transitional zone covered with stratified epithelium and partaking of the characters of both skin and conjunctiva (Chap. XXXI.). There are two layers of epithelium over the palpebral conjunctiva: from the fornices to the limbus the epithelium becomes gradually thicker, forming eventually again a stratified epithelium. Below the epithelium is an adenoid layer, consisting of loose connective tissue containing mononuclear lymphocytes: below this a fibrous layer, much denser and passing insensibly into the underlying tissues—lid or sclerotic. The palpebral conjunctiva is firmly adherent to the tarsus, while the bulbar portion is freely movable over the sclerotic except close to the cornea.

Bacteriology.—The conjunctival sac is practically never free from organisms. Owing to the relatively low tempera-

ture of the conjunctival sac due to exposure, evaporation of lacrymal fluid and moderate blood supply, bacteria do not propagate themselves readily. The tears are not a good culture medium, though since they contain no hæmolysins or bacteriolysins, they cannot be regarded as actively bactericidal. They contain no agglutinin, and diphtheria antitoxin does not pass into them when present in the blood. Hence they act principally in a mechanical manner, washing away deleterious agents and their products. The presence of dust, even if it be sterilised, augments the bacterial content of the conjunctival sac. It is also increased by bandaging owing to the arrest of movement of the lids and the raised temperature of the sac. Most of the organisms normally present are non-pathogenic, but some of these are morphologically identical with pathogenic organisms. Diplococci indistinguishable from pneumococci are found; they may be innocuous to animals or prove themselves true pneumococci. The conjunctiva of many people is immune to pneumococcic infection, though the same germs transferred to other persons will excite a violent inflammation. It may be stated at once that the pneumococcus is one of the most dangerous organisms in the pathology of the eye. Another bacterium, the so-called xerosis bacillus, is morphologically identical with the diphtheria bacillus; it can only be distinguished by skilled examination of cultures. Staphylococci are found: they are relatively innocuous in the absence of other organisms, but play an important part in mixed infections. Staphylococcus albus and xerosis bacilli are practically always present in the normal conjunctival sac. There are reasons for thinking that their presence favours the growth of pathogenic bacteria: thus xerosis bacilli promote the multiplication of Koch-Weeks' bacilli both in cultures and on the conjunctiva. Streptococci, Bac. coli, &c., are pathogenic, but rare. Other pathogenic organisms– gonococci, Koch-Weeks' bacilli, diplobacilli –will be discussed later.

Inflammation of the Conjunctiva.

Conjunctivitis. –Inflammation of the conjunctiva manifests itself in many grades and many types. It is always accompanied by hyperæmia and by increased secretion. The hyperæmia varies in degree and in distribution: the secretion varies in nature and amount.

Hyperæmia may be transitory, or recurrent and chronic. The former is caused by temporary irritation, as by a foreign

body in the conjunctival sac (which includes the surface of the cornea), concretions in the palpebral conjunctiva (*vide* p. 179), in-growing lashes (*vide* p. 565), &c. : in such a case the increased secretion is almost wholly a reflex secretion of tears. A foreign body, especially a grain of corn or the wing capsule of an insect, may be retained in the fornix and set up a violent unilateral conjunctivitis. Irritation limited to the lower fornix may be artificial in malingerers and hysterical patients.

Recurrent or chronic congestion may be caused by the conditions of life—dusty, ill-ventilated rooms, exposure to strong light, &c. Bright light, especially exposure to tropical sunlight, acts partly by the glare due to the luminous rays, partly by the chemical action of the actinic, especially the ultra-violet rays, and partly by the heat, which is chiefly due to the infra-red rays. Chronic congestion is often due to conditions remote from the conjunctiva itself. Very frequently it is a reflex irritation due to errors of refraction : in such cases the edges of the lids may participate. Other causes are found in errors of metabolism—gout, over-eating and drinking, and so on. It is a characteristic symptom of hay fever, and in this case there is an excess of eosinophile leucocytes in the conjunctival secretion.

Simple hyperæmia of the type described causes a sense of discomfort, often described as tightness, grittiness, inability to keep the eyes open, tiredness, &c. Bright light is resented, but there is seldom true photophobia. The conjunctiva often looks quite normal until the lower fornix is exposed, when it will be seen that the parts in contact are congested and sticky. The discomfort frequently comes on only in the evening or after near work. In the gouty cases there may be œdema

chemosis.

Chemosis affects the most loosely attached parts of the conjunctiva, *i.e.*, principally the bulbar conjunctiva and fornices. The mucous membrane becomes swollen and gelatinous in appearance. The swollen membrane forms a wall around the cornea, which it may overhang in severe cases. The palpebral conjunctiva is little affected, but the tissues of the lid are often also œdematous, so that the lids are swollen and the upper hangs down over the lower.

Whenever watering of the eyes is complained of, and whenever only one eye is congested or shows signs of conjunctivitis, the lacrymal passages must be investigated. Pressure with the finger backwards and inwards over the lacrymal sac may

cause regurgitation of fluid—tears, mucus, or pus—showing that the outflow into the nose is obstructed. If no regurgitation can be detected, the position of the lower punctum must be noted. It ought to be invisible until the lid is slightly everted.

The *treatment* of simple hyperæmia consists primarily in the removal of the cause. Defective conditions of life must be ameliorated if possible. The irritation of strong light must be removed, or modified by the use of dark glasses. If the light is not very excessive ordinary neutral tinted (" smoked ") glasses will suffice. They are better than blue or other coloured glasses as they reduce the intensity of the luminous rays more uniformly throughout the spectrum. Nearly all kinds of glasses cut off a large percentage of the ultra violet rays, especially those of shortest wave length. Sir William Crookes has prepared a series of synthetic glasses which have various absorptive powers. Some of these cut off practically all the infra-red and ultra-violet rays, while absorbing the luminous rays to only a slight degree. Tinted Crookes' glasses are specially indicated for use in tropical climates.

Errors of refraction must be corrected. It should be remembered that the error may be artificial, through the use of wrong spectacles. The amount and conditions of near work should be specifically stated.

Defects of the lacrymal apparatus must be treated (Chap. XXXII.). If no defect is noted, local treatment of the hyperæmia is ordered for a time: but if the condition does not improve, the patency of the lacrymal passages must be demonstrated by syringing. The beginner must be careful, however, that he does not do harm rather than good.

Errors of metabolism must be treated on general medical principles. Such causes are easily overlooked: hence they should be specially borne in mind.

Local treatment consists in bathing the eyes frequently with warm boric lotion, with or without a mild astringent, *e.g.*, zinc sulphate, gr. i. or ii. to ℥ i. A drop of a mixture of equal parts of tincture of opium and distilled water, night and morning, will be found soothing. Hazeline, 20 ♏ to ℥ i., is sometimes useful, but varies in its effect in different people. Cocain must be used with diffidence: its effects are transitory, and it has a deleterious action upon the corneal epithelium, but in quite weak doses often affords much comfort.

In cases where temporary alleviation—usually of the disfiguring signs—is insistently desired, a drop of adrenalin

solution (1 in 1000) instilled into the eye will remove the discomfort and reduce the redness of the conjunctiva. The effect is, however, very transitory, but it will often earn gratitude. It is especially useful after the removal of a foreign body from the cornea.

The *nature of the secretion* in conjunctivitis is of diagnostic importance. It may be watery, mucous, muco-purulent, or purulent, and the disease is often classified accordingly. Most forms of acute conjunctivitis are due to bacterial agency. Unfortunately, each pathogenic organism does not produce a specific clinical picture. It is therefore wise in the meantime to retain the old clinical terminology.

Watery secretion is usually due to reflex secretion of tears. The other types of secretion show some relation to the bacterial cause, and must be distinguished on account of the information they convey as to the probable severity of the condition and the indication they provide for special measures of treatment.

The chief forms of conjunctivitis may be divided into two groups: acute, and sub-acute or chronic. Acute conjunctivitis may be classified as simple acute (including muco-purulent), purulent, membranous, and phlyctenular. Sub-acute or chronic conjunctivitis includes simple chronic conjunctivitis, angular conjunctivitis, follicular conjunctivitis, trachoma, tubercle, and syphilis.

Simple Acute Conjunctivitis.—(*Syn. Catarrhal Conjunctivitis*).—The condition described as hyperæmia of the conjunctiva passes imperceptibly into a condition characterised by greater and more general hyperæmia and a thicker mucous discharge which gums the lids together. The lids are usually described as being stuck together in the mornings, because the condition is most noticed after they have been closed for a considerable period. The causes, symptoms, and treatment are the same as in simple hyperæmia.

Various more intense forms of simple acute conjunctivitis are met with: they are probably all of bacterial origin, the organisms differing in different cases. Among the lower classes the disease is called "blight" and it is commonly attributed to a "cold in the eye." Cold probably acts only by lowering the resistance of the tissues to the action of organisms.

The commonest form is **Muco-purulent Conjunctivitis**. Here, as the name implies, the secretion is muco-purulent: it is more profuse than in the simpler forms. As in most cases

of conjunctivitis the disease is contagious, being transmitted directly by the discharge and possibly by the air of ill-ventilated rooms, though this method is doubtful, since most of the organisms are non-sporing and are easily destroyed by drying. The whole conjunctiva is a fiery red ("pink eye"); all the conjunctival vessels are congested, except the circumcorneal zone in the milder cases (*vide* p. 78). Flakes of muco-pus are seen in the fornices, and often between and upon the margins of the lids. If the discharge is allowed to dry the lashes become matted together by dirty yellow crusts. These may be easily mistaken for the condition found in blepharitis, but if the crusts are bathed off the underlying lid margins will be found healthy. Flakes of mucus passing across the cornea often give rise to coloured haloes, owing to their prismatic action. These "haloes" must be carefully distinguished from those met with in glaucoma (*q.v.*).



Fig. 95.—Koch-Weeks' bacilli (× 1000).

The discharge is at first mucous, but gradually becomes more purulent. Beginners are liable to mistake muco-purulent for true purulent conjunctivitis. In the former, the more purulent masses are found among the lashes and at the inner canthus, whilst the fornices and bulbar conjunctiva show only flakes of translucent or yellowish secretion. In the latter, crusts of inspissated pus may be seen among the lashes and at the canthi, but when the lids are separated fluid pus wells out.

The disease reaches its height in three or four days: if untreated it is liable to pass into a less intense, chronic condition. Complications are rare, but abrasions of the cornea are liable to become infected and to give rise to ulcers. Occasionally marginal ulcers form—in debilitated or old people, or as the result of improper treatment.

Muco-purulent conjunctivitis frequently complicates phlyctenular conjunctivitis.

Pathology.—Muco-purulent conjunctivitis is frequently

caused by the Koch-Weeks' bacillus (Fig. 95). This is a very slender rod, varying much in length. It stains badly with the ordinary basic dyes, *e.g.*, Löffler's methylene blue; it is decolourised by Gram. Groups of bacilli found in much degenerated "skeletonised" pus cells are very characteristic.

The organism is rapidly destroyed by drying. It has been known to give rise to very definite epidemics. In England the cases are usually sporadic, though moderate transmission is common. An attack confers immunity for some time.

The Koch-Weeks' bacillus is by no means the only cause of muco-purulent conjunctivitis. Diplococci which are indistinguishable from pneumococci (Fig. 96) also cause it, probably more frequently in England. Pneumococcic conjunctivitis, though not definitely separable from the other acute forms clinically, shows distinct tendencies which should be borne in mind, the more so since the pneumococcus is the cause of hypopyon ulcer (*q.v.*). There is usually more œdema (chemosis), small ecchymoses are common, and a membranous film may form—"pseudo-membranous conjunctivitis." It is commonest in northern countries and in the cold weather, and is more often found in children than adults. It ends with a crisis, like pneumococcic infection of the lungs, after which the organism rapidly disappears from the secretion. It is often accompanied by nasal catarrh, which may precede or follow the inflammation. Iritis is very rare as a sequel of conjunctivitis, but pneumococcic conjunctivitis is exceptional in this respect. The inflammation of the iris is set up by absorption of toxins (*cf.* Hypopyon Ulcer of the Cornea).



FIG. 96.—Pneumococci (× 1000).

The influenza bacillus, which is distinguished with difficulty from the Koch-Weeks' bacillus, is responsible for conjunctivitis during influenza epidemics, more often in children than adults.

The muco-purulent conjunctivitis associated with phlyctenular conjunctivitis is usually due to staphylococcus aureus.

which may also set up conjunctivitis in cases of blepharitis and eczema or impetigo of the skin. The organism sometimes causes a muco-purulent discharge after cataract extraction and other operations, and also in the new-born, probably owing to the defective resistance of the tissues in these conditions. In the presence of irritating dust a staphylococcic conjunctivitis of slight intensity may be set up. The presence of staphylococci is liable to aggravate conjunctivitis set up by some other organism.

Other organisms have been found, but their ætiological relationship to the disease is not proved.

Muco-purulent conjunctivitis generally accompanies or follows measles and frequently scarlet fever. It also occurs with acne rosacea, when minute nodules, somewhat like phlyctens, form at the limbus and on the cornea. It frequently recurs in these cases, and is intractable. Muco-purulent conjunctivitis sometimes occurs among people who use swimming baths. It often spreads rapidly through schools and other such institutions.

Fig. 97. -Undine.

Treatment.—The best treatment for muco-purulent conjunctivitis is the frequent washing out of the conjunctival sac with lotio hydrargyri perchloridis (1 to 10,000) (called hereafter sublimate lotion). It should be warmed and diluted by the addition of a little boiling water. The lotion must not be simply applied to the lids. It is well for the surgeon himself or a competent nurse to wash out the conjunctival sac the first time. For this purpose an "undine" (Fig. 97) is the most convenient reservoir. The lids are everted and the lotion is poured from a little height over the whole surface, every crevice being irrigated as thoroughly as possible. The nozzle of the undine should not be allowed to touch any part of the eye. The patient may be directed to use the ordinary eye-bath for the application of eye lotions. If a child, the parents should be instructed to hold the lids apart, the child lying upon its back. A pad of cotton wool, dripping with the lotion, which may be warmed, is then held over the eye and the lotion squeezed out: the process is repeated until all discharge has been washed away. The lotion may be warmed by placing the bottle in hot water, but the addition of an equal part of hot

water suffices ; eye lotions act chiefly by washing out deleterious material, since they cannot be used sufficiently strong to act as efficient antiseptics. For this reason boric acid lotion is probably nearly as efficacious and is less irritating.

Boric acid ointment or sterile vaseline is smeared along the lids at bed-time, or, in children, as often as they are put to sleep : it prevents the lids from sticking together—a two-fold benefit, that of preventing discharge from being retained, and that of obviating pain on opening them.

The eyes should never be bandaged, as it prevents the free exit of the secretion. If there is any photophobia a shade or dark goggles should be worn. The patient should spend as much time out of doors as possible.

If this treatment is properly carried out, the patient will be well in a few days. Even if only partially successful there will then be less discharge.

If the case is not progressing as rapidly as could be desired, or the attendants are not reliable, and if the discharge is subsiding, it is wise to paint the lids once with silver nitrate solution (gr. x. to ℥ i.). This is the strength which should always be used for painting lids. Stronger solutions act too vigorously as caustics, and if a caustic effect is desired, it can be obtained with greater precision by other means. Weaker solutions are precipitated by the chlorides in the lacrymal secretion, so that they are practically useless.

Fig. 98.—Glass rod.

The following is the best method of painting lids. A glass rod is used, tapering at each end (Fig. 98). The finger should be passed over the ends each time before use to make sure that they are not chipped. The end is dipped in lotion to damp it. A *very thin* wisp of cotton wool is then tightly wound round the end, starting where it begins to taper : this fixes the wool. The end of the wisp is left loose, so that it may absorb the solution. The other end of the rod is armed in the same manner.*

The patient, if a child, is placed upon its back. The lids are everted ; the wool, dipped in the silver solution, is applied freely to the conjunctival sac, the cornea being protected as much as possible. It is quite unnecessary to neutralise the excess of nitrate with salt solution, as is often taught. The excess may be mopped up with a pad of dry absorbent wool. If, as is usually the case, the other eye is affected, the other

end of the rod is used in the same manner. In the absence of a glass rod the best implement is an ordinary wooden match, used in the same way. The match can be thrown away after use. The glass rod must be sterilised by boiling. A camel's hair brush should not be used : it cannot be kept aseptic.

A single painting with silver nitrate will often produce an excellent result. It is good as a prophylactic if discharge is inadvertently introduced into a normal eye. Other preparations of silver—protargol (20 per cent.), argyrol (25 per cent.), &c. are not so efficacious, but they have the advantage of being less painful.

Silver nitrate acts by forming a thin epithelial eschar and coagulating the muco-purulent discharge. The bluish-white film is cast off in flakes, and until this process is complete the feeling of a foreign body in the eye is experienced. The irritation is reduced if the lid is kept everted for a few minutes and the flakes gently removed with cotton-wool. Silver nitrate is not strongly bactericidal, but the organisms are entangled in the coagulum and removed with it. Moreover there is a powerful physiological response to the caustic ; hyperæmia increases and the tissues are flooded with blood serum, which can thus more effectively exert its bactericidal and antitoxic powers. The slighter efficacy of protargol and the modern colloid substitutes for silver nitrate is to be attributed to their smaller caustic and irritant properties, which are often regarded as their chief advantages.

The conjunctiva generally returns to a perfectly normal condition. If the case has been neglected and chronic inflammatory signs persist, astringents should be used as for chronic conjunctivitis (*q.v.*).

Since the disease is contagious care must be taken to prevent its spread. The patient must keep his hands clean and no one else must be allowed to use his towel, handkerchief, &c.

Purulent Conjunctivitis (*Syn. — Acute Blennorrhœa, Gonorrhœal Conjunctivitis*) is a much more serious condition. It occurs in two forms —as ophthalmia neonatorum in babies, and as gonorrhœal conjunctivitis in the adult. Certainly the former, and probably the latter, is not invariably gonorrhœal.

Gonorrhœal Conjunctivitis (*Syn.—Acute Blennorrhœa of Adults, &c.*).—Gonorrhœal conjunctivitis is even more serious in the adult than in babies ; fortunately, considering the prevalence of gonorrhœa, it is comparatively rare (1 in 700—800 cases). While generally due to the gonococcus it is important

from the medico-legal point of view to remember that the same features may be found with streptococci, diphtheria, and probably with mixed infections. The gonococcus is a bun-shaped diplococcus, staining readily, decolourised by Gram, and found within both leucocytes and epithelial cells (Fig. 99). The micrococcus catarrhalis and the meningococcus, both Gram-negative, are sometimes found in the conjunctival sac. They may be distinguished from the gonococcus by the ease with which cultures are obtained and by agglutination tests. The micrococcus catarrhalis is rarely found in acute conjunctivitis, but more often in chronic and post-operative forms.

The disease is due to direct infection from the genitals. Males suffer most, the right eye before the left in a right-handed person. There is more swelling of the lids and conjunctiva than in children, copious purulent discharge, more tendency to involvement of the cornea, and marked constitutional disturbance—rise of temperature, and so on, but especially very marked depression of spirits. The greater danger to the cornea is due to the chemosis, which produces blood and lymph stasis and facilitates the retention of secretion.



Fig. 99.—Gonococci (× 1000).

The incubation period is a few hours to three days. The upper lid becomes enormously swollen and tense, overhanging the lower, and edged with pus. Eversion, which is difficult, shows that the palpebral conjunctiva is deep red and velvety: rarely there is a membrane. Occasionally the discharge is sanious rather than purulent, especially in streptococcic cases. There is great pain; the preauricular gland is enlarged and tender, and may suppurate.

After two or three weeks the purulent discharge diminishes, but subacute conjunctivitis with much papillary thickening of the conjunctiva persists for several weeks longer. The gonococcus is still present—a point of great importance, both as regards contagion and treatment. No immunity is conferred by the attack.

The most important point in diagnosis is the coincidence of urethritis. The most important point in prognosis is the condition of the other eye.

Corneal complications are the rule, and constitute the causes of blindness. There may be diffuse haziness of the whole cornea, with grey or yellow spots near the centre. Ulcers may occur at any part, and are due to necrosis of the epithelium through direct invasion by the organisms. Marginal ulceration, which may extend completely round the cornea, is due to retention of pus in the angle formed by the chemotic conjunctiva. When ulceration has commenced it progresses rapidly and deeply, since the tissues are bereft of their first line of defence—the epithelium. Perforation is therefore common, with all its attendant dangers (*vide* p. 192). Ulceration commencing late in the history is not so dangerous.



Fig. 100.—Buller's shield.

The greatest care should be taken to prevent injury to the cornea during the manipulation necessary for diagnosis and treatment. Abrasions are easily produced by the finger nails, and even by the rough use of wool swabs. Such abrasions rapidly become dangerous ulcers.

Iritis and iridocyclitis, with attendant complications, may arise independently of perforation of the cornea, and lead to serious diminution of vision.

Gonorrhœal arthritis is not uncommon, and endocarditis and septicæmia may arise in complications.

Treatment must be directed first to protection of the other eye. This is at once sealed up with a Buller's shield, which consists of a watch-glass stuck in a frame of adhesive plaster, or better, rubber (Fig. 100). The rubber is hermetically sealed down to the face and nose except at the lower outer angle, where a small piece of tubing is inserted under the edge. If this means of ventilation is not adopted the glass becomes hazy and the eye cannot be properly observed; moreover, the irritation to the eye is greater. Special attention should be directed to fixing the shield near the nose, *i.e.*, on the side of the source of

infection. Patients should be told to sleep lying on the same side as the affected eye.

If the second eye should show signs of infection it must be treated, but every utensil or dressing which is applied to the eye must belong to a totally different set from that used for the worse eye, otherwise it may be inoculated rather than treated. The less affected eye must always be dressed first.

If pus from a gonorrhœal conjunctivitis spurts into the surgeon's eye, the conjunctival sac should be freely irrigated with sublimate lotion and the lids painted once with silver nitrate (*vide* p. 148). The eye should be carefully watched, but no further drastic treatment applied unless conjunctivitis supervenes. The accident is due to carelessness, for every attendant on a gonorrhœal case should wear protective goggles.

If the disease is fully established and there is abundant purulent discharge, the eye must be irrigated every two hours during the day and every four hours during the night with warm saline (0·85 per cent.), boric, or weak sublimate lotion (1 in 8000). Iced wet dressings are applied in the intervals, but are to be bandaged on quite loosely, so as to avoid retention of discharge. The patient is kept in bed, and if intelligent, can bathe his lids in iced lotion in the intervals. Iced applications afford much relief to the patient, though hot ones are probably more efficacious.

Most reliance must be placed on applications of silver nitrate, but they must be used with discretion. They are specially indicated in gonorrhœal conjunctivitis because the gonococcus is an intracellular organism. As already stated (p. 148), silver nitrate destroys the superficial layer of epithelial cells, which is cast off as filmy eschar, carrying the entangled organisms with it. When not contra-indicated the conjunctiva of the everted lids should be well painted with silver nitrate, gr. x. to ℥ i., not oftener than once a day. Special care must be taken to avoid injury (*vide* p. 83). This treatment is contra-indicated in the very early stage before free discharge has set in, and also in later stages if there is much brawny swelling with comparatively little discharge. Under these conditions the stasis is so great that reaction is inefficient; the caustic, in fact, induces the necrosis which it should be our endeavour to avoid. In such cases hot applications and leeches should be relied upon. The latter are applied over the temporal region near the outer canthus. If the lids are very tight the outer canthus should be split (canthoplasty). The ends of strong

blunt-pointed scissors are inserted between the lids into the angle under the outer canthus, which is then divided by a single snip in a horizontal direction. This has the good effect of bleeding the patient slightly, and also of giving free exit to discharge.

Scarification of the chemosed conjunctiva has been recommended and may be done. The removal of the ring of conjunctiva which overhangs the cornea is good, but demands cocain, which has a bad effect on the corneal epithelium. Mercury, by inunction or small doses of calomel, is said to diminish the chemosis.

Römer advocates passive immunisation by means of the local application of a specific gonococcic serum. It may be dusted into the conjunctival sac or used as a 1 per cent. solution in 0·85 per cent. sodium chloride for irrigation. Since no general immunity is conferred by the organism active immunisation with a gonococcic vaccine is useless.

It is of great importance to attend to the general health. The bowels must be kept freely open. The strength must be reinforced by every available means –good foods, tonics, alcohol if necessary. An occasional sleeping draught and the use of sedatives must be ordered according to general principles. Every effort must be made to combat the depression from which these patients suffer.

In the final stage of the disease silver nitrate should be used at increasing intervals for a week or fortnight after the purulent discharge has ceased. Astringent lotions are then employed (*vide* p. 165).

Atropin should be used in all cases where the cornea is involved, since this is always accompanied by some iritis; but the intraocular tension should be watched.

Corneal complications require very active treatment (*vide* pp. 194, *sqq.*).

Metastatic Gonorrhœal Conjunctivitis sometimes occurs in adults, associated with gonorrhœal arthritis. It is a mild simple conjunctivitis, generally bilateral, and occasionally accompanied by iritis. It is probably due to endogenous infection from gonococci in the blood. It usually responds readily to local treatment, but is apt to recur if the arthritis relapses.

Ophthalmia neonatorum is a preventable disease occurring in new-born children as the result of carelessness at the time of birth; it is responsible for 50 per cent. of blind children and about 7—8 per cent. of all blind people (8 per cent. in

U.S.A., de Schweinitz). It is due to infection by vaginal or fæcal matter, or from dirty rags used to clean the eyes. Purulent discharge is usually noticed on the third day; when it occurs later it is generally due to post-partum infection. In rare cases the disease is already present at birth.

Any discharge, even a watery secretion, from a baby's eyes during the first week should be viewed with suspicion, since tears are not secreted at this early date. In cases of infection the discharge rapidly becomes muco-purulent and then purulent. Both eyes are nearly always affected, though one is usually worse than the other. The conjunctiva becomes intensely inflamed, bright red, and swollen, and pours out thick yellow pus. Marked chemosis is a distinguishing feature from severe muco-purulent conjunctivitis, and when the lids are separated by retractors the cornea is seen at the bottom of a crater-like pit. There is dense infiltration of the bulbar conjunctiva, and the lids are swollen and tense. Later the lids become softer and more easily everted, the conjunctiva becomes puckered and velvety, and the blood stasis gives place to intense congestion, with free discharge of pus, serum, and often blood. In some cases a false membrane forms, so that the case resembles a membranous conjunctivitis.

There is great risk of corneal ulceration in ophthalmia neonatorum, especially, as is usually the case, when it is due to the gonococcus, which has the power of invading intact epithelium. The slightest haziness of the cornea should be viewed with apprehension. Often the cornea is already ulcerated, and not infrequently perforated, when the child comes under observation. Ulceration usually occurs over an oval area just below the centre of the cornea, corresponding with the position of the lid margins when the eyes are closed, and, consequently, rotated somewhat upwards. More rarely oval marginal ulcers are formed as in the gonorrhœal conjunctivitis of adults, or the ulceration may be central. The ulcers extend rapidly, both superficially and in depth, largely owing to lymph stasis due to strangulation of the nutrient vessels.

Perforation is usually signalised by a black spot or area in the ulcer, caused by protrusion of the iris. Sometimes perforation is sudden, a large part of the iris prolapses, and the lens may be extruded. In some cases there is a black hole in the cornea, filled with clear vitreous.

Metastatic stomatitis and arthritis occur rarely, as in gonorrhœal urethritis. The arthritic manifestations usually appear in the third or fourth week and affect the knee, wrist, ankle,

or, sometimes, elbow. The course is benign, abscesses being rare.

The baby's eyes must be examined as described in method (4), p. 76. The surgeon must wear protective goggles lest pus spurt into his eyes when the child's lids are separated. Retractors must always be used for separating the lids, since the slightest pressure on the eyeball may result in perforation (*vide* p. 84).

A bacteriological examination should be made in every case.

Pathology.—Probably 60—70 per cent. of cases are due to the gonococcus. The Bac. coli is responsible for some of the remainder, probably through fæcal infection, pneumococci, streptococci, and mixed infections for others. Streptococcic conjunctivitis is even more virulent than gonococcic, the cornea being in imminent danger. A film should be taken from every case.

Sequelæ of Ophthalmia Neonatorum.—If the corneal ulceration heals without perforation there is always much scarring of the cornea, but the nebula clears more in babies than in older people. Perforation may be followed by anterior synechiæ (*q.v.*), adherent leucoma (*q.v.*), partial or total anterior staphyloma (*q.v.*), anterior capsular cataract (*q.v.*), panophthalmitis, &c. Ophthalmia neonatorum is the commonest cause of anterior staphyloma.

When vision is not completely destroyed, but is very seriously impaired by the corneal opacities resulting from ophthalmia neonatorum, the development of macular fixation, which takes place during the first six weeks of life, is interfered with, resulting in the development of nystagmus (*q.v.*), which persists throughout life. Nystagmus may not become manifest until some considerable period after the ophthalmia.

Treatment.—The disease is preventable; *prophylactic* treatment is therefore of prime importance. The new-born baby's closed lids should be thoroughly cleansed and dried. The lids are then separated, and a drop of silver nitrate solution, 1 per cent., instilled into each eye (Credé's method). The eyes must be carefully watched during the first week.

If the disease is established the eyes must be irrigated every two hours with saline or weak antiseptic lotion, and the lids painted once daily with 2 per cent. silver nitrate (*vide* pp. 147, 151). The surgeon or nurse must wear protective goggles. In irrigating or painting the greatest care must be exercised to avoid touching the cornea, thus injuring the already softened epithelium. Argyrol, protargol, and other colloid

preparations of silver must not be used instead of silver nitrate, but may be used as drops in the later stages when the critical period has passed. The slightest sign of corneal haziness is an indication for the use of atropin, 0·5 per cent. (*vide* p. 194). The use of antigonococcic serum has been advocated.

The Prevention of Ophthalmia Neonatorum and Blindness due to this Cause.—Though ophthalmia neonatorum is a preventable disease, and much attention has been directed to its prevention, it still remains the cause of a large amount of blindness. It was made notifiable throughout England and Wales in 1914, being defined as a "purulent discharge from the eyes of an infant commencing within twenty-one days of birth." Under the rules of the Central Midwives Board every midwife is required, under severe penalties, to call to her assistance a medical man in any case of "inflammation of or discharge from the eyes (of the child), however slight," and to send notice to the supervising authority that medical help has been sought. The administrative measures ensure that midwives' cases come promptly under the supervision of medical officers who are accustomed to deal with this class of case. There is evidence that some medical practitioners fail to recognise early and notify promptly cases of ophthalmia neonatorum, with the result that efficient treatment is not begun early enough to prevent serious impairment of vision or even blindness.

Since most of the worst cases of ophthalmia neonatorum are gonorrhœal, the combating of this disease should diminish the incidence of blindness. The next line of defence is the treatment of the expectant mother, and ante-natal measures should be adopted, whenever possible, to cure disease causing abnormal vaginal discharge in the mother. Next, the greatest care should be exercised to avoid vaginal discharge reaching the eyes of the baby at birth. "As soon as the child's head is born, and if possible before the eyes are opened, its eyelids must be carefully cleansed. They should be thoroughly wiped with clean material such as cotton-wool, lint, or rag, using separate pieces for each eye. . . . When the baby is bathed the discharges with which its body is covered during labour are washed off into the bath-water. If its face is washed in this water matter may get into the eyes." (*Instructions to Midwives, by Central Midwives Board.*)

Prophylactic Drugs.—In the hands of medical men Credé's method has been most successful and is to be recommended, especially in cases where abnormal vaginal discharge is known

to exist. It is not, however, to be recommended for universal use by midwives for the following reasons: (1) The midwife may be apt to think that, having dropped a little solution into the eye, she has done all that is necessary, and consequently neglect the scrupulous cleansing which is even more important; (2) the use of drugs will induce an inflammatory reaction which, on the one hand, she may mistake for the onset of the disease and notify accordingly; or (3) on the other hand, she may regard as "only a little reactionary discharge" what is really a manifestation of the disease itself.

When ophthalmia neonatorum has actually developed, a medical practitioner who has not had very thorough training and experience in the treatment of the diseases should at once obtain the assistance of an ophthalmic surgeon. It is to be remembered that any discharge from a baby's eyes during the first week of life is pathological, and no risks should be taken. Given satisfactory home conditions, competent medical supervision, and skilled nursing, home treatment gives the best results. In many cases these requirements cannot be fulfilled. In these cases probably the ideal method is to transfer both mother and child to a hospital where special provision is made for such cases. This is possible in London at St. Margaret's Hospital, under the Metropolitan Asylums Board, and also in most other large towns, and in many county boroughs; but it is eminently desirable that similar provision should be made in suitably distributed General or Ophthalmic Hospitals, since special hospitals for the purpose are liable to become stigmatised as venereal hospitals. Failing this method of dealing with the case, the child should be taken daily to the out-patient department of a General or Ophthalmic Hospital, and the eyes irrigated as often as possible by a competent nurse at home. This method is fairly satisfactory in all but the worst cases, but is not always practicable in outlying districts.

Membranous Conjunctivitis (*Syn.—Diphtheritic Conjunctivitis*). As in inflammation of the throat the surface may become covered by a fibrinous membrane, so the same may occur in the conjunctiva; and just as the milder clinical varieties in the former were distinguished as croupous from the severer or diphtheritic, so also with conjunctivitis. It has been placed beyond dispute, however, that mild cases may be diphtheritic, and severe non-diphtheritic; hence it is best to speak simply of membranous conjunctivitis until a bacteriological examination has placed the matter beyond

dispute. A variety of organisms other than the diphtheria bacillus, *e.g.*, pneumococcus, streptococcus, can produce a membrane, especially in weakly children, *e.g.*, after measles and scarlet fever, and in association with impetigo; these cases are sometimes called pseudo-membranous. They cannot be distinguished clinically with certainty.

Membranous conjunctivitis occurs chiefly in children, and shows all degrees of severity; it may be as virulent as the worst cases of gonorrhœal ophthalmia. It is uncommon in England, but it is of the utmost importance that it should be recognised when seen, not only on account of the grave danger to the eye, but also from the risks of contagion.

In mild cases there is some swelling of the lids and a muco-purulent or sanious discharge. On everting the lids the palpebral conjunctiva is seen to be covered with a white membrane, which peels off readily without much bleeding.

In severe cases the lids are more brawny: the conjunctiva is permeated with semi-solid exudates, which impair the mobility, compress the vessels, prevent the formation of a free discharge, and tend to necrosis both of the conjunctiva and cornea. Here the membrane separates much less readily, the underlying surface bleeding unless it is too infiltrated and solid. The membrane may be patchy or cover the whole palpebral conjunctiva, often beginning at the edge of the lid. It is seldom found on the ocular conjunctiva. The pre-auricular gland may be enlarged and may suppurate. The temperature is raised, unless the patient is in a moribund condition. Albumin is frequently present in the urine.

For six to ten days there is great peril to the cornea. Then the sloughs begin to separate and the discharge becomes more profuse. In a few days the conjunctiva assumes a red and succulent appearance. There is danger now of adhesions forming between the palpebral and bulbar parts of the conjunctiva (symblepharon).

Post-diphtheritic paralysis, even of accommodation, is rare.

Cases of less severe but more chronic membranous conjunctivitis are occasionally met with. In them the membrane is cast off, but occurs again and again. The pathology of these cases is not understood; it has been seen as a complication of erythema multiforme.

Pathology.—Competent bacteriologists have shown that there is little or no relationship between the severity of the local condition and the presence or absence of the Klebs-Löffler bacillus (Fig. 101). Only series of cases in which positive

results have followed inoculation into animals are absolutely trustworthy, owing to the difficulty of distinguishing the diphtheria from the xerosis bacillus, with which it is morphologically identical. The pseudodiphtheria bacillus forms little or no acid in culture and is not virulent for guinea-pigs, and the different varieties may be distinguished by agglutination tests. Neisser's stain (acetic methylene blue and Bismarck brown) demonstrates blue granules at the poles of true diphtheria bacilli in cultures of nine to twenty-four hours. Inoculation tests only are absolutely reliable.

Other cases may be due to the action of heat, caustics, severe atropin irritation, herpes iris, and other non-bacterial causes. Other bacteria which occasionally form membranes are pneumococcus, streptococcus, Koch - Weeks' bacillus, gonococcus, staphylococcus, Friedländer's pneumonia bacillus, bacterium coli, &c. Streptococcic conjunctivitis, a very virulent form, occurs chiefly in children associated with measles, scarlet fever, whooping cough, and influenza.

Fig. 101. Diphtheria bacilli (× 1000).

It is quite rare to obtain evidence of primary diphtheria of the throat, though the disease may have been derived from a case of faucial diphtheria, and extension to the nose and throat by way of the lacrymal sac and nasal duct occurs. The genitalia should be examined for diphtheria or leucorrhœic discharge.

Treatment. -Every case should be treated as diphtherial unless good negative evidence is afforded by films and serum cultures. In mild cases isolation need not be strict until the bacteriological report is obtained on the second day.

The treatment is essentially that of purulent ophthalmia, with one important exception, viz., that painting with nitrate of silver is not to be resorted to. Further, canthoplasty is not to be done, since both these procedures increase the area from which toxins may be absorbed.

Local treatment consists of hot irrigations and bathings as

FIG. 1.

FIG. 2.

PLATE IV. (*To face p.* 159.)

Fig. 1.—Phlyctenular conjunctivitis, showing two phlyctens on the conjunctiva and one on the cornea.

Fig. 2.—Follicular conjunctivitis.

for purulent ophthalmia (*q.v.*). One drop of atropin should be instilled at the commencement of treatment. Quinine lotion, gr. iii. to ℥ i., with a minimum of acid to dissolve the salt, has been recommended.

The most important general treatment is the administration of antitoxin as in faucial diphtheria. Since the antitoxin is innocuous it should be used at once in every doubtful case. I have obtained benefit by local instillations of antitoxin, which is a rational procedure but seems to have been neglected. Special attention should be paid to the nutrition, and tonics are indicated.

Corneal complications must be suitably treated (*vide* p. 194). Antitoxins are specially useful, as may be shown by experiment. If diphtheria toxin is injected into the cornea of two rabbits and one is given an intravenous injection of antitoxin the cornea of this animal will remain clear, whereas that of the other will become cloudy. Corneal ulceration, however, is usually due to secondary infection with pyogenic organisms. It may start at the middle or margin of the cornea and is not wholly due to interference with nutrition.

In streptococcic membranous conjunctivitis the danger of necrosis of the cornea and even of the death of the patient is such that immediate recourse should be had to a specific antistreptococcic serum.

Phlyctenular Conjunctivitis (*Syn.*—*Eczematous Conjunctivitis*).—In phlyctenular conjunctivitis (Plate IV., Fig. 1) one or more small, round, grey or yellow nodules, slightly raised above the surface, are seen on the bulbar conjunctiva, generally at or near the limbus; they rarely occur on the palpebral conjunctiva. The disease is very frequently complicated with muco-purulent conjunctivitis, in which case the whole conjunctiva is intensely reddened. In pure phlyctenular conjunctivitis the congestion of the vessels is limited to the area around the phlyctens.

The disease is most frequent in children from five or six to ten or twelve years of age, but not very young children: it is rarely seen in adults. The children often have enlarged lymphatic glands in the neck, &c., or other signs of tubercle; on the other hand, every sign of tubercle may not infrequently be lacking: the children, however, are seldom robust. The first attack often follows an exanthem, especially measles. Rhinitis and adenoids are frequently present; signs of congenital syphilis may be found.

Phlyctens, as the name suggests (φλύκταινα, a bleb), at

first much resemble blebs: it is doubtful, however, whether there is a true vesicular stage. They may be so small as to be seen with difficulty, but they usually measure about 1 mm. in diameter, occasionally reaching a diameter of 3 mm. or 4 mm. The larger ones are yellow, and have been described as pustules. In the later stages the epithelium over the surface is often destroyed, small ulcers being formed. When this occurs on the conjunctiva proper it is of little moment, since healing takes place rapidly without the formation of a scar. When it occurs on the cornea, as is very frequently the case, it is much more serious (*vide* p. 210).

Very frequently the skin of the lids and cheeks shows an eczematous condition, and eczema will be found not uncommon, if searched for, in other parts of the body, especially in the scalp. This fact has led the condition to be regarded by some as an ocular manifestation of eczema. The disease has indeed been regarded as an exanthem. It is probable that in most cases the eczema of the lids and face is secondary to the continual irritation of the skin induced by the overflow of tears and the rubbing of the wet surface with the hands.

Pathology throws some light on the disorder, though it by no means settles the causation. A simple phlycten shows in section a triangular area of intense infiltration, the apex of the triangle being towards the deeper layers. The subepithelial adenoid layer normally contains a few mononuclear lymphocytes, but in the phlycten they are very numerous and closely packed together. The epithelium is intact, and it is doubtful if a vesicular stage has ever been observed.

If there is a considerable amount of conjunctivitis of the muco-purulent type, not only are lymphocytes present but there are also many polymorphonuclear leucocytes, both in the sub-epithelial tissues and amongst the epithelial cells. In such cases the epithelium is quickly desquamated.

If a bacteriological examination is made, many of the organisms of muco-purulent conjunctivitis may be found. In the pure phlyctenular cases only staphylococci are found in abundance. Now staphylococci are not so common as might be expected in the normal conjunctival sac; it has, therefore, been concluded that the disease is due to staphylococcic infection. If staphylococci are rubbed into a healthy or excoriated conjunctiva a transitory redness occurs and rapidly passes off. Phlyctens have never been produced in this manner. The nearest approach to the artificial production of phlyctens has been by injecting cultures of tubercle

bacilli, in which the organisms have been killed, into the veins of rabbits. It is doubtful if the infiltrates were real phlyctens in these experiments, but it is certain that the administration of tuberculin has not infrequently been followed by an attack of phlyctenular conjunctivitis in the human subject. On the other hand, the superficial position of the lesions and the analogy to other ulcerative conditions of the cornea favour the view that the disease is ectogenous in origin, the morbid agent being at present unknown.

We are now in a position to judge more accurately what probably occurs in phlyctenular ophthalmia. It is not improbable that the true cause of the outbreak is a toxæmia, the toxins—probably tuberculous—not being very potent, but sufficient if the patient is already in a weakened condition. The irritation of the eye leads the child to rub it vigorously. The lacrymation and rubbing cause an eczematous condition of the skin, in which the staphylococci normally present flourish and increase. These are rubbed into the eye, increasing the irritation, without being primarily responsible for the disease. Other organisms are also rubbed in if they happen to be present; they find a suitable nidus in the debilitated conjunctiva, and an acute muco-purulent conjunctivitis is superposed upon the phlyctenular disease.

Simple phlyctenular conjunctivitis is attended with few symptoms. There is some discomfort and irritation associated with reflex lacrymation. If there is no muco-purulent complication and if the cornea is not involved there is little or no photophobia.

Complications, however, are the rule, partly because the behaviour of the child conduces to them, partly because the favourite situation for the phlyctens is near the cornea. Here they are often astride the limbus. It has already been mentioned that the epithelium of the cornea is closely associated anatomically and developmentally with the conjunctiva. It is not surprising therefore that there is a great tendency for the superficial layers of the cornea to suffer when the conjunctiva is disordered, and this is seen par excellence in phlyctenular ophthalmia. The special corneal complications will be considered later (*vide* p. 210). In all such cases lacrymation is increased, muco-purulent discharge is often present, and photophobia is intense.

The term photophobia (φῶς, light; φόβος, fear; dread of light) is a misnomer. It is the term applied to the blepharospasm which is set up by the conjunctival, or more probably

corneal, irritation, and which becomes greatly increased on the slightest attempt to separate the lids, especially if the attempt is made in bright light. This blepharospasm is not abolished in the dark ; it is abolished by thorough application of cocain, though this is difficult to effect. It must be concluded therefore that it is a reflex due to afferent impulses travelling along the fifth nerve, not along the optic nerve. It has been said that light acts as the stimulus to the fifth nerve endings in the cornea. This conjecture must be regarded as a survival of the old fallacy. There is no evidence to prove—though it is not disproved—that light can stimulate the fifth nerve endings. It is far more probable that " photophobia " is due to a vicious circle of such a nature that movement of the lid over a spot denuded of epithelium, where the nerve endings are laid bare, causes reflex contraction of the orbicularis ; this increases the irritation, increasing in turn the blepharospasm.

Photophobia is more intense when the phlyctens are near the cornea, than when at a distance. It varies rather with their number than their size, and is extreme if they are so numerous as to form a ring round the cornea.

Temporary blindness has been observed occasionally in children after long-continued blepharospasm. It passes off in two or three weeks and is probably functional, induced primarily by the desire not to see and facilitated by the effect of prolonged pressure upon the globe by the tightly closed lids.

Severe blepharospasm makes the greatest care in the first examination imperative. The condition of the cornea is in all cases and at all costs to be placed beyond doubt. It facilitates investigation if the lids are gently separated and a drop of 2 per cent. cocain instilled. The child is left for 5 to 10 minutes, and the eyes are then examined with all the precautions described above (p. 83).

Phlyctenular conjunctivitis shows a very marked tendency to recur at intervals during the age period which is specially concerned. These recurrences usually take place when some intercurrent malady or defective condition in the patient's surroundings leads to lowering of vitality.

Treatment.—Simple phlyctenular conjunctivitis is usually readily amenable to treatment, which must be local and general.

Local treatment consists in bathing the eyes frequently with hot boric or sublimate lotion : yellow oxide of mercury

ointment, gr. iv.—viii. to ℥ i., is used, a piece the size of a hemp seed being placed within the lids three times a day; the eye is gently massaged by a finger placed upon the upper lid, moving the lid upon the globe.

Ointments are best applied on a glass rod. The child is placed upon its back on a couch and an assistant holds the arms against the body, keeping the legs still by pressure with the elbows. The surgeon separates the lids with two fingers of one hand and places the end of the glass rod carrying the ointment between the separated lids. Keeping the rod in position, the lids are allowed to close upon its end, and it is then withdrawn by carrying it outwards towards the temple. The other end of the rod and the surgeon's other hand are used for the other eye.

If there is any corneal complication, or evidence of its imminence, atropin, gr. iv. to ℥ i., is combined with the yellow ointment.

Very frequently the soddening of the skin with tears and the wrinkling of the skin through blepharospasm cause excoriations (rhagadæ) at the outer canthus. They much increase the blepharospasm and should always be looked for and treated. They are very troublesome unless attacked by cauterisation. They should be touched with the sharp point of the solid silver nitrate or the mitigated silver stick.

An efficient substitute for the yellow oxide which has fallen into undeserved disuse is finely powdered calomel, dusted into the eye, best from a camel's hair brush, which is not allowed to touch the eye. It often produces a remarkable improvement to intractable cases, but it must not be employed if iodides are being given internally; under these circumstances the unstable and extremely irritating mercurous iodide is formed in the conjunctival sac.

The photophobia is best treated, when severe, in the following manner. The child's face should be plunged in cold water and the mouth and nose kept under water until he struggles for breath; this is repeated three or four times—daily if necessary. The treatment is useless if not carried out ruthlessly, but no other method is so efficacious. It is not uncommon for a single application to render further treatment of the photophobia unnecessary.

The eyes are not to be bandaged unless corneal ulceration is so severe as to assume the preponderant *rôle*. A shade, covering both eyes and extending well over the temples, should be ordered. Smoked glasses may be substituted, but

they should not be tightly fitting goggles, which will become soiled with the discharge and are cleaned with difficulty.

General treatment is never to be neglected, otherwise recurrence is inevitable. Fresh air is the best tonic, and the children should be kept out of doors as much as possible : the windows of living and sleeping rooms must be kept open. Cold or sea-baths are useful. Good food, with a plentiful supply of fresh vegetables, is indicated. A calomel purge should initiate the general régime.

Cod liver oil and maltine are given in the cool weather and throughout the year if well tolerated. Phosphates and iodide of iron form substitutes or supplementary tonics.

General régime must be continued for a prolonged period in order to prevent recurrence, and, in any case, phlyctenular ophthalmia should be regarded as a sign of debility which requires attention (*vide* p. 212).

Simple Chronic Conjunctivitis occurs as a continuation of simple acute conjunctivitis, sometimes in spite of orthodox treatment, especially in the " gouty " type of patient. It is frequent when the cause of irritation is continuous—smoke, dust, heat, bad air, late hours, abuse of alcohol, and so on. A very common cause is the chronic reflex irritation induced by errors of refraction, overuse of the eye in bright electric light, &c. Permanent irritation from concretions (*vide* p. 179) in the palpebral conjunctiva, misplaced lashes, dacryocystitis, chronic rhinitis, &c., must be remembered and as far as possible eliminated. Unilateral chronic conjunctivitis should suggest the presence of a foreign body retained in the fornix, or inflammation of the lacrymal sac. It is often necessary to make a thorough and systematic investigation of the local and general conditions before the cause can be found. It is not infrequently associated with chronic intranasal trouble. The disease is too frequently regarded as trivial, but it is a source of great discomfort to the patient, who is duly grateful for permanent relief.

Burning and grittiness are complained of, especially in the evening, when the eyes often become red. Difficulty in keeping the eyes open is a common symptom. The lids may or may not be stuck together on waking.

The discharge is slight, most frequently subnormal, so that the edges of the lids feel hot and dry.

The eyes may look quite normal on first examination. When the lower lid is pulled down the posterior conjunctival vessels are seen to be congested, and the surface of the mucous

membrane is sticky. The palpebral conjunctiva, upper and lower, may be congested, with velvety papilliform roughness. Occasionally it is succulent and fleshy.

Treatment consists in eliminating the cause and restoring the conjunctiva to its normal condition. Errors of refraction and chronic nasal catarrh are perhaps most likely to be forgotten ; they should be sought out as a matter of routine. When heat is a prominent ætiological factor, *e.g.*, in cooks, spectrum blue glasses may be ordered, since they cut off the heat rays to a large extent. The treatment of the special local conditions mentioned above will be discussed in their proper place. A gouty tendency should be treated by an appropriate régime.

Local treatment consists essentially in diminishing congestion and restoring the conjunctiva to its normal suppleness and secretory activity. It must be remembered that the condition is largely one of lack of tone, due to defective response to prolonged irritation. A stimulating treatment is therefore indicated, and is supplied by astringent applications, which not only act by relieving the congestion, but also promote a more healthy lymph flow and glandular secretion.

In mild cases weak astringent lotions suffice, *e.g.*, boric lotion with zinc sulphate, gr. i.—ii. to ℥ i., alum lotion, gr. iv. to ℥ i., &c. They should be used two or three times a day, not immediately before going to bed. Adrenalin has a transient effect in diminishing redness and itching. Boric acid ointment or sterile vaseline should be applied to the margins of the lids at bed-time.

In severer cases a preliminary painting with silver nitrate solution is indicated, repeated once or twice a week if necessary, or the milder protargol (5—10 per cent.) may be used. Silver preparations should not be ordered for application at home, since prolonged use may lead to staining of the conjunctiva (argyrosis).

Atropin, which is always resorted to by the inexperienced in intractable diseases of the eye, does much more harm than good. It causes great inconvenience from paralysis of accommodation, and has little effect upon the conjunctiva, such as it has being deleterious. Apart from this, it is extremely dangerous in elderly patients, who are specially liable to chronic conjunctivitis. In them, more than in others, there is grave danger that atropin may induce an acute attack of glaucoma, a disaster which it is impossible to overrate.

In the more severe cases of chronic conjunctivitis there

is often an abnormal amount of secretion from the Meibomian glands. It should be squeezed out of the glands by pressure on the lid with the thumb against a glass rod laid upon the conjunctival surface.

Angular Conjunctivitis (*Syn.—Diplobacillary Conjunctivitis*) is one of the few forms in which a specific organism causes a typical clinical picture. In it the reddening of the conjunctiva is limited almost exclusively to the intermarginal strip, especially at the inner and outer canthi, and to the bulbar conjunctiva in the same neighbourhood. Besides the conjunctivitis there is also excoriation of the skin at the inner and outer angles, which may be very slight, a mere scurfiness, but is nearly always present. After a few cases have been seen the typical picture is very easy to recognise, but the condition is not always typical. There is discomfort, with slight muco-purulent discharge; blinking is often complained of. Not infrequently there is nasal catarrh, and diplobacilli are found in the nasal secretion. If untreated the condition becomes chronic and may give rise to definite blepharitis. Clear, shallow corneal ulcers may occur, but are rare. They are usually marginal, but may be central and associated with hypopyon (*vide* p. 203). A single attack does not confer immunity, and relapses are not uncommon.



Fig. 102.—Diplobacilli (× 1000).

Pathology.—The disease is due to the Morax-Axenfeld diplobacillus (Fig. 102). The bacilli consist of pairs of large, thick rods, placed end to end. They stain well with basic stains, are decolourised by Gram, and are easily recognised in films. There is an incubation period of four days. The diplobacilli are strongly resistant to drying. They have been found in the nasal tract of healthy persons, and are often present in the nasal discharge in cases of angular conjunctivitis.

Treatment.—Diplobacillary conjunctivitis responds readily to zinc salts. These may be applied in the form of boric lotion with zinc sulphate, gr. ii. to ℥ i., or as drops, preferably

the former. It is not known how zinc salts act in these cases, for the diplobacillus will grow in cultures containing them. Boric, zinc oxide, or ichthyol (2—5 per cent.) ointment is applied to the lids at night.

Follicular Conjunctivitis occurs frequently in children and young adults, both eyes being affected. It is characterised by the formation of small round or oval translucent bodies, 1 mm. or 2 mm. in diameter, in the lower fornix (Plate IV., Fig. 2); they are less commonly seen in the upper fornix, especially near the outer and inner canthi, never on the plica semilunaris or bulbar conjunctiva. They are raised above the surface, are often arranged in parallel rows, and consist of localised aggregations of lymphocytes—follicles, sometimes wrongly termed granulations—in the sub-epithelial adenoid layer.

Microscopically follicles are indistinguishable from the solitary lymph patches in the intestine, and often also from the follicles of trachoma (*q.v.*). They do not occur in the normal conjunctiva in man. The conjunctiva is not reddened or swollen. They persist for an indefinite time, causing few symptoms, and disappear without leaving any trace, such as scarring (cf. Trachoma).

Follicular conjunctivitis is usually due to overcrowding and living in badly ventilated rooms, especially schoolrooms, but also occurs among better class school children. Isolated follicles may occur in the outer part of the lower fornix in any chronic conjunctivitis of long standing. They may be due to overuse of atropin or eserin (*vide* p. 176). The children are seldom robust, and adenoid vegetations in the throat, which are of a similar nature, are often present. The disease is probably not contagious, and never develops into trachoma, as has been held by some good observers.

The symptoms are slight, consisting chiefly of slight irritation of the eyes, worse in bright lights and after near work.

Treatment.—Follicular conjunctivitis seldom requires local treatment. A weak astringent lotion may be ordered, and yellow oxide of mercury introduced within the lids once or twice a day. If the follicles are very large painting with silver nitrate solution will do good; single follicles may be touched with the alum pencil. Atropin, if in use, should be stopped or replaced by an equivalent mydriatic (*vide* p. 176).

Special attention should be directed to the refraction, and any errors corrected.

The general health and surroundings must be investigated and put upon a sound basis.

Trachoma (*Syn.—Granular Conjunctivitis*) is a much more serious form of folliculosis, which is responsible for the blinding of enormous numbers of people in places where it is endemic Both eyes are almost always affected. It may be stated at once that in England it is a rare disease except where large numbers of Irish or aliens are herded together.

In making a diagnosis the relative frequency of various forms of disease should always be borne in mind. It is a truism, but it is often disregarded, that an unusual type of case is less likely to be a rare disease than an unfamiliar manifestation of a common one. A better class child with follicles in the conjunctiva is most unlikely to be a case of trachoma, whereas, if the child goes to a board school in the east end of London, where there are large numbers of trachomatous aliens, the distinction of follicular conjunctivitis from trachoma becomes a matter of great difficulty.



Fig. 103. — Trachoma. (Nettleship.) *a*, trachoma follicles; *b*, scar in typical position, parallel with the edge of the lid.

The changes met with in the conjunctiva in trachoma are of two types, which are often present simultaneously. The *papillary* type is not specially characteristic; it is usually a more definite form of the papillary enlargement and congestion which is met with in other severe forms of conjunctivitis. The conjunctiva covering the upper tarsus is most affected, and appears red and velvety. This condition may pass into one with more uniform jelly-like thickening. Only in the comparatively infrequent cases in which no follicles can be seen will the true disease pass wholly unsuspected.

The *follicular* type (Fig. 103) manifests itself in the presence of follicles in the conjunctiva. When small they cannot be distinguished from the follicles of follicular conjunctivitis, and microscopical examination shows that they are fundamentally identical. They often, however, assume a size and appearance which is seldom or never seen in follicular conjunctivitis, but they differ most in having a characteristic distribution. The large follicles may be 5 mm. or 6 mm. in diameter. They are translucent and look like grains of boiled sago (" sago grains ").

The follicles usually commence in the lower fornix, but in most cases they quickly appear in the upper also. Unlike

what obtains in follicular conjunctivitis, they are not limited to the fornices. They often form a row along the upper margin of the upper tarsus, whence they invade the palpebral conjunctiva, appearing upon the tarsal surface, though they are less numerous in this situation. They are common about the caruncle, and may be seen on the plica semilunaris. Follicles do not occur on the upper tarsal conjunctiva in follicular conjunctivitis except at the inner and outer angles. They are very rare on the bulbar conjunctiva, but when seen here they are pathognomonic of trachoma.

The disease is very chronic and leads to much irritation, photophobia, lacrymation, &c., with some muco-purulent discharge. In certain districts abroad trachoma is endemic, and a very acute form is observed. It is very doubtful if true acute trachoma is ever seen in England: what usually passes for it is of quite different origin. It is due to the lowering of resistance of the trachomatous conjunctiva whereby it becomes specially liable to intercurrent attacks of other forms of acute conjunctivitis. This is facilitated by the irritation, which leads to rubbing of the eyes, so that contamination is readily brought about. The so-called acute trachoma met with in England is therefore chronic trachoma upon which an acute muco-purulent or purulent conjunctivitis has been engrafted. Even in Egypt the acute symptoms are most often caused by the Morax-Axenfeld diplobacillus, the Koch-Weeks bacillus, or the gonococcus (MacCallam).

Trachoma is an extremely contagious disease. Pathological anatomy reveals nothing characteristic, for it is doubtful if the recently discovered "trachoma bodies" are truly pathognomonic: there is lymphocytic infiltration of the whole of the adenoid layer of the parts of the conjunctiva affected. Special aggregations of lymphocytes, without a definite capsule, form follicles which are generally indistinguishable from those of follicular conjunctivitis. In late stages and large follicles the stroma and cells tend to become hyaline and gelatinous; sometimes the surface becomes broken and the contents are extruded into the conjunctival sac. In other cases a fibrous capsule forms around the follicles, which thus become isolated: more and more fibrous tissue is laid down, giving rise to cicatricial bands such as are never formed in follicular conjunctivitis, and are very characteristic.

Bacteriology has failed to elucidate the pathology of the disease. Various specific organisms have been described, but all have failed to resist criticism.

Trachoma is endemic in many parts of the world, *e.g.*, Russia, Poland, East Prussia, parts of Austria-Hungary, Egypt, Syria, Persia, China and Japan. It shows a predilection for certain races, *e.g.*, Irish, Jews, but it is not a racial disease; the predilection depends upon the mode of life of the individuals. Extended observation militates against the view that any race is exempt, though it is certainly uncommon among negroes.

The disease flourishes among people who are crowded together in unhealthy rooms—in armies, navies, asylums, workhouse schools, &c. –wherever the lower classes are herded together. Children and debilitated adults are most susceptible, but the robust are not exempt. It is commoner in low-lying, damp districts.

The disease is spread by transference of conjunctival secretion by means of fingers, towels, &c. The presence of much discharge, whether of true trachomatous origin or due to intercurrent conjunctivitis, increases the liability to contagion. On the other hand, scrupulous cleanliness suffices to prevent the extension of the disease to healthy subjects.



Fig. 104.—Trachomatous pannus. (Nettleship.)

Trachoma in the early stages may be easily mistaken for a simple chronic conjunctivitis. This error will be avoided if it be made an invariable rule in all cases of conjunctivitis to evert the upper lids and examine the upper fornices (*vide* p. 76).

Complications.—While trachoma very rarely affects the bulbar conjunctiva, it not infrequently attacks the cornea.

Trachomatous pannus is a lymphoid infiltration, with vascularisation, of the margin of the cornea, usually limited to the upper half (Figs. 104, 70 C), but tending to spread towards the centre and to involve the whole cornea. The upper part of the margin of the cornea becomes cloudy, and minute superficial vessels, springing from the corneal loops, grow inwards towards the centre. The haziness and vascularisation increase until the upper half of the cornea is affected. The vessels are all superficial (*vide* p. 86), and microscopic examination has shown that they lie at first between Bowman's membrane and the epithelium. They carry in with them a small amount of granulation tissue. In later stages Bowman's membrane

disappears and the superficial layers of the substantia propria become involved.

In progressive pannus the vessels are mostly parallel to each other and directed vertically downwards, anastomosing little. They extend to a level which forms a horizontal line, and beyond this line there is a narrow strip of infiltration and haze. In regressive pannus, on the other hand, the vessels extend a short distance beyond the area which is infiltrated and hazy : this difference is useful in estimating the results of treatment.

In more severe cases the vascularisation is not limited to the upper part, but superficial vessels grow in from all sides and the whole cornea becomes vascularised and opaque.

Pannus is not due to the rough upper lid rubbing upon the cornea. This is doubtless a predisposing factor, but in many conditions in which the lids are rough from some other cause pannus does not occur. It is not due to continuity, since the bulbar conjunctiva from the limbus to the fornix is unaffected. It is induced by contiguity, probably by direct infection, and this part of the cornea is most affected because it is covered by the lid both day and night.

It may resolve completely, leaving the cornea quite clear, but only in cases treated early, when the vessels have not yet destroyed Bowman's membrane. In other cases a permanent opacity results. Occasionally the corneal substance becomes weakened so that the cornea bulges under normal intraocular pressure and ectasia follows (keratectasia).

Corneal Ulcers are commonest at the advancing edge of the pannus. They are shallow, little infiltrated, and very irritable, causing much lacrymation and photophobia. Indolent central ulcers may form, or there may be ulcers in any part of the cornea, but especially over the pannous area.

Sequelæ.—Apart from the results of pannus and corneal ulceration the most malign effects of trachoma are caused by distortions of the lids. A peculiar drooping of the upper lids is very characteristic (*trachomatous ptosis*). It gives a sleepy appearance to the patient. There is always some scarring (Fig. 103), and when this is extensive the shape and position of the lids, especially the upper, are altered. Pressure on the everted upper lid will cause the appearance of white bloodless areas which may be mistaken for trachomatous scarring. They are easily distinguished by relieving the pressure. Through the great swelling of the conjunctiva the lids may be turned outwards (ectropion). In the late stages the follicles invade the tarsus, causing softening and absorption of the

dense fibrous tissue ; through the later contraction of the new-formed scar tissue the lids may be turned inwards (entropion), causing the lashes to rub against the cornea (trichiasis), &c. (*See* Chap. XXXI.)

Treatment. If there is much discharge the case must first be treated with a view to diminishing it and reducing congestion. The lids should be painted once daily with silver nitrate solution ; sublimate lotion should be thoroughly used three or more times a day, and boric ointment applied to the edges of the lids at bed-time.

In a few days the conjunctivitis will be much less, and an attempt must be made to get rid of the follicles. In relatively slight cases this is best effected by scouring the conjunctiva with a smooth crystal of copper sulphate. The crystal is fixed in a wooden holder and is pointed at the end. The lids are everted, and the point of the copper stick is pushed well up into the upper fornix and moved from one side to the other, the lid being lifted away from the globe by the stick during the manœuvre so as to avoid touching the cornea. The stick is then rubbed firmly over the whole of the palpebral conjunctiva.

The application of the copper stick is very painful, especially during the first few applications. Cocain may be previously instilled, but it does not prevent the intense smarting. It is important to start with the upper fornix, since this is most affected and most difficult to reach ; if it is not done first blepharospasm and the struggles of the patient will make it impossible afterwards. It is useless to apply copper too gently ; it must be firmly rubbed into the conjunctiva. Some of the sulphate dissolves in the tears, and should be mopped up with a pad of dry wool, since it is very irritating to the cornea. Pannus is no contraindication to the use of the copper stick, but quite the reverse ; copper sulphate applied to the lids is the best treatment for pannus. The pannus itself is not treated directly at all. On the other hand, corneal ulceration of any kind or degree is an absolute contraindication ; in these cases, even if pannus is present, reliance must be placed on silver nitrate until the ulcers are healed. Unfortunately silver has little specific influence over the trachomatous process, whereas that of copper is undoubted.

The copper stick must be applied once daily, otherwise the case is certain to run on indefinitely. Sublimate lotion is ordered for home use as before. Active treatment should be carried out for many weeks and should be continued for

several weeks after apparent cure. In the later stages copper sulphate drops (gr. i. ad $\overline{3}$ i.), or copper sulphate ($\frac{1}{2}$—1 per cent.) or copper citrate (10 per cent.) ointment may be used.

In most cases of trachoma the treatment advised will suffice to bring about that condition of amelioration which is usually described as cure. Relapses are common, occurring sooner or later, according to the length and assiduity of treatment. I am doubtful myself whether trachoma is ever really cured by the methods adopted.



FIG. 105.—Graddy's forceps.

In severer cases more drastic remedies must be employed. A method seldom employed now is to instil an infusion of jequirity (Abrus precatorius), which sets up a violent reaction. Jequiritol, the active principle, can be standardised and its action controlled by an anti-body. This treatment is more risky and probably not more efficacious than the methods about to be described. Probably the best of the stronger applications is a concentrated solution of perchloride of mercury in glycerine. It has been used as strong as 4 per cent.; under these circumstances it acts as a caustic. It is painted on the fornix and everted lids. This treatment is almost unbearably painful, and is followed by intense reaction, the conjunctiva and lids becoming enormously swollen. The pain lasts for several hours, gradually diminishing in intensity. Ice, or lint wrung out in iced water, should be applied to the lids immediately after the application. There is no doubt that much benefit is derived from the treatment.

The follicles may also be destroyed by touching them with a solid stick of carbon dioxide snow. Care must be taken to allow the tissues to thaw before the everted lid is replaced. X-rays and radium have also been used, but do not give better results than the ordinary treatment.

When the follicles are numerous and very prominent the treatment is shortened by attacking them mechanically. It may be done by various forms of scarification or expression. In performing any operation upon trachomatous patients

protective goggles must be worn by the surgeon and the immediate attendants.

The conjunctiva is first thoroughly cocained. Scarification may be performed with the finger nails—an objectionable practice—or by a knife, needle, sharp spoon or stiff tooth-brush. The follicles may also be destroyed by the galvano-cautery or by electrolysis.

FIG. 106.—Knapp's roller forceps.

In *expression* one of the many forms of expressor is used—*e.g.*, Graddy's forceps (Fig. 105), Knapp's roller forceps (Fig. 106). The upper lid is everted, and one limb of the forceps is pushed up into the fornix, the other being laid upon the palpebral conjunctiva. The two limbs are then pressed together with moderate force and are drawn in a horizontal direction from one end of the fold to the other. The fold of the fornix should in this manner be thoroughly mangled, and all the follicles squeezed out. Some recommend the immediate application of silver nitrate or sublimate solution (1 in 2000), but it is best simply to apply cold compresses, since there is always considerable reaction.

If the follicles in the upper fornix are very large and closely packed it is well to commence treatment by excising the fornix. There is always a redundance of tissue here, and no evil results ensue. The upper lid is everted doubly (*vide* p. 76) so as to expose the retro-tarsal fold completely. A silk suture is then passed through the fold at each end. By dragging on the sutures the whole fold is drawn out. It is then excised with scissors. If the tarsal plate is much diseased or distorted it also may be excised.

Pannus requires no special treatment, but if it fails to respond a *peritomy* may be performed. In this operation a collar of conjunctiva, 5 mm. broad, is excised round the corneal margin. The raw surface may be seared with the galvano-cautery, the object being to destroy the vessels and prevent their re-formation. Corneal ulcers must be treated on general principles (*vide* p. 194).

Hygienic treatment must be carried out so far as practicable.

The influx of aliens into the east end of London has made trachoma a serious menace to our own population. There is evidence that the operation of the Aliens Immigration Acts has reduced the prevalence of the disease: *e.g.*, the number of school children treated in the Metropolitan Asylums Board's trachoma schools has diminished.

Trachoma, then known as Egyptian ophthalmia, was spread far and wide in Europe by the French armies during the Napoleonic wars. Brilliant success attended the efforts to deal with trachoma among the Chinese labourers during the Great War, and not only was there no dissemination of the disease, but most of the cases were alleviated or cured.

Tubercle of the Conjunctiva is rare: it nearly always produces ulceration. Conjunctival ulceration should always suggest either the presence of an imbedded foreign body or a tuberculous or syphilitic lesion.

Fig. 107.—Tubercle of the conjunctiva. (After Eyre.)

Tubercle occurs in several forms: (1) small miliary ulcers usually on the palpebral conjunctiva; (2) granules on the palpebral conjunctiva resembling trachoma follicles (Fig. 107); (3) gelatinous cocks-comb-like excrescences on the fornices; (4) polypoid pedunculated outgrowths. The conjunctiva may be affected by extension of lupus from the face.

The preauricular gland is often enlarged and may suppurate. The disease is chronic, and the ulcers are indolent. The patients are usually young, and often free from clinical signs of tuberculous disease elsewhere in the body. There is very little doubt that the lesion may be the primary seat of tubercle, the bacilli being inoculated into minute abrasions, which are probably always present in the conjunctiva, caused by dust. There is little pain or irritation unless the ulceration is extensive.

It is not improbable that the second type is frequently mistaken for trachoma, and it is possible that it is cured by

the treatment founded on the wrong diagnosis. I have seen cockscomb tubercle in the lower fornix associated with tuberculous disease of the lacrymal sac in a child.

Pathology.—Scrapings may show tubercle bacilli. Sections show typical giant-cell systems. In doubtful cases inoculation experiments should be made. If a piece of tuberculous tissue is introduced into the anterior chamber of a rabbit's eye a typical tuberculous iritis will ensue in two or three weeks. Intraperitoneal inoculation of guinea-pigs is more satisfactory. Von Pirquet's reaction or injections of old tuberculin may be tried.

Treatment.—The disease should be eradicated, more especially as being often the primary focus. The affected conjunctiva should be excised, or if this is not feasible, thoroughly scraped and cauterised.

Injections of the latest form of tuberculin have given encouraging results, and subconjunctival injections of 2 per cent. guaiacol cacodylate may be employed. The application of 50 per cent. solution of lactic acid has been advocated.

Extension to the globe is probably rare. If prolonged treatment fails and the disease spreads it may be necessary to excise the eye.

Syphilis rarely manifests itself in the conjunctiva in the form of a primary chancre or gummatous ulceration. In the former case it may be due to the removal of a foreign body with the tongue. Ulceration of the palpebral or still more of the bulbar conjunctiva is always suggestive of the condition. Scrapings should be taken and examined for spirochætes, and the Wassermann test should be applied. (*See* Chap. XXXI.)

Conjunctivitis caused by Drugs (Atropin Irritation, &c.).—Some people are particularly susceptible to atropin, and more rarely to eserin and other drugs, locally applied to the conjunctiva. The lids become swollen, tense, and red, in fact erysipelatous. Examination of the conjunctiva will often show follicles, and even rarely a membrane. The cause is obscure : sometimes the condition seems to be due to a special solution or brand of the drug, owing to some impurity. Workers with crysophanic acid suffer from conjunctival irritation, and prolonged internal administration of arsenic causes the same effect. The dust of teak wood is particularly irritating to the conjunctiva. Conjunctivitis is also caused in some people by contact with horses or cats, or certain flowers, especially *Primula obconica*.

Treatment.—In atropin irritation the drug should be avoided.

If a mydriatic is imperative, some other should be substituted, *e.g.*, duboisin sulphate, 0·5 per cent.; scopolamin hydrobromide, 0·25 per cent. to 0·5 per cent., or hyoscin, 0·5 per cent., may be used, but toxic symptoms sometimes occur. It is said that atropin may be continued if it is scrupulously kept away from the skin by covering the lids with glycerine or ointment up to the margins.

The conjunctiva soon recovers after cessation of the cause, but astringent lotions accelerate the cure.

Action of Caustics.—See p. 396.

Some Rarer Forms of Conjunctival Inflammation.—One or two rarer forms of inflammation of the conjunctiva demand passing notice since they may be mistaken for those already described.



Fig. 108.—The palpebral form of spring catarrh.

Spring Catarrh (*Syn.*—*Vernal Catarrh*).—This is a recurrent conjunctivitis occurring with the onset of hot weather, and therefore rather a summer than a spring complaint. It is found in young people, usually boys, associated with the usual symptoms of mild conjunctivitis. Burning, itching, some photophobia, and lacrymation are the chief symptoms. Both eyes are affected. In the cooler months the condition subsides and gives no trouble, but recurs with the return of heat. The disease is met with amongst all classes, is sporadic, and non-contagious. It has been attributed on insufficient grounds to the action of the actinic rays of the spectrum.

Two types of objective signs are met with: (1) the palpebral form; (2) the bulbar form: both may be combined, but this is relatively rare. The palpebral form is easily recognised if typical. On everting the upper lid the palpebral conjunctiva is seen to be hypertrophied and mapped out into polygonal raised areas, not unlike cobble stones (Fig. 108). The colour is bluish white, like milk, and this appearance is seen also over the lower palpebral conjunctiva.

The flat-topped nodules are hard, and consist chiefly of dense

fibrous tissue, but the epithelium over them is thickened, giving rise to the milky hue. In vertical section they resemble circumvallate papillæ. Eosinophile leucocytes are present in them in great numbers and may be found in the secretion.

The palpebral form cannot be mistaken if typical, but it may resemble trachoma. The type of patient, the milky hue, the freedom of the fornix from implication, and the characteristic recurrence in hot weather will usually prevent mistake.

The bulbar form is less characteristic. In it there is a wall of thickening at the limbus, more gelatinous in appearance, and also milky. It may be mistaken for phlyctenular conjunctivitis.

In both forms the lesions persist during the cold months, though they are less marked.

Serious complications never supervene, and the ultimate prognosis is good. Occasionally a peculiar corneal opacity, resembling arcus senilis in having a clear zone between it and the limbus, is left, and some thickening and discoloration of the conjunctiva may remain.

Treatment is purely symptomatic. Well-fitting goggles with tinted glass should be worn. The irritation is best relieved by very weak acetic acid, gt. i. to ʒ vi. Adrenalin gives temporary relief. Gentle massage with the upper lid after application of yellow oxide of mercury ointment or 1—2 per cent. guaiacol ointment is beneficial. Astringents are harmful: boroglyceride has been recommended, with arsenic internally. Radium applications have given relief in some cases. Excision of the nodules, sometimes advised, seems to be useless. The general health should be attended to, and adenoids and enlarged tonsils, if present, removed.

Ophthalmia nodosa is a nodular conjunctivitis which may be mistaken for tubercle—pseudo-tuberculous disease of the conjunctiva. It is due to the irritation of the hairs of certain caterpillars, and therefore always commences in the late summer months. Small semitranslucent, reddish, or yellowish grey nodules are formed in the conjunctiva and sometimes in the iris. On microscopical examination hairs surrounded by giant cells and lymphocytes are found.

Treatment.—The nodules in the conjunctiva should be excised. Otherwise the condition is treated on general principles.

Parinaud's Conjunctivitis may be mistaken for tubercle or trachoma. Usually one eye only is affected. Granulations occur on the tarsal conjunctiva or fornices, and the preauricular and submaxillary glands are enlarged and may suppurate. The disease commences with slight constitutional disturbance and may last for months. It has been attributed to bovine tubercle and to a leptothrix (Verhoeff).

Pemphigus of the conjunctiva is a rare but very serious disease. Vesicles occur, but more commonly greyish white membranous patches. Progressive cicatrisation of the conjunctiva follows, leading eventually to *essential shrinking of the conjunctiva*, with consequent opacification of the cornea from malnutrition. Vesicles may be found in the nose, mouth and pharynx, but rarely in the skin. Treatment, such as transplantation of mucous membrane, is unavailing.

Electric Light Ophthalmia (Photophthalmia).—The ultra-violet rays of the electric light may cause extreme burning pain, lacrymation, photophobia, blepharospasm, and swelling of the palpebral conjunctiva and retrotarsal folds, coming on a few hours later. The condition is generally caused by the bright flash of a short circuit, but may result from exposure to a naked arc light, as has happened, for instance, in cinema studios. It is rarely due to exposure to enclosed arc or other lights, since the glass globe absorbs the most deleterious of the ultra-violet rays.

Treatment.—Cold compresses, dark glasses, and astringent lotions.

Snow Blindness.—The cause and symptoms of snow blindness are the same as of electric light ophthalmia, viz., exposure to ultra-violet rays, especially from 310 $\mu\mu$ to 290 $\mu\mu$, an unusually large percentage of which is reflected from snow surfaces. Smoked or orange tinted (not blue) glasses should be used as a prophylactic measure, and they are most efficacious when made with Crookes's glass (*vide* p. 142). The treatment is the same as for electric light ophthalmia.

Degenerative Changes in the Conjunctiva.

Concretions (*Syn.*—"*Lithiasis*").—Concretions occur as minute hard yellow spots in the palpebral conjunctiva. They are due to the accumulation of epithelial cells and inspissated mucus in depressions which are called Henle's glands. They never become calcareous, so the term is a misnomer, but they are so hard that when they project from the surface they scratch the cornea and give the sensation of a foreign body in the eye. They are common in elderly people. There is no reason to attribute them to gout, but uric acid deposits have been observed in the palpebral conjunctiva of gouty patients. Concretions should be removed with a sharp needle.

Pinguecula is a triangular patch on the conjunctiva, found usually in elderly people, especially those exposed to dust,

wind, and so on. It occurs in the direction of the palpebral aperture, the apex of the triangle being away from the cornea. It affects the nasal side first, then the temporal. It is yellow in colour and looks like fat, whence the name (pinguis, fat). It is not due to fat, but to hyaline degeneration of the connective tissue and an excessive development of yellow elastic fibrous tissue. It is particularly conspicuous when the eye is inflamed, since the pinguecula remains relatively free from congestion : mistakes in diagnosis may then occur. It requires no treatment, but may be removed if the disfigurement is great.

Pterygium (*πτέρυξ*, a wing).—This is a peculiar encroachment of the conjunctiva on the cornea (Fig. 109). It is triangular in shape, and when single is always upon the nasal



Fig. 109.—Pterygium.

side ; when double the temporal one has developed later. It is derived from pinguecula (Fuchs). It must be carefully distinguished from *pseudo-pterygium*, which is due to the tip of a fold of chemotic conjunctiva becoming adherent to an ulcer within the corneal margin. It may occur at any part of the cornea. The conjunctiva then forms a bridge over the limbus, and a fine probe can always be passed beneath it, which is not possible with a true pterygium.

The apex of the pterygium is usually blunt ; there is no ulcer in the cornea beyond it, as formerly described, but there are often small opacities. In the early stage the pterygium is thick and vascular ; it advances over the cornea and may reach the pupillary area and interfere with vision. When it ceases to grow it becomes thin and pale, but never disappears.

The true pterygium is a single layer of conjunctiva, adherent in its whole length to the sclerotic and cornea, though only loosely, except at the apex. The area of adhesion is always smaller than its breadth, so that there are folds at the upper and lower borders.

Pterygium is not due to a fold of conjunctiva being dragged across the cornea by a progressive ulcer, as was once taught. It is probably due to malnutrition of the cornea, resulting from the pinguecula. Granulation tissue grows in and invaginates the epithelium, destroying Bowman's membrane. When it ceases to progress dense fibrous tissue is formed.

Pterygium is rare in England, but is common in dry climates with sandy soils, *e.g.*, parts of Australia, South Africa, Texas, &c. It was common in soldiers during the Boer War, and it is not infrequently seen in sailors.

Treatment.—Pterygium is best left alone unless it is progressing rapidly towards the pupillary area, or is very disfiguring. The latter reason is not of much weight, since it cannot be removed without leaving a scar.

The apex of the pterygium may be destroyed with the actual cautery. Removal is effected by seizing the neck, near the corneal margin, with fixation forceps, raising it, and shaving or dissecting the apex from the cornea. Care must be taken not to go too deep. The pterygium is freed from the sclerotic for about half the distance towards the canthus. Two converging incisions with scissors separate the apex and greater part of the body. The conjunctiva is then freed from the sclerotic above and below so as to admit of the two edges being sutured together.

Pterygium sometimes recurs after removal. This may be only apparent, owing to vascularisation of the denuded area. If it actually re-forms and extends towards the pupillary area, the apex should be turned down under the bulbar conjunctiva and sutured in this position (McReynolds).

Symptomatic Conditions.

Subconjunctival Ecchymosis, due to the rupture of small vessels, frequently occurs. It may be the result of direct injury, or, more commonly, occur spontaneously. Very minute ecchymoses, or possibly thromboses, are seen in severe conjunctivitis, especially pneumococcic. Larger ones accompany severe straining, especially in old people, *e.g.*, on lifting heavy weights, vomiting, &c. In these circumstances they

indicate a weakness of the vessel walls, and should be regarded as a danger signal. Other signs of arteriosclerosis should be sought in the fundus oculi and elsewhere, and, if found, warning of the possibility of cerebral hæmorrhage given, with appropriate instructions for its avoidance. Very frequently no such signs can be discovered, and the condition, though unsightly, is trivial. Subconjunctival ecchymoses are not infrequently seen in children with whooping cough ; as a rule they need arouse no anxiety, but retinal hæmorrhages and hyphæmia also occur in association with them, as well as cerebral hæmorrhage.

More serious are the large subconjunctival ecchymoses which sometimes follow blows or falls on the head. They may then be due to extravasation of blood along the floor of the orbit, the result of a fractured base. In fractures of the sphenoid the blood appears later on the temporal side than elsewhere. Hæmorrhage also results from severe or prolonged pressure on the thorax and abdomen, as in persons squeezed in a crowd.

Subconjunctival hæmorrhages may occur in scurvy, purpura, malaria, and so on.

The importance of subconjunctival hæmorrhage is always symptomatic, since the eye itself is never endangered.

Treatment. The blood becomes absorbed in from one to three weeks without treatment. Boric lotion is usually ordered as a placebo. Dionin may be used to accelerate absorption in young patients, but should not be employed in elderly subjects with arteriosclerosis.

Chemosis, or œdema of the conjunctiva, may occur in (1) acute inflammations ; (2) in cases of obstruction to the lymph flow ; (3) in abnormal blood conditions.

In the first group of cases the inflammation may be in the conjunctiva, *e.g.*, gonorrhœal conjunctivitis, or within the eyeball, as in panophthalmitis, hypopyon ulcer. It is also found in acute glaucoma. The inflammation may be in the accessory structures of the eye, *e.g.*, a stye, a parasitic bite on the lid, dacryocystitis, periostitis, orbital cellulitis, cerebro-spinal meningitis. The chemosis of dionin is probably due to a specific irritation of the conjunctival vessel walls.

In the second group the pressure of an orbital tumour may so interfere with the lymph and blood streams as to produce chemosis, and it is found in pulsating exophthalmos.

To the third group belong Bright's disease and the anæmias. It is sometimes due to urticaria or angioneurotic œdema.

Recurrent chemosis is sometimes associated with menstruation, and it has been observed in trigeminal neuralgia and migraine.

Xerosis (*Syn.*—*Xerophthalmia*) (ξηρός, dry) is a dry, lustreless condition of the conjunctiva which occurs in two groups of cases: (1) as a sequel of local ocular affection; (2) associated with general disease.

The first type is a cicatricial degeneration of the conjunctiva —(a) following trachoma, burns, pemphigus, diphtheria, &c., commencing in isolated spots, ultimately involving the whole conjunctiva and cornea; (b) following exposure, due to ectropion or proptosis, whereby the eye is not properly covered by the lids. As the result of the rare affection of the conjunctiva with pemphigus the cicatricial contraction of the conjunctiva may be extreme and progressive, a sort of keloid condition being induced (*essential shrinking of the conjunctiva*): the lids may then be quite adherent to the globe, the cornea being opaque like skin (*vide* p. 179).

In the other group of cases xerosis occurs in a mild form, found in children, usually boys, accompanied by night blindness (Chap. XIX.), and characterised by small triangular white patches on the outer and inner sides of the cornea, covered by a material resembling dried foam, which is not wetted by the tears (Bitot's spots). The cases usually occur during the summer months, and the children are not conspicuously ill-nourished. A similar mild form, also associated with night blindness, is met with in adults in some countries, *e.g.*, India, but seldom if ever in England. A severe form is found in marasmic children, associated with keratomalacia (*vide* p. 213) and necrosis of the cornea.

Pathology.—The chief changes are in the epithelium, which becomes epidermoid, *i.e.*, exactly like that of skin, with granular and horny layers. The foamy spots are due to horny epithelium, which becomes cast off into the conjunctival sac and accumulates in the lower fornix. Owing to this change the epithelium ceases to secrete mucus. It becomes dry; a certain amount of vicarious activity is set up in the Meibomian glands (Chap. XXXI.), which cover the surface with their fatty secretion. The watery tears then fail to moisten the conjunctiva. The so-called xerosis bacilli, which are pseudo-diphtheria bacilli, grow profusely under these conditions, but they have no causal relationship and are of no importance.

It is to be noted that xerosis has nothing to do with any failure of function on the part of the lacrymal apparatus. The conjunctiva can be quite efficiently moistened by its own

secretions alone. If the lacrymal gland is extirpated xerosis does not follow. If on the other hand the secretory activity of the membrane itself is impaired xerosis follows, in spite of normal or increased lacrymal secretion.

Treatment.—Xerosis is a symptom, and its treatment must therefore be purely symptomatic. Local treatment consists in relieving the dryness with boric acid ointment, olive oil, milk, weak alkaline solutions, &c.: smoked glasses should be worn.

In the marasmic cases restoration of normal nutrition is the all-important aim, though the patients have often gone too far.

Argyrosis is the staining of the conjunctiva from prolonged application of silver salts (nitrate, protargol, &c.) for the treatment of chronic conjunctivitis, and especially trachoma. The conjunctiva, particularly the lower fornix, is stained deep brown. The staining is due to the impregnation of the elastic fibres in the membrane and vessel walls with reduced metallic silver. The condition is very difficult to get rid of, but subconjunctival injections of potassium iodide (℥ iij.—iv. of a 30 per cent. solution) and internal administration of hexamin (urotropin) over a prolonged period have been advocated.

Cysts and Tumours.

The only common **cysts** found in the conjunctiva are due to dilatation of lymph spaces. When small these often form rows of little cysts on the bulbar conjunctiva (*lymphangiectasis*). Occasionally single, though multilocular, cysts occur (*lymphangioma*). Larger retention cysts of Krause's accessory lacrymal glands occur in the upper fornix (Chap. XXXI.). Subconjunctival cysticercus and hydatid are rare in England. Non-parasitic cysts require simple removal of the anterior walls. *Epithelial implantation cysts* occur rarely after injuries or operations, *e.g.*, tenotomy, and may burrow into the cornea.

Tumours of the conjunctiva have all a tendency to be polypoid, owing to the perpetual movements of the globe and lids.

Papillomata occur at the inner canthus and in the fornices. In the latter position they may be mistaken for the cockscomb type of tubercle, but the individual leaflets may be separated by a probe. They also occur sometimes at the limbus in old people and are then liable to become malignant. They should be removed.

Simple Granulomata, generally polypoid, often grow from tenotomy wounds or the sites of foreign bodies. They consist of

ordinary exuberant granulation tissue. They are common in empty sockets after excision, and at the site of chalazions which have been insufficiently scraped (Chap. XXXI.). They should be removed by scissors.

Fibromata, also generally polypoid, occur in sockets. They may be soft or hard, and require simple removal.

Nævi or *congenital moles* are not uncommon. They are white gelatinous or pigmented nodules situated by preference at the limbus or near the plica semilunaris. They have the same structure as in the skin—groups, often alveolar, of "nævus cells" in close connection with the epithelium. They are congenital and tend to grow at puberty, rarely becoming malignant. They should be excised. Pigmentation at the limbus occurs normally



Fig. 110.—Dermoid of the conjunctiva, in a somewhat unusual situation. Note the hairs.

in dark races, and patches in this situation are not uncommon in people with dark complexions.

Dermoids are lenticular yellow tumours, usually astride the corneal margin, most commonly at the outer side (Fig. 110). They are often wrongly called dermoids of the cornea. Not infrequently there is a notch in the upper lid corresponding with the position of the tumour. They consist of skin in an abnormal situation, with epidermoid epithelium, hairs, sebaceous glands, &c. They are congenital and tend to grow at puberty; the hairs also grow and often cause irritation. Dermoids should be dissected off the globe if troublesome, though their removal does not produce much improvement in appearance, as the site of attachment to the cornea remains densely opaque.

Dermo-lipomata or *fibro-fatty tumours* are congenital tumours found at the outer canthus in babies. They consist of fibrous

tissue and fat, sometimes with dermoid tissue on the surface. They are not encapsuled. The main mass should be removed, but it will be found that the fat is continuous with that of the orbit: care must be taken not to injure the extrinsic muscles.

Sarcoma is rare. It occurs at the limbus, is usually pigmented, and most of the patients have been old. Sarcomata spread over the surface of the globe, but rarely penetrate it. Recurrence and metastasis occur as elsewhere in the body. They may be alveolar —derived from nævi— or round or spindle celled. They must be

Fig. 111.—Epithelioma of the conjunctiva, from a section (× 6).

removed as freely as possible and examined microscopically. On the slightest sign of recurrence the eye must be excised, and if recurrence again takes place the orbit must be exenterated. A diffuse spreading pigmentation of the conjunctiva occurs rarely in elderly people, and has been known to give rise to metastatic sarcomatous tumours. It should always be viewed with grave suspicion.

Epithelioma occurs par excellence where one kind of epithelium passes into another, hence in the eye chiefly at the limbus (Fig. 111), and at the edges of the lids. Papillomata in old people often take on malignant proliferation. Like sarcomata, epitheliomata spread over the surface and into the fornices, rarely penetrating the globe. They have the characteristic structure. The treatment is the same as for sarcoma: it is well in both cases to cauterise the base with the actual cautery after the first removal.

CHAPTER XI.

DISEASES OF THE CORNEA.

The special importance of diseases of the cornea depends upon the fact that they often leave permanent opacities which seriously lower the visual acuity, while the complications which not infrequently attend them may lead to the loss of the eye.

Inflammation of the Cornea (Keratitis).

Inflammation of the cornea may be purulent or non-purulent. An immense amount of research has been devoted to inflammation of the cornea from the earliest investigations of Bowman (1849) onwards, and much of our knowledge of inflammation in general is derived from these researches.

Purulent Keratitis, Ulceration of the Cornea.—Purulent keratitis is nearly always exogenous, *i.e.*, it is due to pyogenic organisms which invade the cornea from without. The first line of defence is the epithelium. It has been pointed out that the only organisms which are known to be able to attack normal epithelium are the gonococcus and the diphtheria bacillus; but many other bacteria are capable of producing ulceration, notably the pneumococcus.

When we remember the exposed position of the cornea it is not surprising that minute abrasions are extremely common. They are probably of everyday occurrence, but other factors have also to be reckoned with. Pathogenic organisms of high virulence are not always present in the conjunctival sac, and if they are, as is often the case, the resistance of the normal tissues has to be taken into account. It is too often forgotten that normal tissues with a healthy blood supply and lymph flow are well armed against any but the most virulent invaders. Even with these prolonged contact is usually necessary.

Apart from actual abrasions many causes of diminished resistance of the epithelium are met with. Such are drying, as in xerosis, necrosis due to deficient nutrition, as in keratomalacia, desquamation as the result of œdema or neuroparalytic keratitis. If cocain is instilled too freely, especially

if the lids are not kept closed in the intervals, the epithelium becomes dull and is finally thrown off. Hence this drug is to be used with caution and only under supervision ; it should seldom if ever be given in lotions for use at home.

Malnutrition affects not only the epithelium but also the whole cornea. In badly nourished corneæ ulceration is apt to be deep, leading to early perforation, and also extensive, resulting in widespread necrosis. Such septic ulcers, due to the attack of ordinary pyogenic organisms, occur after injuries in asthenic conditions, in keratomalacia, lagophthalmia, &c. Corneal ulcers are much commoner in the lower orders, doubtless owing to the defective resistance of the tissues and the more frequent exposure to injury.

In the commonest form of suppurative keratitis—the corneal ulcer—there is localised necrosis in the most anterior layers of the cornea. The sequestrum is partly disintegrated and cast off into the conjunctival sac, and partly adheres to the surface of the ulcer. Usually the epithelium is destroyed and cast off over an area considerably larger than the ulcer itself, and the same applies to Bowman's membrane. The epithelium, however, rapidly advances towards the ulcer, grows over its edge, and even over the slough or pus which forms the floor.

The ulcer is usually saucer-shaped, and the walls project above the normal surface of the cornea (Fig. 112), owing to the imbibition of fluid by the corneal lamellæ, which causes them to swell. The spaces between the lamellæ are packed with leucocytes for some distance around the ulcer, appearing as a grey zone of infiltration. This is the progressive stage.

A line of demarcation forms as in necrosis elsewhere in the body. The toxins are most concentrated near the centre where there are most organisms. A wall of polymorphonuclear leucocytes forms a second line of defence. At a certain distance the tissues are protected ; here the leucocytes are not paralysed or killed by toxins, but exert their digestive functions, macerating and dissolving the necrotic tissues. When the dead material has been thrown off the ulcer is somewhat larger, but the cloudiness has disappeared, the floor and edges are smooth and transparent, and the regressive stage is reached.

Meanwhile vascularisation has been going on (Plate V., Fig. 1). Minute superficial vessels grow in from the limbus near the ulcer. They supply the pabulum to restore the loss of substance : they also supply protective substances—anti-

FIG. 1.

FIG. 2.

PLATE V. (*To face p.* 189.)

Fig. 1.—Nebulæ, with recent vascular ulcer.

Fig. 2.—Phlyctenular pannus.

bodies—and therefore play an important *rôle* in combating bacterial infection. Sometimes they are so exuberant as to overstep the limits of utility, *e.g.*, in fascicular ulcer (*vide* p. 210).

When the ulcer has become vascularised, everything is prepared for cicatrisation, which is carried out exactly as in other connective tissues. The fixed connective tissue cells, here the corneal corpuscles, divide and form masses of nucleated spindle-shaped cells, over which the epithelium grows and is lifted to its normal level. The nuclei and vessels gradually disappear, and a mass of fibrous tissue is formed. The fibres are not arranged regularly like the normal lamellæ, so that they refract the light in various directions : the scar is, therefore, more or less opaque according to its thickness. If it is very large and dense some of the larger vessels persist permanently ; the smaller ones disappear. Bowman's mem-

Fig. 112.—Vertical section of a corneal ulcer, showing infiltration of the substantia propria

brane is never regenerated, and if it has been destroyed, as is the case in all but very superficial abrasions, some degree of permanent opacity remains.

During the progressive stage there is lacrymation, photophobia, and pain, owing to the exposure of the fibrils of the ophthalmic division of the fifth nerve. Some of the toxins elaborated by the bacteria diffuse through the cornea into the anterior chamber, just as atropin does when instilled into the conjunctival sac. Here they exert an irritative effect upon the vessels of the iris and ciliary body, so that hyperæmia of the iris, with or without definite ciliary injection, occurs. The irritation may be so great that leucocytosis takes place, and polymorphonuclear leucocytes are poured out by the vessels of the iris and ciliary processes. They pass into the aqueous and gravitate to the bottom of the anterior chamber, where they form an *hypopyon* (Fig. 128).

There are several important facts about hypopyon which

must be borne in mind. The pus cells do not come from the cornea, as was once thought. This is proved by the following facts : (1) Descemet's membrane is impermeable to leucocytes, though fluids readily pass through it ; (2) the cells sometimes contain pigment granules, obviously derived from the uveal tract ; (3) if the ulcer has not perforated the hypopyon is sterile. The last fact is of the greatest importance, both theoretically and practically. It shows that the leucocytosis is due to toxins, not to actual invasion of bacteria, which, indeed, are as incapable of passing through the intact Descemet's membrane as are leucocytes. It accounts for the ease and rapidity with which hypopyon is often absorbed : it may develop in an hour or two, rapidly disappear, and as readily reappear. Such hypopyons are very fluid, always moving to the lowest part of the anterior chamber if the position of the patient's head is changed. The fact that the hypopyon is sterile has great practical importance—it is unnecessary to remove the pus, as is the rule in all other parts of the body ; if the ulcerative process can be stopped the hypopyon will be absorbed.

The hypopyon may be so small that it is scarcely visible, being hidden behind the rim of sclera which overlaps the cornea. It may reach half way up the iris, having a flat upper surface in accordance with the laws of gravity. It may fill the anterior chamber, wholly obscuring the iris. The larger hypopyons are usually less fluid, owing to the formation of a fibrinous network which imprisons the leucocytes in its meshes. Such hypopyons are much less readily absorbed, and it may be necessary or advisable to evacuate them.

The scar tissue which replaces the destroyed portions of the cornea usually fills in the gap exactly, so that the surface is level. If it is thin the resulting opacity is slight and is called a *nebula* (Plate V., Fig. 1) ; if rather more dense it is sometimes called a *macula ;* if very dense and white it is called a *leucoma.* A thin, diffuse nebula covering the pupillary area interferes more with vision than a strictly localised dense leucoma, so long as the latter does not block the whole pupillary area. The reason is that the leucoma stops all the light which falls upon it (Fig. 113), whereas the nebula refracts it irregularly, allowing many of the rays to fall upon the retina, where they blur the image formed by the regularly refracted rays. An opacity does not necessarily prevent the light from being focussed upon the retina immediately behind it. Thus,

a central opacity of the cornea will not prevent the focussing of an object upon the macular region, for the rays passing through the clear peripheral parts of the cornea will be refracted towards the macula, only those rays being cut off which are incident to the corneal surface at the opaque region. There is thus a loss of brightness rather than of definition, though definition will also be impaired by the superposition of a diffuse entoptic image of the opacity upon the clear image of the external object.

When Bowman's membrane has been destroyed the opacity is permanent, but even then it tends to clear more or less. The younger the patient the more clearing may be anticipated. The deeper the cicatrix the less it clears ; perforating wounds remain permanently opaque. This fact is well illustrated by the punctures made by discission needles in needling cataract ;



Fig. 113.—Optical effect of a corneal opacity.

they remain as grey spots in the cornea throughout life. Vascularisation plays a part in the clearing of corneal opacities, as is shown by the fact that they clear first in the immediate vicinity of the vessels.

Extremely thin cicatrices may be almost or quite transparent. In them there is often deficient scar tissue formed so that the surface is flattened or even indented. Such corneal *facets* can only be seen by carefully examining the corneal reflex (*vide* p. 81), but they cause considerable defect of visual acuity.

From the same cause—deficiency of scar tissue—the cornea may be markedly thinner at the site of the ulcer than elsewhere. The scar may then bulge above the surface, owing to ineffectual resistance to the normal intraocular pressure. As the cicatrix becomes consolidated the bulging may disappear, or it may remain permanently as an *ectatic cicatrix* (keratectasia from ulcer).

Some ulcers, especially those due to the pneumococcus and septic organisms, extend rapidly in depth. There is then grave danger of perforation. The whole thickness of the cornea except Descemet's membrane and a few corneal lamellæ may be destroyed. Descemet's membrane, like other elastic membranes, offers great resistance to inflammatory processes. It is, however, unable alone to support the intraocular pressure: it therefore becomes protruded through the ulcer, appearing upon the surface as a transparent vesicle, which is called a *keratocele*. This may persist, surrounded by a white cicatricial ring, or it may eventually rupture.

Perforation and its effects.—When an ulcer perforates the aqueous suddenly escapes and the intraocular pressure sinks to zero, *i.e.*, to the atmospheric pressure. The iris and lens are driven forwards into contact with the back of the cornea.

The effect of perforation upon the nutrition of the cornea is good: owing to the diminution of intraocular pressure the diffusion of lymph through the cornea is facilitated. Ulceration usually ceases, pain is alleviated, and cicatrisation proceeds rapidly. The complications which attend perforation are however of extreme danger to sight and even to the preservation of the eye. These complications vary according to the position and size of the perforation.

Usually the perforation takes place opposite some part of the iris, which therefore lines the aperture when the aqueous escapes. The iris becomes gummed down to the opening by lymph, which gradually organises, and an *anterior synechia* is formed. The blocking of the perforation with iris allows the anterior chamber to be re-formed, fresh aqueous being rapidly secreted.

The aqueous often escapes very quickly owing to some sudden exertion on the part of the patient, *e.g.*, coughing, sneezing, straining at stool, or spasm of the orbicularis. Any such sudden exertion causes a rise in general blood pressure, which at once manifests itself by a rise in intraocular pressure. The weak floor of the ulcer is unable to support the sudden strain and gives way. In such a case, especially if the perforation is large, a portion of the iris is carried not only into the opening but through it, and a *prolapse of iris* is produced. The prolapse may not include the pupillary margin, in which case it is hemispherical; or the pupillary margin may also prolapse, a tag of iris lying free upon the cornea. In either case the colour of the iris soon becomes obscured by the deposition of grey or yellow lymph upon the surface. In large

prolapses the stroma becomes thinned and the black retinal pigment epithelium is thrown into relief.

Sometimes the whole cornea sloughs, with the exception of a narrow rim at the margin, and *total prolapse of iris* occurs. The pupil usually becomes blocked with exudate, and a false cornea is formed, consisting of iris covered by lymph.

If the perforation takes place very suddenly the suspensory ligament of the lens is much stretched. It may rupture partially, causing tilting or dislocation of the lens, or wholly, so that the lens may be expelled through the perforation.

If prolapse of iris has occurred cicatrisation may still progress. The lymph which covers the prolapse or pseudo-cornea becomes organised, and forms a thin layer of connective tissue over which the conjunctival or corneal epithelium rapidly grows. The contraction of the bands of fibrous tissue tends to flatten the protruding prolapse or pseudo-cornea. It rarely, however, becomes quite flat; more commonly the iris and cicatricial tissue are too weak to support the restored intraocular pressure, which is often increased, owing to the advance in position of the iris (*vide* p. 264). The cicatrix tends therefore to become ectatic, and such an ectatic cicatrix in which the iris is incarcerated is called an *anterior staphyloma* (Fig. 114). If the prolapse of iris is partial the resulting staphyloma will be partial; if total, a total staphyloma will ensue. The bands of scar tissue on the staphyloma vary in breadth and thickness, producing a lobulated surface, hence the name (σταφυλή, a bunch of grapes).



FIG. 114.—Anterior staphyloma, showing also an anterior capsular (pyramidal) cataract (R. L. O. H. Museum.)

If the perforation happens to be opposite the pupil it cannot

be covered with iris. The pupil often becomes adherent to the margin, and the aperture becomes filled with exudate. The anterior chamber is then re-formed very slowly: the lens remains long in contact with the ulcer, and permanent opacity may occur in it—*anterior capsular cataract* (Fig. 114). As the anterior chamber re-forms the exudate filling the opening is submitted to strain. It frequently ruptures, especially if the patient is restless. This process may be repeated again and again, so that the opening may become permanent—*corneal fistula.*

The sudden reduction of intraocular pressure when the perforation occurs removes support from all the intraocular blood vessels. They become dilated and may rupture, *intraocular hæmorrhage* taking place. The retinal vessels may rupture, causing vitreous hæmorrhage, or the choroidal, causing sub-retinal or sub-choroidal hæmorrhage. It may be so profuse that the contents of the globe are extruded with the outflowing blood: indeed, in very rare cases the hæmorrhage may endanger life, for it is most likely to occur in old people with atheromatous vessels.

Finally, the organisms which have caused the ulceration of the cornea may gain access to the interior of the eye as the result of perforation, the vitreous acting as an excellent culture medium. *Purulent irido-cyclitis* or even *panophthalmitis* may thus be set up, a result especially prone to occur in gonorrhœal ophthalmia and in hypopyon ulcer (*q.v.*).

Treatment of Uncomplicated Ulcers.—Cleanliness, heat, rest, and protection are the fundamental principles of the treatment of corneal ulcers. Surgical cleanliness is the principle which should regulate the use of lotions; heat is employed to prevent stasis and encourage repair; local rest is attained by the use of atropin; rest and protection from deleterious external agencies are aimed at in the use of bandages.

The ordinary treatment of a simple uncomplicated ulcer is as follows. The conjunctival sac is washed out carefully three or more times a day with a considerable quantity of a mild antiseptic lotion, which should be used as hot as can be borne comfortably (*vide* p. 146). It acts principally by washing away secretions and necrotic material, which carry with them many of the organisms and their toxins. It is impossible to apply antiseptics sufficiently strong to kill the organisms, hence it is of little importance whether weak sublimate lotion (1 in 8000) or simple boric lotion be used.

After each irrigation a drop of 1 per cent. atropin solution

or a small lump of 1 per cent. atropin ointment is introduced between the lids.

A protective bandage is then applied. It consists of a pad of sterile gamgee tissue, or a layer of cyanide gauze covered by a pad of cotton wool, kept in place by a bandage, firmly but not tightly applied. A simple tie bandage suffices. This is of sufficient length to pass round the head and tie behind. It passes obliquely above the sound eye, over the ulcerated eye, and under the ear of the side of the affected eye, where the bandage is given a single turn ; the ends are tied just above the occipital protuberance.

This treatment suffices for mild cases.

In more severe cases hot fomentations should be applied in the intervals between bathing with lotion, which should be done more frequently—every three or four hours. The compresses should be made of large round pads of plain or boric lint, on one surface of which gutta-percha tissue is sewn. The compresses are placed in a cloth and immersed in boiling water ; by keeping the ends of the cloth out of the water and turning them in opposite directions the excess of the water is wrung out without scalding the fingers. The compress is applied as hot as can be borne. It is at once covered with a large pad of hot cotton wool, and bandaged into position.

Atropin has a twofold function in the treatment of corneal ulcers. In the first place it keeps the eye at rest by paralysing the intrinsic muscles, both the sphincter iridis and ciliary muscle. In the second place it prevents most of the dangerous results of iritis (*vide* p. 241). Corneal ulcers are always accompanied by more or less iridic and ciliary hyperæmia, and actual inflammation often occurs.

Eserin has been recommended instead of atropin in the treatment of some ulcers. It has been held that prolapse is less likely to occur when a peripheral ulcer perforates if the pupil is contracted. This is a fallacy. Even with complete dilatation under atropin the pupil at once contracts when the aqueous escapes. The objections to eserin are that it prevents rest by keeping the sphincter iridis and ciliary muscle in a state of tonic contraction, that it irritates the iris and tends to increase iritis, with consequently greater risk of the formation of posterior synechiæ, and that it causes discomfort or even pain in the eye. It is never to be used for a simple ulcer, though it may have beneficial effects in special chronic types (*vide* p. 212).

Any contributory cause for ulceration must of course receive attention. Prominent among such causes are conjunctival

conditions and general malnutrition. Thus, the ulcers associated with trachomatous pannus will not heal if the lids are neglected (*vide* p. 171).

In purulent conjunctivitis and trachoma the lids should be painted with silver nitrate, even during the progressive stage. The copper stick must not be used in trachoma, owing to the irritative effect upon the cornea. The presence of much conjunctival discharge is a contraindication to the use of a bandage; the benefit derived from it is more than counteracted by the retention of secretions (*vide* p. 147). It must be replaced by a shade.

To restrain children from touching their bandages a straight splint may be applied to the front of the arm, reaching a little beyond the elbow, thus preventing flexion.



FIG. 115.—Lang's speculum. This form is preferable to the ordinary type, as it keeps the lashes out of the field of operation.

In debilitated adults or old people and marasmic children the building up of the constitution by good food, fresh air, and tonics is often more important than the treatment of the local condition.

When cicatrisation is complete and all irritative signs have passed off an attempt must be made to render the scar as transparent as possible. The results are usually disappointing, but cicatrices clear considerably in young patients, and in many others a gratifying improvement may be noticed in the course of months or years. Stimulating treatment is indicated, beginning with weak irritants and passing cautiously to stronger. Insufflations of finely powdered calomel may be used first. If it is well borne dilute yellow oxide of mercury ointment is employed. A small lump of the ointment, gr. iv. to ʒ i., is placed in the conjunctival sac and rubbed in gently

by rotatory movements of the upper lid by means of the finger. The massage with the lid should be employed three

Fig. 116.



Fig. 117.

Fig. 116.—Ordinary fixation forceps. Fig. 117.—Claw fixation forceps. The latter should be used only when a particularly firm grip of the episcleral tissue is desired, *e.g.*, in operating upon an unsteady patient under cocain.

times a day for periods of 5 to 10 minutes. If it is not resented stronger ointment, up to gr. xvi. to ℥ i., is used. As the eye becomes accustomed to one form of irritation it is well to

change the drug from time to time. Dionin, 5 to 10 per cent., may be added to the ointment, or used in solution. On application, especially for the first few times, it causes great œdema of the conjunctiva and a burning sensation. The surgeon should make the first application, as the patient is often alarmed at its severity. The stimulation of the blood and lymph flow induced by this drug is undoubtedly beneficial.

In very intractable cases the same effect may be produced by subconjunctival injections of 2 to 10 per cent. sodium chloride solution after instillation of cocain. Ten to fifteen minims of sterile solution are injected under the bulbar conjunctiva as far as possible behind the upper part of the limbus. The pain, which may be severe, is diminished by the addition of 2 per cent. cocain, or 0·1 per cent. acoin to the solution. The injection should not be repeated more than once a week.

Fig. 118.

Fig. 119.

Broad needles.

Old, degenerated, often calcareous cicatrices are best left alone, since they are liable to break down and form very dangerous ulcers, owing to the lack of resistance in the scar tissue.

If a small dense leucoma covers the pupillary area vision may be much improved by an optical iridectomy (Chap. XXII.).

Keratoplasty, the excision of a disc of scarred cornea and its replacement by a disc of clear cornea from a rabbit's or human eye, is practically never successful. The new tissue rapidly becomes opaque.

Some improvement in appearance may be obtained by tattooing dense leucomata. It is only suitable for firm smooth scars in perfectly quiet eyes, and is even then not without danger. Indian ink is tattooed into the scar with a grooved needle, under cocain, and with thorough antiseptic precautions. The eye must not be held with fixation forceps, since the punctures in the conjunctiva will be tattooed: smooth forceps must be used or none at all. More justifiable is the tattooing of small central nebulæ; it has the effect of cutting off the irregularly refracted rays, so that vision is improved (*vide* p. 190).

Treatment of Complicated Ulcers.—If perforation is imminent

special means must be adopted to prevent it. The patient should be confined to bed, and laxatives given.

It has already been pointed out that perforation improves the nutrition of the cornea. Perforation may sometimes be anticipated with advantage by *paracentesis*. By this procedure the aqueous is evacuated slowly, and the more dangerous results of perforation may be avoided. Another indication for paracentesis is extreme pain.

Paracentesis may be performed through the floor of the ulcer or just inside the periphery of the cornea. When the aqueous has escaped and the inflamed iris comes in contact with the cornea the most acute pain is felt. If, therefore, the patient is very nervous a general anæsthetic should be used, but it is to be avoided if possible on account of the straining and vomiting which may attend its administration. Whenever possible the eye is thoroughly cocained by instilling one drop of a 2 per cent. solution every five minutes for a quarter of an hour, the lids being closed during the intervals; or deep infiltration with novocain may be used (*vide* p. 629).



FIG. 120.—Keratome.

FIG. 121.—Desmarres' paracentesis needle.

FIG. 122.—Spatula.

The patient lies upon his back upon the operating table. The conjunctival sac is washed out with warm boric lotion or saline. The speculum (Fig. 115) is inserted, and the eye is fixed with fixation forceps (Figs. 116, 117) by taking up a fold of conjunctiva at a convenient spot close to the corneal margin. The points of the forceps should be pressed well into the conjunctiva so as to include the episcleral tissue, otherwise they are liable to tear the conjunctiva if much traction is exerted, as by an involuntary movement of the

patient. The incision is made with a broad needle (Figs. 118, 119), preferably one bent on the flat, or a keratome (Fig. 120), or a paracentesis needle (Fig. 121). If the floor of the ulcer is to be incised the point is inserted here so that the blade makes an angle of about 45° with the cornea. Directly it is pushed through the floor the plane of the blade is altered so that it lies against the back of the cornea; if this is not done there is imminent danger of wounding the lens. The instrument is pushed on until the incision is sufficiently long. It is then very slowly withdrawn, so that the aqueous may flow off very gradually. If the aqueous escapes suddenly the lens may be wounded, intraocular hæmorrhage may occur, or the iris may prolapse. If the last misfortune occur the prolapse must be excised (*vide* p. 201). If the operation is performed well, probably little aqueous will escape. The spatula (Fig. 122) is then applied to the edge of the wound which is nearer to the corneal margin, and this lip is gently depressed. The aqueous then escapes slowly and with a minimum disturbance to the eye. The aqueous can be evacuated on the following day by simply opening the wound with the spatula and depressing the lip.



Fig. 123. Fig. 124.

Iris forceps, straight and bent.

If the incision is made near the periphery of the cornea it should be 1 to 2 mm. inside the lower margin, especially if there is an hypopyon present. The keratome is then entered in the plane of the iris, and its direction changed as before as soon as the point is seen to be inside the anterior chamber.

In deep ulcers, such as are liable to perforate, the removal of necrotic material may be hastened by scraping the floor with a spatula, or the ulcer may be cauterised (*vide* p. 206). If the actual cautery is used it may be made deliberately to perforate the centre of the floor of the ulcer, so that the aqueous may escape and better conditions of nutrition be set

up. This procedure can only be recommended in special cases.

Another procedure is to scrape the floor of the ulcer and then to cover it with a flap of conjunctiva. The conjunctiva is separated from the limbus near the ulcer. A second incision is made with scissors parallel to the first, so that a strip of conjunctiva rather wider than the breadth of the ulcer is separated from the globe but remains attached at the two ends. This is then shifted on to the cornea so as to cover the ulcer, being retained in place by a stitch at one or both ends, thus keeping it somewhat stretched. The wound in the bulbar conjunctiva may be closed by a stitch. The lids are then carefully lifted over the strip, so as not to displace it, and the eye is bandaged.

If perforation has occurred the treatment depends upon its size and situation. If it is small or in the pupillary area prolapse of iris is not to be feared. Rest in bed, atropin, and a firmly applied bandage suffice: all forced expiration—blowing the nose, coughing, &c.—must be avoided. A sneeze can often be inhibited by firm pressure with the finger upon the middle of the upper lip close to the nose. If a small perforation is over the iris adhesion to the cornea usually occurs. It may become detached when the anterior chamber re-forms, or may be drawn out into a fine thread. No special treatment is required.



Fig. 125.—de Wecker's scissors. They should be blunt-pointed for iridectomy, one blade sharp-pointed for iridotomy.

If prolapse of iris has occurred it must be excised. No attempt should be made to replace the prolapse in these cases because the iris has become soiled with pus and replacement may result in infection of the interior of the eye and panophthalmitis.

Iridectomy of prolapsed iris is performed as follows. Instruments required: speculum, fixation forceps, two pairs of iris forceps (Figs. 123, 124), de Wecker's scissors (Fig. 125), iris repositor (Fig. 126).

If the patient is a child or highly excitable, a general anæsthetic is used, otherwise cocain suffices (*vide* p. 199). The conjunctival sac is washed out (*vide* p. 146). The speculum is inserted, and the eye fixed with fixation forceps held in the left hand. The iris repositor is passed into the wound between the iris and the cornea, with a view to freeing any adhesion. The longer the prolapse has existed the firmer will be the adhesion of the iris to the cornea. It may be impossible to free it, and in this case effectual iridectomy cannot be performed. Having freed the iris as much as possible, the fixation forceps are handed over to an assistant. The prolapse is seized with iris forceps held in the right hand, as close to the cornea as possible, and drawn well out from the wound. The second pair of iris forceps, held in the left hand, is then applied again as near the cornea as possible, and the iris drawn still further out. (Capsule forceps (Fig. 127) are very good for this purpose.) de Wecker's scissors are then taken in the right hand, and the iris is cut off close to the cornea. If the operation has been successfully performed the stump of iris retracts into the anterior chamber and is quite free from the wound. Atropin is instilled, and a pad and bandage applied.

Fig. 126.—Iris repositor, made of silver; it can be bent to any required angle.

It is to be noted here that the iris is extremely ductile; it can be dragged out much farther than might be expected, and it must be dragged out as far as possible in order that the incision may be through clean iris tissue, all the soiled part being removed. Some operators prefer to retain the fixation forceps in the left hand throughout, drawing out the iris with iris forceps in the right hand. In this case the assistant cuts off the prolapse. The method has the advantage that any sudden movement of the eye may be counteracted or followed by co-ordinated movements of the two hands of the same individual. Such sudden jerks have been known to drag the whole iris out of the wound, since it tears away at the thinnest part, viz., near the ciliary border. In ordinary cases the greatest danger is that of wounding the lens. A conjunctival flap may be used after excision of the prolapse (*vide* p. 201).

Iridectomy of prolapsed iris is only possible within the first few days, before adhesion has become firm. It is not to be



FIG. 1.



FIG. 2.

Fig. 1.—Ulcus serpens, with crescentic infiltrated advancing border above. There was hypopyon when the figure was drawn. (From a drawing by Dr. S. H. Habershon.)

Fig. 2.—Iritis, with irregular pupil and ring synechia, causing bulging forwards of the ("iris bombé").



FIG. 3.



FIG. 4.

Fig. 3.—Persistent pupillary membrane. Note the origin of the strands from the position the minor arterial circle.

Fig. 4.—Persistent pupillary membrane. The strands are adherent to the lens capsule, wh has finely granular pigment upon its surface.

[*To face p.* 203

performed after this has occurred, nor in the case of very large prolapses. In the latter there is so large an opening in the cornea that a permanent fistula may result, with loss of the eye from diminished tension and shrinking.

In very large prolapses there is much bulging and the base is often constricted. Every attempt should then be directed to obtaining a flat cicatrix. In addition to rest in bed and the means already advised, a pressure bandage must be applied for a prolonged period. A pressure bandage differs from an ordinary protective bandage only in that the space around the eye is packed carefully with cotton wool to the level of the nose and that considerable pressure is exerted in applying the bandage.

Fig. 127. — Couper's capsule forceps.

Keratocele is treated first by rest and a pressure bandage. If this fails the vesicle may be punctured, and the case treated like a perforated ulcer.

Fistula of the cornea is treated first like a perforated ulcer. If this treatment fails the edges of the fistula may be cauterised with the actual cautery, or a point of lunar caustic. In order that this may be done there must be some trace of an anterior chamber, otherwise the lens will be injured. A conjunctival flap may be drawn over the fistula.

Commencing staphyloma should be treated by a pressure bandage. If this fails a paracentesis may be done, or an iridectomy opposite the clearest part of the cornea.

Hypopyon Ulcer.—When an eye is injured so that an abrasion of the cornea is produced there is grave danger of infection from virulent pyogenic organisms. The probabilities of this occurrence depend upon the presence of such organisms and upon the amount of resistance which the tissues possess. Of all the organisms which are capable of producing deep ulceration by far the most dangerous, because the most widely spread, is the pneumococcus. It is not infrequently present in the normal conjunctival sac, but it is particularly likely to be present if there is any inflammation of the lacrymal sac (dacryocystitis). The presence of dacryocystitis is therefore a standing menace to the eye. The pneumo-

coccus, more than any other organism, tends to give rise to hypopyon, but other pyogenic organisms may also produce this result.

The substance which produces the injury may carry the infecting agent. The commonest causes are scratches with the finger nail, leaves or branches, grains of corn, and minute foreign bodies, especially stone.

Unless the organism be very virulent some lack of resistance on the part of the tissues must be predicated. Hence hypopyon ulcers are much commonest in old people and alcoholic subjects, and in the lower rather than the upper classes. The debilitating effects of hot weather are noticeable. Hypopyon ulcers also occur during or after acute infectious diseases, such as measles, scarlet fever, small-pox, and so on. In small-pox it is not a variolous pustule upon the cornea, but it differs from the typical hypopyon ulcer.

Hypopyon ulcers vary in type according to the infective agent and the age of the patient. In about 70—80 per cent. of all cases in adults the cause is the pneumococcus, and the ulcer is then of a very characteristic type, and has been called *ulcus serpens* from its tendency to travel over the cornea in a serpiginous fashion.

The typical ulcus serpens is a greyish-white or yellowish disc near the centre of the cornea (Plate VI., Fig. 1). The opacity is greater at the edges than at the centre and is particularly well marked in one special direction. A cloudy grey area, made up of fine lines, surrounds the disc, but is also more marked in the same direction. The whole of the cornea may be lustreless or hazy. There is violent iritis, and the aqueous is cloudy, or there may be a definite hypopyon. The lids are slightly œdematous, and there is conjunctival and ciliary congestion. The subjective symptoms at the early stage are pain in the eye and brow and a variable amount of photophobia.

The ulcer increases in size and depth. On the side of the densest infiltration, which often looks like a yellow crescent, the tissues break down and the ulcer spreads ; on the other side it may be undergoing simultaneous cicatrisation. In this manner it travels forwards. Meanwhile the hypopyon has become more evident, but it may vary in size from hour to hour (*vide* p. 190).

If left to pursue its natural course the hypopyon will increase and become fibrinous, the ulcer will perforate, usually forming a large opening through which the iris prolapses. The whole

cornea, except the narrow rim nourished by the corneal loops (Plate II.), may necrose, and panophthalmitis destroy the eye. In other cases an extremely dense cicatrix in which the iris is incarcerated (*adherent leucoma*) destroys sight. This may be flat or ectatic (anterior staphyloma). Sometimes the iris is bound down to the lens before perforation occurs. In such cases there are posterior synechiæ, which may be annular or total (*vide* p. 245), and the pupil may be blocked by exudates which organise into fibrous tissue (occlusion of the pupil).

Though hypopyon ulcer occurs sometimes in children, it never assumes the typical form of an ulcus serpens. In them and in some cases in adults the serpiginous character of the ulcer is not apparent, but it is distinguished by its great



Fig. 128.—Diagram of hypopyon ulcer. *a*—*b*, advancing infiltrated border; *b*—*c*, ulcerated surface; *d*, upper level of hypopyon.

tendency to extend in depth, so that perforation readily occurs. On the whole, such ulcers have a milder course than the ulcus serpens, and this is especially the case in children. It is doubtless due partly to the fact that less virulent organisms are at work, especially applicable to adult cases, and partly to greater resistance of the tissues, especially applicable to children.

The milder type of hypopyon ulcer is often due to the diplobacillus of Morax or to the allied diplobacillus liquefaciens of Petit. It usually commences as a central grey infiltration, which develops into an ulcer covered by a grey membrane and surrounded by radiating grey striæ. It generally spreads in all directions, but does not show the same tendency as the pneumococcic ulcer to spread in depth. It can only be diagnosed with certainty by bacteriological examination.

Pathology. - The ulcus serpens is due to the pneumococcus, either alone or mixed with other organisms. There is no doubt that the essential features are caused by the pneumococcus alone. A variety of organisms—staphylococci, streptococci, gonococci, &c.—have been found in atypical hypopyon ulcers. As already stated, a relatively mild form is due to the diplobacillus.

Anatomically, the ulcus serpens shows at first a depressed surface covered with slough (Fig. 128). The corneal lamellæ around and below the ulcer are separated by masses of polymorphonuclear leucocytes.

In the progressive stage the infiltration is chiefly limited to an area, wedge-shaped in section, corresponding with the yellow crescent. In other parts the edges are clean, and may be covered with epithelium. Often there is infiltration just anterior to Descemet's membrane at a spot exactly opposite the floor of the ulcer, while the intervening lamellæ are normal.



Fig. 129.—Ophthalmic galvano-cautery, which may be worked off the main with a transformer, or off a portable accumulator.

This fact accounts to some extent for the great tendency to perforation, since the inflammatory process is going on as it were from both surfaces of the cornea.

The hypopyon consists of polymorphonuclear leucocytes massed together in the lower angle. In the later stages they are enmeshed in a network of fibrin. It has been pointed out that the leucocytes are derived from the iris and ciliary processes (*vide* p. 190).

Treatment.—In all cases of hypopyon ulcer in adults treatment must be initiated at once and must be energetic. The first stage is cauterisation of the ulcer. If it is performed skilfully it does no harm and may save the eye.

Cauterisation may be performed with pure carbolic acid or with the actual cautery, the most convenient form of the latter being the galvano-cautery (Fig. 129). In my opinion the latter method, which requires more skill, possesses no advantages and has some disadvantages. Apart from the dangers attending the use of the actual cautery, carbolic acid has the advan-

tage of penetrating a little more deeply than it is actually applied, thus extending its antiseptic properties more widely : it acts both as a caustic and an antiseptic. No harm is done even if the acid spreads over the normal cornea. Although the parts touched become at once quite white, the normal tissues rapidly recover without detriment. The acid must not, however, touch the conjunctiva, otherwise very acute conjunctivitis is set up and adhesion between the lids and globe may occur.

Pure carbolic acid is applied as follows. The patient is seated or lying upon a couch. The ulcer is first stained with fluorescein (2 per cent.), in order that its limits may be more clearly defined. The conjunctival sac is thoroughly cocained (*vide* p. 199). The surgeon stands behind or at the head. With his left hand he separates the lids as in removing a foreign body, steadying the globe at the same time. The ulcer is scraped with a spatula and together with the surrounding cornea is dried with the point of a piece of blotting paper. A wooden match, somewhat pointed, is dipped into the carbolic acid. Care is taken that the wood is thoroughly wet, but has no drop of acid hanging to it which may run over the cornea. The ulcer is then touched over the whole of its surface with the point of the match. If there is sufficient carbolic acid on the match the spot touched becomes white. Special care is taken thoroughly to cauterise the advancing edge of an ulcus serpens, *i.e.*, the part marked by a yellow crescent.

Cauterisation with carbolic acid may be repeated two or three times at intervals of one or two days if the ulcer still progresses. If this treatment, combined with the use in the intervals of mild antiseptic lotions, atropin, and hot compresses, as for less severe ulcers, does not check the progress, the actual cautery may be tried. It has been recommended to heat the ulcer with the cautery without actually touching it ; or hot air may be blown on to it from a rubber ball, such as is used by dental surgeons. This treatment is certainly beneficial in some cases.

If these means fail, and especially if the tension of the eye is raised (*vide infra*), yet more drastic measures must be resorted to. Of these the most important is *paracentesis*. It may be performed as already described, but in the case of hypopyon ulcers what is called *Saemisch's section* (though it was devised by Guthrie) offers some advantages. It consists in completely dividing the ulcer from one side to the other, the ends of the incision being in healthy corneal tissue. It

must be remembered that if the operation is performed under cocain the most excruciating pain is felt when the iris comes into contact with the cornea. For this reason it is best to use a general anæsthetic, or infiltrate deeply with novocain.

Saemisch's Section. –Instruments required : speculum, fixation forceps, Graefe knife (Fig. 130), smooth iris forceps ; in case the iris should prolapse, the following should also be at hand : two pairs of iris forceps, de Wecker's scissors, iris repositor.

After cocaining or anæsthetising, the eye is washed out and the speculum is inserted. The eye is fixed with fixation forceps in the usual manner (*vide* p. 199). The point of the Graefe knife is inserted in healthy cornea just outside the edge of the ulcer, preferably at the advancing part where the yellow crescent is densest. The edge of the knife is directed forwards, so that if the lens advances before the section is complete it will touch the back of the knife and not be wounded. The knife is passed across the anterior chamber until the point is seen beyond the opposite edge of the ulcer. The counter-puncture is made in healthy cornea here, and the knife is pushed on so as to cut out. The aqueous pours out, and carries with it much of the hypopyon. Atropin is instilled, and the eye bandaged.

Fig. 130.—Narrow Graefe knife.

If the ulcer is very large it is impossible to make the puncture and counter-puncture in healthy cornea. In these cases, too, the knife usually cuts out as it is in the act of passing across the ulcer. If the hypopyon is very fibrinous it may be necessary to pull the coagulum out with the smooth iris forceps.

This procedure improves the conditions of nutrition of the cornea and evacuates the hypopyon. It is not to be advocated in mild or in the early stage of severe cases, but as a last resource it may prove beneficial.

Instead of the ordinary sublimate lotion hydrogen peroxide may be used. Optochin or ethyl hydrocuprein, a quinine derivative, is said to have a specific action on pneumococcic ulcers : a 1 per cent. solution is dropped into the eye frequently or it may be applied directly to the ulcer.

The diplobacillary hypopyon ulcer should be treated with

zinc salts. Zinc sulphate solution (0·25—1 per cent.) should be used every hour or two, and an ichthyol (1·5 per cent.)-zinc sulphate (0·5 per cent.) ointment applied to the lids.

The results of treatment of the severer forms of hypopyon ulcer are disappointing. This is largely due to the fact that they are seen too late. In rabbits it is possible to control the development of pneumococcic ulcers by intravenous injection of an anti-pneumococcic serum, if this be used sufficiently early. This method has been adopted in man, but usually fails. Another cause of failure is the development of secondary glaucoma during the treatment. The patients are usually elderly and therefore often have shallow anterior chambers. In the absence of an ulcer one would hesitate to put atropin into such an eye from the dread of causing glaucoma (*vide* p. 268). Now the presence of an ulcer, with the accompanying iritis and hypopyon, increases the risks of glaucoma from the use of atropin. We are therefore on the horns of a dilemma, for atropin will have a beneficial effect in keeping the iris at rest and tending to diminish the iritis, and therewith the hypopyon. The routine use of atropin is therefore justified, but the tension of the eye should be watched much more carefully than is usually done. If the tension rises the effect on the cornea is extremely bad, for it diminishes the lymph-flow and therewith the resistance of the tissues to bacterial toxins. It is indeed a definite indication for prompt paracentesis or Saemisch's section ; otherwise the condition will go from bad to worse.

If there is a mucocele the lacrymal sac should be excised as soon as possible (*vide* p. 598).

Keratitis disciformis is probably a milder inflammation of the same type as hypopyon ulcer, and is only rarely accompanied by a small hypopyon. It is characterised by a central grey disc, lying in the middle layers of the substantia propria. The disc is sharply defined and often shows several concentric grey lines, rather like a target. In the centre there is usually a denser "bull's eye." This form of keratitis is not common : it occurs generally in adults and is unilateral. It is accompanied by moderate irritation, which however persists for several weeks or months, leaving a permanent opacity. Owing to the central situation vision is considerably impaired. There is no ulceration, but the inflammation is probably caused by ectogenous infection through a defect in the epithelium. Keratitis disciformis has been observed as a result of vaccinia affecting the lid margin, but is not always due to this cause. It is little amenable to treatment.

Mycotic Hypopyon Ulcer.—A rare form of hypopyon ulcer due to a fungus, the aspergillus fumigatus, is occasionally met with. In it the slough is dry in appearance, and is surrounded by a yellow line of demarcation, which gradually deepens into a gutter. As the name implies, there is an hypopyon. *Treatment* is the same as for other hypopyon ulcers.

Phlyctenular Keratitis.—It has already been pointed out that phlyctens are commonly found seated upon the limbus. They may also occur within the corneal margin. The fact must be emphasised that the disease is essentially conjunctival, and when the cornea is affected it is the conjunctival element of the cornea, viz., the epithelium and the superficial layers immediately underlying it, which suffers. Phlyctenular keratitis does not necessarily result in ulceration, so that in these cases it is incorrect to classify it as a purulent keratitis, but it is convenient to consider the corneal manifestations of the disease under this heading because the complications and their treatment are similar to those of corneal ulcers in general.



Fig. 131. — Fascicular ulcer, travelling inwards towards the centre of the cornea. (Nettleship.)

Corneal phlyctens are localised infiltrations of exactly the same nature as conjunctival phlyctens. They cause more pain and reflex blepharospasm (photophobia) than do the conjunctival ones, symptoms which are worse in the morning. They may become absorbed without destruction of the overlying epithelium: in this case they cause no permanent opacity. The tendency for the epithelium to be destroyed or rubbed off is very great (cf. p. 160), and the denuded surface easily becomes infected, usually by staphylococci (*vide* p. 160). In this manner a small superficial ulcer is formed.

The corneal phlycten is a grey nodule, slightly raised above the surface. If the epithelium breaks down and an ulcer is formed the surface becomes covered with polymorphonuclear leucocytes and looks yellow. The subsequent course depends probably upon the nature of the infection and the condition of nutrition of the patient. It may deepen rapidly and even perforate, though this is comparatively uncommon.

A very characteristic form of phlyctenular ulcer is the *fascicular ulcer* (Figs. 70 B, 131). This is a serpiginous ulcer

which steadily creeps over the cornea, usually towards the centre, advancing slowly for weeks. It is supplied by a leash of vessels which lie in a shallow gutter and follow the advance of the ulcer. The ulcer starts near the limbus and heals on the peripheral side, while the central margin remains grey and infiltrated. As long as this infiltrated crescent is seen the ulcer is progressing. It always remains superficial and never perforates. When healing finally takes place the vessels gradually disappear, but the whole of the track of the ulcer remains as a permanent opacity, densest, however, where the ulcer stops.

The severest cases of phlyctenular keratitis are accompanied by a diffuse deep-lying infiltration. The large greyish area is dotted over with minute spots. The deep infiltration may disappear entirely, or it may become yellow and break down, forming a large ulcer.

Sometimes the phlyctens are so closely packed at the limbus that they become confluent and may even surround the cornea. If they break down and form a *ring ulcer* an extremely dangerous condition is set up. The nutrition of the whole cornea is endangered, and even if total necrosis does not occur an extensive perforating ulcer may be formed at the margin.

More commonly, however, a continuous infiltration of the limbus leads to the development of superficial vessels at the periphery of the cornea, a condition which is called *phlyctenular pannus* (pannus eczematosus or scrophulosus) (Plate V., Fig. 2). Unlike trachomatous pannus (*q.v.*), it shows no special predilection for the upper part of the cornea. It is thin and not very vascular, and usually undergoes complete resolution, though the course is generally very tedious. It is accompanied by intense blepharospasm.

Treatment of phlyctenular keratitis is the same as that of phlyctenular conjunctivitis (*q.v.*) until ulceration has occurred. In these cases atropin is to be combined with the yellow oxide of mercury ointment. If the latter causes much irritation, or if the ulcer is at all deep, simple atropin ointment should be substituted. The yellow ointment is particularly beneficial in fascicular ulcers, but may fail to stop their progress. The crescentic infiltrate should then be cauterised with pure carbolic acid or the actual cautery, or the vessels may be destroyed with the actual cautery. Pure carbolic does not destroy vessels efficiently, and whenever this is desired the actual cautery must be used. These extreme measures are seldom required in phlyctenular keratitis. If a fascicular ulcer has already

reached the centre of the cornea and lies over the pupillary area it is not a bad plan to allow it to progress still farther, because the opacity left by the track of vessels is less dense than that left at the final site of ulceration.

Ring ulcer must be treated by hot compresses, atropin, and a pressure bandage.

Deep ulcers and the scars left by phlyctenular keratitis must be treated on general principles (*vide* p. 194).

Phlyctenular ophthalmia is responsible for a vast amount of seriously impaired vision due to corneal opacities. The disease is a manifestation of general debility and is largely preventable. The worst effects would undoubtedly be avoided if the children could be transferred to better hygienic conditions in the country, and steps are now being taken to provide treatment in convalescent homes for such cases.

Marginal Ulcer.—Apart from the ring ulcers of phlyctenular keratitis, ulcers not infrequently occur in this situation in old people, especially in the gouty. They are shallow and little infiltrated, often multiple. They may be caused by the Morax-Axenfeld Diplobacillus. Sometimes they heal rapidly, but as rapidly recur, so that the process drags on indefinitely, to the detriment of the patient's health. Frequently the ulcers become vascularised and the vessels persist. More serious rare forms of deep marginal ulceration also occur in old people, resembling the phlyctenular ring ulcer, sometimes leading to necrosis of the whole cornea. Chronic serpiginous ulcer (*vide* p. 217) commences at the margin of the cornea. Marginal ulcers are often accompanied by severe neuralgic pains in the face and head.

Treatment.—Diplobacillary ulcers should be treated with zinc salts (*vide* p. 166). The recurrent marginal ulcers of gouty people are best treated by painting the ulcer with weak silver nitrate solution, gr. v. to ℥ i. They often heal up quickly after this application, combined with weak antiseptic lotions. Recurrence, however, can only be prevented, and that with difficulty, by constitutional treatment, which includes the special diets and drugs for gout, as well as fresh air and moderate exercise. The eyes should be protected with smoked glasses. If this treatment fails, and the ulcers are vascular, destruction of the vessels with the actual cautery is most likely to succeed. Sometimes eserin does good in these cases.

In the deep ring ulcer of old people silver nitrate or the actual cautery may be used. Paracentesis may be employed

to avert perforation and improve the nutrition of the cornea. Every attempt should be made to build up the constitution by a nutritious diet and tonics.

Central Ulcer.—Symmetrical central ulcers of an extremely indolent type are not infrequently met with in badly nourished children. They probably occur here because this part of the cornea is farthest from the nourishing vessels. They are quite superficial, show little infiltration and no vascularisation: they form shallow round pits or facets about 2 mm. in diameter. There is little or no reaction, either in the form of lacrymation or photophobia. They do not spread either superficially or in depth, nor do they show any tendency to heal. When they finally heal they often leave permanent clear facets, showing the absence of a proper supply of pabulum during the cicatricial stage. They appear to have nothing in common with phlyctenular keratitis except that both conditions are indications of defective nutrition. They are not uncommonly associated with trachoma.

They sometimes occur outside the centre of the cornea, and may perforate, allowing a knuckle of iris to prolapse. This prolapse should not be cut off, owing to the relatively large gap in the cornea, and the defective powers of repair in the debilitated patient.

Treatment must be directed especially to improving the general nutrition. A few weeks in a convalescent home in the country will effect more than any local treatment. Atropin and boric lotion are used locally. Trachoma, if present, must be suitably treated.

Keratomalacia is a rare disease in England, affecting badly nourished children, usually early in the first year of life. The conjunctiva becomes dry and shows xerotic spots (*vide* p. 183). The cornea becomes dull and insensitive; the haze increases and yellow infiltrates form. Finally the whole cornea necroses and may seem to melt away within a few hours. A characteristic feature is the absence of inflammatory reaction. In the rare cases in which the children are old enough to exhibit this symptom the disease commences with night blindness; they are able to see much better in the daytime than in the dusk. The children are usually extremely ill and very frequently die. Owing to their apathetic condition they do not close the lids, so that the cornea is continually exposed. Both eyes usually become affected. Streptococci have been found in the cornea and sometimes in the blood; in other cases the pneumococcus is present. Many of the children are syphilitic.

Experiments on animals tend to show that the disease is due to the absence of fat-soluble vitamine in the diet.

Treatment must be directed to the general health and environment. Subcutaneous injections of large quantities of normal saline solution are beneficial. The lids should be kept closed under moist warm compresses. The nutrition of the cornea is sometimes benefited by the use of eserin.

Atheromatous Ulcers occur in old dense leucomata, especially such as have undergone degenerative changes resulting in the formation of hyaline fibrous tissue and calcareous deposits. Such scars have little vitality, and the deposits act as foreign bodies. They readily succumb to infection, the epithelium being badly nourished. When ulceration once commences it proceeds rapidly and deeply, with little or no effort at repair. Perforation takes place, and is often followed by panophthalmitis.

Treatment.— The eye is frequently blind and disfiguring. In such cases it is well to excise it at once, thus relieving the patient of much unnecessary misery. If it is worth saving the condition must be treated on general principles.

Keratitis e Lagophthalmo occurs in eyes insufficiently covered by the lids. The epithelium of the exposed cornea becomes desiccated and the substantia propria hazy. Owing to the drying the epithelium is cast off, and the cornea falls a prey to infective organisms.

The condition is due to any cause which may produce lagophthalmia, *e.g.*, extreme proptosis as in exophthalmic goître or orbital tumour, paralysis of the orbicularis, and so on. The absence of reflex blinking is an important factor, as well as defective closure of the lids during sleep. Patients extremely ill from any disease are liable to this form of keratitis.

Treatment consists in keeping the cornea well covered. In mild cases it is sufficient to bandage the eyes at night. If possible the cause of the exposure must be removed. In the meantime it may be necessary to keep the lids closed with plaster and a bandage, or partially to sew them together (*vide* p. 216).

Neuroparalytic Keratitis occurs in some cases in which the fifth nerve is paralysed. It is relatively rare in nuclear and fascicular lesions within the central nervous system, unless the facial nerve is simultaneously involved. Nor does it occur in all cases of peripheral lesion ; thus, if the Gasserian ganglion is removed or the fifth nerve injected with alcohol for trigeminal neuralgia with proper precautions only a small proportion of

the cases get neuroparalytic keratitis. The disease has been known since the time of Magendie, and was attributed to special trophic impulses conducted by the nerve. This theory was combated, and the view advanced that the condition depended upon the loss of sensation in the eye. As a result, reflex blinking was more or less abolished, minute foreign bodies were not felt and therefore not removed, abrasions were unnoticed and untended, so that ulceration was readily induced and pathogenic organisms had free play. Neither of these theories satisfactorily accounts for the facts. It will be seen that the disease of the cornea usually has a very characteristic nature, quite different from the ulceration of neglected injuries. It is probable that in some manner not fully understood the disease is due to irritative changes in or about the degenerating nerve, and that mere section or paralysis of the nerve is unable to produce the disease in the absence of such irritative conditions. On the other hand, the importance of reflex closure of the lids is shown by the fact that keratitis occurs in all cases of paralysis of both fifth nerves. When only one is affected the consensual reflex from the other eye affords protection. Moreover the disease is more likely to occur if there is proptosis or paralysis of the orbicularis palpebrarum.

Besides these cases which result from radical treatment of trigeminal neuralgia, neuroparalytic keratitis is caused by intracranial tumours, gummatous basal meningitis, and fractures of the skull. (*See* Chap. XXIX.)

The characteristic feature of neuroparalytic keratitis is the desquamation of the corneal epithelium. The surface of the cornea becomes dull, and the epithelium is thrown off, first at the centre, then over the whole surface except a narrow rim at the periphery. The whole of the epithelium may peel off intact. The substantia propria then becomes cloudy and finally yellow, breaking down into a large ulcer, which is usually accompanied by hypopyon. There is no pain, owing to the anæsthesia, but ciliary injection is marked. A large perforation occurs if the case is not speedily treated. In any case the resulting leucoma is generally so large as to destroy useful vision. Relapses are the rule, the healed scar quickly breaking down again, and the whole process being repeated.

After complete removal of the Gasserian ganglion sensation is never regained.

Treatment.—The ordinary treatment of corneal ulcer should be tried as a preliminary, special care being devoted to the protection of the eye with a bandage. Improvement is often

marked, but directly the bandage is relinquished the ulceration starts anew. In these cases it is best to suture the lids together, and to keep them sutured for a long period, only separating them when the conditions of the weather—warmth, absence of high winds, and so on—are favourable. It is usually insufficient to narrow the palpebral aperture by lateral or median tarsorrhaphy, *i.e.*, suturing the lids at the canthi (Fig. 132). Too great an area of cornea is left exposed. It is best to suture them for 8—10 mm. in the middle (central tarsorrhaphy).

No anæsthetic is necessary, since sensation is lost in the



FIG. 132.—Tarsorrhaphy. A, lateral, by Fuch's method ; B, central ; C, median.



FIG. 133.—Insertion of stitches in central tarsorrhaphy.

conjunctiva and lids. The mucous membrane is dissected up from the margin of the lower lid just posterior to the lashes over a rectangular area about 8—10 mm. long in the middle of the lid. The edge of the upper lid is similarly treated at the corresponding position. Two or three sutures are then passed through the skin so that they come out at the posterior edge of the bare surface (Fig. 133), not on the posterior surface of the lid where they would rub against the cornea. After being similarly carried through the skin of the other lid they are tied, and the freshened surfaces are brought into contact. In a few days, if the stitches have not torn out, the lids will be firmly adherent.

In the worst cases the eye is useless, and tends to become a source of perpetual trouble. It is then best to excise it, which may be done without any anæsthetic if the ophthalmic branch of the trigeminal nerve is completely paralysed.

Dendritic Ulcer is the result of herpes febrilis, more rarely of herpes ophthalmicus. The vesicular stage is rarely seen, the epithelial wall of the vesicle being quickly broken. Minute shallow clear facets, like abrasions, are found in the first stage. They may be easily overlooked. They generally cause much pain, lacrymation and blepharospasm. They may spread in all directions, coalescing with others and forming a large shallow ulcer with crenated edges. More often grey striæ extend in one or more directions, increase in length, and send out lateral branches, which are generally knobbed at the ends (Fig. 134). In this manner a dendritic figure, not unlike a liverwort, is formed. The surface over the infiltrates breaks down, and an extremely irritating and chronic type of ulcer is produced. Generally only one or two of the infiltrates stains with fluorescein at any given time, but fresh spots are continually being formed. It is often associated with frontal neuralgia. Such an ulcer may persist in spite of treatment for weeks or months, sending out fresh branches but never extending in depth. The disease not infrequently recurs.

Fig. 134. — Dendritic ulcer, somewhat diagrammatic.

Dendritic ulcer is really a result of herpes, and therefore probably due to lesion of the fifth nerve or its branches. It is sometimes found in subjects of malaria.

Treatment.—The ulcer should be carefully cauterised with pure carbolic acid. Absolute alcohol has been recommended for the purpose in these cases, but causes much pain after the cocain has ceased to act. If carbolic fails to stop the progress the actual cautery should be used. Atropin, &c., and a pad and bandage are used, but in many cases it is advisable simply to use smoked glasses and get the patient out in the fresh air as much as possible. Sometimes for no known reason eserin is successful when atropin has failed to produce a good result. The general health must not be neglected, especially as the patient often becomes very depressed.

Chronic Serpiginous Ulcer (*Syn.*—*Rodent Ulcer, Mooren's Ulcer*).—This is a rare superficial marginal ulcer, usually occurring

in elderly people, and spreading, if not checked, over the whole cornea. It commences by one or more grey infiltrates. These break down, forming small ulcers, which spread and sooner or later coalesce. The ulcer undermines the epithelium and superficial lamellæ at the advancing border, forming a whitish overhanging edge, which is very characteristic. The base becomes quickly vascularised. It never perforates, but goes on with intermissions for months, until eventually a thin nebula is formed over the whole cornea, and sight is much diminished. There is sometimes iritis, and very rarely a small hypopyon. In about a quarter of the cases both corneæ are affected, but not always simultaneously. The cause is unknown.

Treatment. The overhanging edge should be cut off with



Fig. 135.—Distribution of the branches of the ophthalmic division of the fifth nerve on the face.

scissors, and then the whole surface of the ulcer, and especially the margin, should be well cauterised with the actual cautery. A few cases have responded to repeated applications of absolute alcohol to the ulcers. More commonly treatment fails to stop the process, which has even been known to recur in the cicatrised cornea.

Non-suppurative Keratitis.—This occurs in two types, superficial and deep. The superficial forms include some purely symptomatic conditions, such as pannus. Some of them, such as herpes, may lead to the formation of ulcers and pass into the purulent type. The deep forms never suppurate.

Herpes Corneæ.—Herpes of the cornea occurs most commonly in herpes febrilis, more rarely in herpes ophthalmicus.

In *herpes febrilis* vesicles form upon the cornea quite similar in nature to those which may be present, but are not always so, upon the lips or angle of the nose. The patients are usually suffering from some febrile condition, *e.g.*, influenza, pneumonia, whooping cough, &c., but it may be so trivial as to escape observation. It sometimes occurs after typhoid inoculations. The vesicles are scarcely the size of a pin's head and are often arranged in rows or groups. They quickly rupture, forming abrasions which heal rapidly, leaving no opacity.

Fig. 136.—Herpes ophthalmicus.

Usually, however, fresh crops of vesicles appear, and the condition may prove very obstinate. In severe cases, ulcers are formed which may be of dendritic type (*vide* p. 217), or may be due to secondary infection. The acute stage is accompanied by great irritation, lacrymation, and blepharospasm. The ocular affection is usually unilateral and on the same side of the face as vesicles on the lips, &c. The cornea is generally not anæsthetic except at the spots attacked.

Herpes corneæ may be mistaken for phlyctenular keratitis. The former occurs usually in adults, the latter in children. The clear vesicles differ in appearance from the grey infiltra-

tions of phlyctenular keratitis. After the vesicles have burst, the shape and the total absence of vascularisation are distinguishing features. They are then liable to be mistaken for traumatic abrasions, from which they are distinguished by their grouping, the crenated edges when several have coalesced, persisting shreds of the ruptured vesicles, and the absence of history of any injury.

In *herpes ophthalmicus* one or more of the branches of the ophthalmic division of the fifth nerve (Fig. 135) is marked out by rows of vesicles or the scars left by them, exactly as in herpes zoster in other parts of the body. The supra-orbital, supra- and infra-trochlear branches are nearly always involved; frequently the nasal branch; only rarely the infra-orbital branch. It is very rarely bilateral. There may be fever and malaise at the onset, and the eruption is preceded by severe neuralgic pains along the course of the nerves. These are so characteristic that they should arouse suspicion of the nature of the disease before the vesicles appear. The pain sometimes ceases after the outbreak of the eruption, but not always, and it may continue for weeks or months. The skin of the lids and other areas affected becomes very red and œdematous, so that the disease is often mistaken for erysipelas. The characteristic distribution and especially the strict limitation to one side of the middle line of the head should obviate this error. The vesicles often suppurate, bleed and cause permanent scars. The active eruptive stage lasts about three weeks and is followed by some anæsthesia of the skin. Ocular complications arise during the subsidence of the eruption.

Corneal vesicles are rare, but when they occur are exactly like those of the febrile form, and behave in the same manner. More commonly there is a diffuse deep infiltration of the cornea (*keratitis profunda*), associated with iridocyclitis (*q.v.*). The cornea is usually insensitive. This is tested by touching it with a wisp of cotton wool, and comparing with the opposite eye. The slightest touch is followed by reflex closure of the lids if the cornea is sensitive. The intra-ocular tension is not infrequently somewhat diminished in the early stage. The eye lesions are very obdurate and often persist long after the disease has otherwise passed away. In some cases there is associated paralysis of motor cranial nerves, especially the third, sixth and seventh. Facial palsy adds seriously to the risk of the eye, owing to its partial exposure.

Herpes ophthalmicus occurs at any age, but generally in elderly people. Not infrequently there is a history of contact

with patients suffering from chicken-pox, and evidence is accumulating that the relationship is more than a mere coincidence. It may account for the fact that herpes zoster is apt to occur in epidemics, which are more common in the spring and autumn than at other times.

The disease is due to lesions in the Gasserian ganglion of the same type as those found in the posterior root ganglia in herpes zoster, *i.e.*, microscopic thromboses and hæmorrhages similar to those found in the anterior cornual nuclei in anterior poliomyelitis. This disease shows many analogies on the motor side to herpes zoster. Probably both are caused by infection, the lesions being due either to bacterial invasion or more probably toxins. Herpes ophthalmicus has been known to follow involvement of the Gasserian ganglion in a malignant growth.

Treatment.—In herpes corneæ the eyes should be protected with a bandage. When the vesicles have ruptured, atropin and warm compresses give most relief. It is sometimes necessary to use cocain when the pain is acute, though it has a bad effect upon the epithelium. Oily applications—ol. ricini –are sometimes grateful. Prolonged ulceration must be treated according to the type of ulcer, whether dendritic (*vide* p. 217) or septic. The general health must be attended to ; quinine, valerian, salicylic preparations, *e.g.*, salicylates, aspirin, &c., are useful.

In herpes ophthalmicus the skin may be treated with cocain ointment (1 per cent.) and dusted with starch powder. Quinine is generally given internally and aspirin to relieve the pain. The patient should be kept in bed during the acute stage.

Superficial Punctate Keratitis is an uncommon condition allied to herpes febrilis, and, like it, usually associated with influenza or catarrh of the respiratory tract ; but no vesicles are formed. It commences as an acute conjunctivitis. At this stage, or after it has passed off, groups or rows of superficial, slightly raised, grey dots are seen scattered over the cornea, least at the periphery. They may or may not stain faintly with fluorescein. They remain unchanged for months, but gradually disappear. The disease affects one or both eyes, usually in young people.

Treatment as for herpes.

In *acne rosacea*, generally in elderly women, keratitis associated with much irritability and lacrymation may occur. In addition to slight muco-purulent conjunctivitis (*vide* p. 146), greyish white

infiltrates and small ulcers appear in the cornea. They are very intractable and frequently recur.

Interstitial Keratitis (*Syn.—Parenchymatous Keratitis*) is a deep keratitis usually affecting children between the ages of five and fifteen, the subjects of congenital syphilis. It is commoner in girls than boys.

After slight irritative symptoms, with some ciliary congestion, one or more hazy patches appear in the cornea, near the margin or towards the centre (Plate VII., Fig. 1). If they are near the margin they push forwards towards the centre; if at the centre, others appear and fuse, until finally the whole cornea looks lustreless and dull. Minute examination shows that the patches are in the deep layers and are made up of denser spots and fine streaks. In two to four weeks the whole cornea is hazy with a steamy surface, giving a general appearance like that of ground glass. Denser spots can always be seen in the general mist. In the severest cases the whole cornea becomes quite opaque, so that the iris is hidden; as a rule the iris can be seen dimly.

Meanwhile vascularisation has occurred. It is of the deep type (*vide* p. 86), consisting of radial bundles of brush-like vessels. The larger marginal plaques may be very vascular; as they are covered by a layer of hazy cornea their bright scarlet is toned down to a dull reddish pink ("salmon patches"). The separate vessels can only be seen by magnification. The small salmon patches are often crescentic; when larger they are sector-shaped. The opacity extends a little beyond the vessels, which seem to push the opacity in front of them. In the acme of the condition the vessels run in radial bundles almost, but seldom quite, to the centre of the cornea. Vascularisation is probably brought about for the supply of protective substances which cannot reach the cornea by the normal process of diffusion. There is often a moderate degree of superficial vascularisation, greater in some cases than in others, but never extending far over the cornea. The conjunctiva may be heaped up like an epaulette at the limbus, so that some slight resemblance to phlyctenular keratitis may be seen. Indeed, it is probable that these patients are both syphilitic and "strumous."

After the disease has reached its acme it commences to subside. The cornea clears from the margin towards the centre, which may long remain hazy, though it too finally clears, except in the worst cases. As the cloudiness disappears the

PLATE VII. (*To face p.* 222.)

Fig. 1.—Interstitial keratitis, showing general infiltration of the cornea with vascularisation at the periphery, especially above ("salmon patch").

Fig. 2.—Episcleritis.

FIG. 1.

FIG. 2.

vessels become obliterated, but though they cease to carry blood they remain permanently as fine opaque lines; they can be demonstrated only by magnification either with oblique illumination or more definitely by the direct method (*vide* p. 116). The cornea should be examined in the dark room by oblique illumination with a corneal loupe, when the vessels will be seen as grey lines on a black background. It should then be examined by the direct method with a + 20 D lens behind the mirror. The edge of the pupil is brought into focus, and then either the head is withdrawn slightly or plus lenses are added, the head being kept still, until the cornea is in focus. The vessels will then be seen as black lines against a red background. The characteristic radial course and distribution (*vide* p. 86, Fig. 71) affords permanent proof of previous occurrence of interstitial keratitis, and is important evidence of congenital syphilis.

At the height of the disease vision may be reduced to finger-counting or hand movements, and there is much lacrymation and reflex blepharospasm. Very mild cases are met with in which the patches of cloudiness are thin, scarcely vascularised at all, and clear up quickly. In the marginal type the opacity may be limited to a sector.

The surface very rarely becomes ulcerated. It is frequently stippled, steamy, and slightly uneven, and this condition may persist. In the worst cases the cornea may be enormously thickened and gelatinous in appearance. The impression given is that the cornea is very ectatic, and that the eye is in a hopeless condition; it will probably clear up well with useful vision. In no case must an eye be removed on account of interstitial keratitis.

Interstitial keratitis is almost invariably symmetrical, though an interval of three or more weeks usually intervenes before the onset in the second eye. Rarely the interval is several months or even years. The acute stage lasts at least six weeks and may extend to several months. The clearing of the cornea takes many weeks or months, but not much improvement can be expected after eighteen months. Delayed interstitial keratitis, *i.e.*, in patients over thirty, occasionally occurs and is more liable to be unilateral; it is often very severe. Interstitial keratitis in acquired syphilis is uncommon, though probably not so rare as is generally thought; it is also generally unilateral: several cases have been reported in which the primary lesion has been on the lids or face, and the keratitis has been limited to the same side. Most of the

reported cases have occurred about two years after the primary sore, but the interval may be much shorter or longer.

There is evidence of relatively high mortality among the children of mothers who have had interstitial keratitis.

In some cases interstitial keratitis follows an injury to the eye, such as a blow or operation, *e.g.*, tenotomy. I have seen it follow an attack of phlyctenular conjunctivitis. Apparently the injury acts as an exciting cause in a subject naturally prone to the disease, usually a congenital syphilitic.

Pathology.—The rare cases of interstitial keratitis which have been examined anatomically show that the infiltration of

FIG. 137.—Section of interstitial keratitis, showing infiltration confined to the deeper layers of the substantia propria (× 60). *E*, epithelium; *B.m.*, Bowman's membrane; *S.p.*, substantia propria; *D.m.*, Descemet's membrane with endothelium.

the cornea is almost entirely limited to the deeper layers lying immediately anterior to Descemet's membrane (Fig. 137).

It has been pointed out that corneal conditions which are secondary to conjunctivitis are superficial. Similarly deep keratitis is secondary to disease of the uveal tract. Thus, in tubercle of the iris (*q.v.*) it is not uncommon in the late stages to meet with an interstitial keratitis fundamentally the same as in congenital syphilis. Further, anatomical investigation has shown that in the latter form the uveal tract is profoundly affected. Thus, it is the rule for a considerable degree of iritis to be present. Sometimes there is severe cyclitis, as shown by the presence of precipitates ("keratitis punctata," *vide* p. 257) on the back of the cornea. Not infrequently there is choroiditis. If the pupil of the unaffected eye

is dilated and the periphery of the fundus carefully examined, patches of anterior choroiditis will not uncommonly be found. All these facts support the view that the disease is fundamentally a uveitis, and that the keratitis is secondary, *i.e.*, merely symptomatic. Clinically it masks the uveitis, which is thus liable to be overlooked, and hence the disease is called a keratitis. It is very important to realise the true pathology, since treatment must be directed to avoiding the deleterious results of iridocyclitis rather than those of keratitis.

It is probable that anaphylaxis plays some part in the pathogenesis of interstitial keratitis (Wessely, von Szili), and may

Fig. 138.

Fig. 139.

Fig. 140.

Fig. 141.

Fig. 142.

Fig. 143.

Figs. 138 to 143.—Hutchinson's teeth. (Nettleship.)

account for the occurrence, after a latent period, of the disease in the second eye.

Ætiology.—It has already been stated that nearly all cases which are not obviously secondary to tubercle of the iris, &c. —and these are rare—are due to congenital syphilis. Many surgeons, however, consider that interstitial keratitis is not infrequently due to tubercle. Owing to the importance of diagnosing congenital syphilis in doubtful cases the principal signs of this complaint may be enumerated.

(1) General features. Prominence of frontal eminences, flatness of the bridge of the nose, breadth of face, stupidity or undue precocity, deafness.

(2) Hutchinson's teeth (Figs. 138—143). There is nothing characteristic in the first dentition. In the permanent teeth only two, the upper central incisors, are to be relied upon,

but the other incisors and first molars are often deformed. The characteristic change in the central upper incisors appears to depend upon defective formation of the central lobe (Figs. 139, 142—143). Soon after eruption this lobe wears away, leaving a vertical notch at the centre of the cutting edge (Fig. 138). If the cause has acted so intensely as to prevent the development of the central lobe, there is, instead of a notch, a narrowing and thinning of the cutting edge as compared with the crown, and this according to its degree produces a resemblance to a screwdriver or peg (Figs. 140, 141). The teeth are too small in every dimension, so that the incisors are separated from one another by considerable spaces. In extreme cases all the incisors are peggy and much dwarfed.



Fig. 144.—Arcus senilis.

The changes are usually symmetrical, but Fig. 142 shows one tooth typically deformed and the other normal.

(3) Linear cicatrices, relics of former rhagades, are met with at the angles of the mouth. Cicatrices on the hard or soft palate or elsewhere in the mouth point to antecedent syphilitic ulcerations.

(4) Shotty, painless lymphatic glands are felt in the neck, especially in the posterior triangle.

(5) Hard periosteal nodules, best found upon the shins, and chronic synovitis of the joints, especially the knee joints, may be present.

Evidence of acquired syphilis may be sought in the parents. In cases of doubt, the Wassermann test should be applied.

As already mentioned, interstitial keratitis also occurs in

acquired syphilis, and it has been attributed to malaria, myxœdema, trypanosomiasis, &c. It frequently occurs in foxhounds and other highly-bred dogs, and I have seen it in dogs after the thyroid gland has been removed.

Treatment.—It is usual to order antisyphilitic remedies, but it is doubtful if they have any influence over the course of the disease, partly because the cornea is non-vascular. Mercury inunctions, or powders or pills of mercury combined with chalk and ipecacuanha, or perchloride of mercury may be given. Iodide is best administered in children in the form of syrup of iodide of iron, and may be combined with syrup of phosphates as a tonic. Salvarsan and its substitutes have proved disappointing. The administration of thyroid gland has proved serviceable in some cases. Cod liver oil or maltine may be used when there is a tuberculous element about the condition, and in all cases general hygienic régime must be instituted.

Local treatment consists in guarding against the evil effect of the uveitis which is an invariable accompaniment of the disease. Atropin is ordered as a routine measure, with the double purpose of keeping the ciliary body and iris at rest and preventing the formation of posterior synechiæ. There is often great difficulty in getting the pupil to dilate, probably owing to defective penetration of the drug through the diseased and vascularised cornea. Hot bathings or radiant heat should be used frequently in the acute stage. In obstinate cases with much lacrymation and blepharospasm, especially if the pupils will not dilate with atropin, leeches applied to the temple do good (*vide* p. 253). Smoked glasses are ordered.

In later stages the means used for clearing corneal cicatrices (*vide* p. 196), depending essentially upon improving the lymph flow through the cornea, are also indicated for the opacities of interstitial keratitis. Prolonged use of yellow oxide of mercury ointment, combined with atropin in the earlier stages, is commonly made : it should be well massaged into the eye.

Other forms of deep keratitis, *e.g.*, the central deep keratitis of adults, senile marginal deep keratitis, sclerosing keratitis (*vide* p. 238) occur.

Keratitis profunda is the name given to a central deep infiltration of the cornea of indefinite origin. Such a condition occurs after contusion of the eye, and in this case usually clears up rapidly. In other cases a deep grey opacity, seen on magnification with the loupe to be composed of dots and striæ, occurs in

the centre of the cornea in adults. There is little irritation, and very little, if any, vascularisation. It begins to clear up after four to eight weeks, but may leave permanent diffuse opacities. It has been attributed to cold, malaria, and other causes. It should be treated with the same local remedies as for interstitial keratitis.

Keratitis marginalis profunda occurs rarely in old people, commencing as a greyish yellow infiltration, one or two millimetres broad, continuous with the sclera. It is usually limited to the upper part of the cornea, but occasionally forms a complete ring. The slight irritation subsides in a week or two, leaving a permanent opacity, resembling arcus senilis, but unlike it, continuous with the sclerotic.

Degenerative Changes in the Cornea.

Arcus senilis is a fatty degeneration of the cornea met with in old people (Fig. 144). It commences as a crescentic grey line concentric with the upper and lower margin of the cornea. The extremities of the crescents finally meet, and an opaque line, thicker above and below, is formed completely round the cornea. It is characterised by being separated from the margin by a narrow line of comparatively clear cornea. It is sharply defined on the peripheral side, fading off on the central. It is never more than about 1 mm. broad, and is of no importance, either from the point of view of vision or of the vitality of the cornea.



Fig. 145.—Band-shaped opacity of the cornea, from an eye with iridocyclitis.

Arcus juvenilis is exactly similar to arcus senilis, but is a rare condition found in children. Even arcus senilis may develop at a comparatively early age, but the juvenile condition is probably congenital. It is of no importance. The characteristic diagnostic feature of both these opacities is the presence of a line of clear cornea between them and the limbus. This is occasionally found in old sclerosing keratitis, but in this case the opacity is usually localised to some one part of the cornea and extends farther towards the centre.

Band-shaped Opacity (*Syn.—Transverse Calcareous Film*

Zonular Opacity).—This is a common condition in old, blind, shrunken eyes. It is due to defective nutrition and exposure. It lies entirely in the interpalpebral area, commencing at the inner and outer sides, and progressing until it forms a continuous band across the cornea (Fig. 145). Near the corneal loops, just inside the limbus, the cornea is generally relatively clear, as in so many degenerative conditions, probably owing to the better nutrition close to the blood-vessels. The condition is due to hyaline degeneration of the superficial parts of the substantia propria, followed by the deposition of calcareous salts.

As a rare condition it is found in otherwise healthy eyes, sometimes as a horizontally oval area in the palpebral fissure, usually in both eyes.

Treatment. In the rare form last mentioned, improvement of vision may be obtained by scraping off the opacity, which is usually calcareous and quite superficial. In the common form the eye is blind, and nothing remains but to remove it if it is painful or unsightly.

Other Degenerative Changes are frequently met with in old leucomata, anterior staphylomata, and so on. They consist of hyaline degeneration, calcification, &c. Such scars are liable to a serious form of ulceration (*vide* p. 214).

Rare degenerative conditions are *nodular* and *reticular* (grill or lattice-like) *opacities*. They occur as a familial disease, usually picking out the young males of a family and commencing at about the age of ten or twelve. Opaque dots or forked lines appear under the epithelium in the centre of the cornea and slowly increase in number, but never invade the ring of cornea close to the margin. There is little inflammatory reaction and the cause is unknown. Vision is gradually obscured and treatment is of little avail. I have seen grey moss-like opacities, apparently of similar nature, with the same distribution in the cornea of elderly people.

Senile marginal atrophy, in which a gutter forms in the periphery of the cornea in the situation of an arcus senilis, occurs rarely in one or both eyes of old people. The gutter may become ectatic.

Ectatic Conditions.

It has already been stated that ectatic conditions of the cornea may result from inflammation, viz., keratectasia (*vide* p. 191), and anterior staphyloma (*vide* p. 193). Two forms of ectasia of non-inflammatory origin are known, viz., keratoconus and keratoglobus.

Anterior Staphyloma is a protuberant cicatrix arising from a prolapse of the iris; it may be partial or total (Fig. 114). It may follow a perforating wound, but is usually due to perforation of an ulcer, especially such as is caused by ophthalmia neonatorum. The primary protrusion occurs at the moment of prolapse. Cicatrisation follows, and in the case of small prolapse may lead to flattening of the scar. In other cases of small, and in all cases of large prolapse the contraction of the scar tissue is insufficient to bring this about, and the soft cicatrix yields to the normal intraocular pressure. Generally the prolapse of the iris leads to blocking of the angle of the anterior chamber, the intraocular tension rises, and the cicatrix yields still more, or if it was previously flat, secondary protrusion takes place.

Partial staphylomata are usually conical, rarely hemispherical; they usually extend to the margin on one side. Total staphylomata are usually hemispherical, rarely conical. There is invariably a rim of cornea around the pseudo-cornea, this rim being well nourished by the peripheral blood vessels and never necrosing through ulceration. The thickness of the staphyloma varies very greatly in different cases, and often in different parts of the same staphyloma. In the latter bands of cicatricial tissue develop, while the intermediate parts project; in this manner a racemose staphyloma is produced. Owing to the rise of intraocular tension the whole eyeball expands, especially in children, in whom the walls are relatively plastic. If the lens has not been expelled when the ulcer perforated, as sometimes happens, it is flattened, the expansion of the ciliary ring causing stretching of the suspensory ligament. Owing to contact of the lens with the inflamed cornea after perforation the anterior capsular cells of the lens often proliferate and form an anterior capsular cataract (*q.v.*). The high intraocular pressure also causes cupping of the optic disc.

The pseudo-cornea is formed by organisation of the exudates on the surface of the prolapsed iris. It consists of fibrous tissue covered by epithelium, and lined by rarefied iris pigment epithelium. The epithelium on the anterior surface is often very thick and epidermoid; the fibrous tissue often undergoes degeneration. The anterior chamber is obliterated in total staphyloma, while the posterior chamber is enormously enlarged, and filled with yellow albuminous fluid.

The sight is always diminished, and in total staphyloma is reduced to perception of light or total blindness. The eye may project between the lids, so that a xerotic condition is

set up and atheromatous ulcers may form. Ectropion of the lids may occur mechanically. The tension is raised, either as the cause of the protrusion, or as a result of the blockage of the angle of the anterior chamber (Chap. XIV.). This may lead to pain. The staphyloma may be so thin that rupture occurs on the least injury, and may be several times repeated.

Treatment.—Total anterior staphyloma is best treated by excision of the eye, with or without the insertion of a glass globe in Tenon's capsule. Patients will often prefer to keep the eye if it is painless and not too disfiguring.

Various methods of ablation, in which the anterior part of the eye is removed and the contents scooped out, have been devised as an alternative for excision. They give a movable stump on which to set an artificial eye, but they are open to the objection that they are not entirely free from the danger of causing sympathetic ophthalmia (*q.v.*).

Treatment of partial staphyloma is directed to obtaining flattening of the cicatrix, preventing or relieving increased intraocular tension, and improving sight. The attempts which should be made at the early stage have already been described (*vide* p. 203). Iridectomy is sometimes advisable with a view to improving vision and preventing or curing glaucoma (*q.v.*). It cannot be performed at the site of the staphyloma, since there is no anterior chamber here, but must be done over the clearest part of the cornea.

Keratectasia differs from anterior staphyloma in that the iris takes no part in it. Its causation and prevention have already been discussed (*vide* p. 191). Sometimes the whole cornea expands, producing a condition which is almost identical with the keratoglobus (*vide* p. 232) of infantile glaucoma, though due to a different cause. When it is fully developed treatment is useless.

Keratoconus (*Syn.—Conical Cornea*) is probably due to a congenital weakness of the cornea, though often it only manifests itself after puberty. The cornea is thin and weak near the centre, and gradually bulges forwards more and more; the apex is always slightly below the centre. Sometimes it pulsates synchronously with the arterial pulse, and this may cause a subjective apparent pulsation of the objects looked at. The pulsation may be demonstrated with Schiötz' tonometer. The cornea is at first perfectly transparent, and vision is impaired through the protrusion and alteration in curvature. If advanced the conical shape is easily recognised in profile. In the less advanced cases distortion and diminution in the

size of the corneal reflex over the centre is the chief guide. These changes are best seen when the reflex from a large flat disc painted with broad concentric black and white lines (Placido's disc or keratoscope) is observed through a hole in the centre of the disc, or when the cornea is examined with the ophthalmometer. With the ophthalmoscope mirror at a distance of 1 metre a ring of shadow concentric with the margin is seen on the red reflex, altering its position on moving the mirror. It is due to a zone through which few rays pass into the observer's eye owing to the emergent rays on the central side being convergent whilst those on the peripheral side are divergent.

The patient becomes myopic, but the error of refraction cannot be satisfactorily corrected with glasses owing to the hyperbolic nature of the curvature. The condition is almost invariably bilateral, though frequently more advanced on one side than the other. It may be extremely slight and very slowly progressive, or the reverse. In the later stages the apex may become opaque, but ulceration, rupture, increase of tension, and so on, do not occur.

Treatment.—In the early stages every effort should be made to improve vision with glasses and the progress should be carefully watched. Various methods have been adopted to stop the process. Miotics are probably useless. The best treatment is cauterisation of the apex with the actual cautery. The cauterisation at the extreme apex must be deep, and perforation here has been advocated. The latter procedure is not without danger to the eye from the formation of a corneal fistula, anterior synechia, infection, &c. Some operators remove a small wedge-shaped slice from the thickness of the cornea at the apex. It is more difficult to perform and does not give better results on the whole than the cautery. The scarring from cauterisation is much less than might be anticipated, but it may be advisable to do an optical iridectomy if vision is still very bad. Anterior sclerectomy (trephining) has been performed for conical cornea, but without advantage.

Keratoglobus is the hemispherical protrusion of the whole cornea which is met with in infantile glaucoma (*q.v.*).

Symptomatic Conditions.

There are many pathological conditions of the cornea which are merely evidence of disease in other parts of the eye or of extension of disease. Some are often described as true

diseases of the cornea, notably as forms of "keratitis." This involves a wrong principle and a misuse of terms which can only lead to confusion. Since it is of great importance to distinguish these conditions from primary affections of the cornea, both from the points of view of diagnosis and treatment, it will be well to review the more common here.

In **glaucoma** there is nearly always uniform diffuse bluish haze of the whole cornea. It is due to alterations in the refractive conditions of the corneal elements, brought about by the increased intraocular pressure, and not to any gross pathological change. If the condition persists the cornea becomes



FIG. 146.—Œdema of the cornea and vesicular "keratitis," from a glaucomatous eye (× 60).

hazy throughout, and this haziness, unlike the former, does not immediately disappear when the intraocular tension is relieved. Here there is definite œdema, due to the impediment which prolonged tension causes to the diffusion of lymph. It manifests itself first in the epithelium, which becomes steamy and stippled. Sections show that this is due to accumulation of fluid in and between the cells, especially the basal cells (Fig. 146). Fluid also accumulates between the lamellæ and around the nerve fibres. If the œdema lasts for a very long period, as in eyes blinded with absolute glaucoma, the epithelium may be raised into vesicles or bullæ (*vesicular or bullous "keratitis"*). The evidence of prolonged tension

makes it easy as a rule to distinguish this condition from herpetic vesicular keratitis; moreover the vesicles or bullæ have firmer epithelial walls and show less tendency to burst.

Nearly allied to bullous "keratitis" is the formation of epithelial threads, which adhere to the cornea by one end, while the other, which is often club-shaped, hangs down free. This is commonly called *filamentary "keratitis."* It occurs in the same conditions, *i.e.*, usually associated with glaucoma, but it may be seen rarely without any apparent cause in cases of the herpetic type.

Keratic precipitates, usually badly termed "keratitis punctata" or "k.p." in England, are often deposited upon the back of the cornea in cyclitis and iridocyclitis. The greatest care must be taken not to overlook them, since they may be almost the only objective sign of serious disease. They may be on the back of a clear cornea or the deeper layers may be infiltrated as a result of the iridocyclitis; thus, they are not uncommon in interstitial keratitis. Their appearance and nature will be described in discussing their cause (*vide* p. 257).

Fig. 147. — Striate opacity of the cornea, following section of the cornea above for extraction of cataract.

Opacities of the cornea are often secondary or symptomatic. Such are the grey or white, usually tongue-shaped marginal opacities which follow scleritis. Owing to their resemblance to the sclerotic they are called *sclerosing "keratitis"* (*vide* p. 238).

Congenital opacities of various kinds are sometimes met with. Many are not truly congenital, but are due to injury received at birth. Birth injury of the cornea takes the form of temporary diffuse opacity due to œdema, or of permanent linear opacities due to ruptures of Descemet's membrane.

Striate opacity occurs in various forms. The commonest form is that seen after operations upon the globe in which a peripheral section has been made, as in cataract extraction. Here, when the section is above, delicate grey lines run down vertically from the wound and may pass completely across the cornea (Fig. 147). They can be seen clearly only by magnification. They disappear spontaneously as the wound heals. They are due to slight folding of the cornea, whereby Descemet's membrane and the adjacent lamellæ become wrinkled. Radial striæ are seen around wounds or ulcers;

they are partly referable to the same cause, partly to distension of the interlamellar spaces by œdema. The fine hatching which is seen around ulcers and sometimes after tight bandaging are to be referred to similar causes.

Opacities of the cornea may be due to improper treatment. One of the commonest is due to *deposits of lead salts* upon an abrasion or ulcer. An insoluble, densely opaque white film is precipitated and adheres very firmly. The spot is sharply defined and looks like white paint. Probably the deposit is always thrown off eventually, but a very long time may elapse. An attempt should be made to scrape it off, but it is wisest never to use lead lotions at all in the treatment of any affections of the eye. They can always be replaced by equally efficient substitutes.

Pigmentation of the cornea may also occur from improper treatment. Prolonged use of silver nitrate as a paint or in the form of drops is followed by dark brown staining of the conjunctiva and slight staining of the cornea. The condition is called *argyrosis* (*vide* p. 184), and in the conjunctiva is found to be due to impregnation of the elastic fibres with metallic silver ; in the cornea Descemet's membrane is stained. It is permanent ; hence silver solutions, including protargol, argyrol, &c., should never be ordered as drops or for use at home, at any rate without stringent injunctions.

Blood in the cornea is rare. It may occur as a bright red spot quite superficially at the margin or as a greenish or rusty stain over the whole surface. In the latter case it is derived from blood in the anterior chamber, associated with high tension –a relatively infrequent combination (*vide* p. 398).

Tumours of the cornea, so-called, are probably always secondary extensions, most commonly from the conjunctiva, the limbus being a favourite situation for these growths (*vide* p. 184).

CHAPTER XII.

DISEASES OF THE SCLEROTIC.

Redness of the white of the eye is caused by a variety of conditions. The commonest is conjunctivitis: the next common some inflammation of the anterior part of the uveal tract, viz., the iris and ciliary body. Some of the distinctive characteristics of the redness in conjunctival and ciliary inflammation have already been pointed out (*vide* p. 77). Redness of the white of the eye may also be caused by inflammation of the sclerotic, and it is a frequent error among beginners to ascribe the other forms to this disease. It must therefore be borne in mind that episcleritis and scleritis are relatively uncommon.

Inflammation of the Sclerotic.

Two forms of inflammation of the sclerotic are described, superficial or episcleritis, and deep or scleritis. They might equally well be considered mild and severe forms of the same disease, but the distinction is convenient clinically since they usually differ in the course they take.

Episcleritis is an inflammatory affection of the deep subconjunctival connective tissues, including the superficial scleral lamellæ. A circumscribed nodule, which may be as large as a lentil, appears usually two or three millimetres from the limbus (Plate VII., Fig. 2). It is hard, immovable, and very tender, the conjunctiva moving freely over it. It is traversed by the deeper episcleral vessels, and therefore looks violet, not bright red. It is extremely chronic, never ulcerates, and may be entirely absorbed, but more frequently leaves a slate-coloured scar behind, to which the conjunctiva is adherent. The cornea and uveal tract rarely participate in the inflammation.

There may be little or no pain, but usually there is a feeling of discomfort and tenderness on pressure, and severe "neuralgia" is often complained of. The nodule becomes gradually

absorbed in the course of days, or, more frequently, weeks, but during the process of absorption, or soon after, fresh nodules of the same type arise. In this manner the disease may drag on for months. Both eyes are frequently affected. In the worst cases the disease extends into the deeper parts of the sclerotic, and thus passes almost imperceptibly into scleritis.

Anatomically dense lymphocytic infiltration of the subconjunctival and episcleral tissues is found.

Rheumatism and gout are commonly indicated as the chief causes of episcleritis. A history of acute rheumatism is rarely obtained; more commonly there has been well marked "muscular rheumatism," which is to be ascribed to septic absorption (*vide* p. 258). Tuberculous or syphilitic patients seem predisposed, and almost invariably there is some cause of general debility. It is commonest in elderly people (cf. Scleritis), and in women.

Treatment.—General treatment is of more avail than local. Even in cases in which no history of rheumatism can be elicited, salicylic preparations—salicylates, salicin, aspirin—seem to do good, and should be tried. If they fail resort should be had to iodides. Diaphoresis by pilocarpin injections and vapour baths does good in some cases, and inunctions of mercury are useful apart from syphilitic infection.

The most useful local treatment is massage by the finger applied to the upper lid. A simple boric ointment may be used or a weak yellow oxide of mercury ointment, but strong stimulants can rarely be borne. In the more severe cases warm compresses, dionin, and leeches to the temple should be employed. In the worst cases the constant current, scarification or superficial cauterisation are indicated. Every effort must be made to build up the constitution, and success often depends upon these measures.

Scleritis is rarer than episcleritis. There are usually nodules, or a single nodule, but the area affected is much less circumscribed. The swelling is at first dark red or bluish, later it becomes pale violet and semitransparent, like porcelain. It may extend entirely round the cornea, forming a very serious condition known as *annular scleritis*. Scleritis differs from episcleritis in that the cornea and uveal tract are involved, some iritis, but more cyclitis and anterior choroiditis being present. There is no ulceration, but much absorption, so that the sclerotic is thinned, a dark purple cicatrix being formed, which is often too weak to withstand the intraocular

pressure, so that ectasia follows (*ciliary staphyloma*). In many cases of diffuse deep scleritis hard whitish nodules develop in the inflamed zone. They are the size of a pin's head and lie beneath the conjunctiva, all at about the same distance from the corneal margin; they disappear without disintegrating.

Anatomically scleritis is the same as episcleritis, but extends deeper; there is dense lymphocytic infiltration of the sclera, the lamellæ being separated by cords of cells.

Both eyes are usually affected. Young adults are the most common subjects (cf. Episcleritis), and women more often than men. The causes are obscure, but of the same type as in episcleritis. It is often associated with disturbance of menstruation.

Scleritis is most serious on account of its sequelæ and complications. Uveitis of some kind is probably an invariable accompaniment. It is uncertain whether it may be a result or a cause of the scleritis; most probably it is neither, but both are due to a common cause. This cause is probably the absorption of toxins from some septic focus. It is often difficult or impossible to find the focus, but special attention should be directed to the nasal sinuses and generative organs, the latter especially in women. In other cases the alimentary canal is probably at fault, and many of the drugs which have been found beneficial probably act chiefly by disinfecting the intestinal tract. In some cases the disease is undoubtedly tuberculous and in others syphilitic. Stock has produced scleritis, sclerosing keratitis and lesions in the uveal tract by the intravenous injection of tubercle bacilli in rabbits. There is little doubt that tubercle plays a larger part in disease of these structures in man than has hitherto been recognised. Ciliary staphyloma leads to distortion of the globe, and vision is impaired by it and by the many deleterious effects of the uveitis. Secondary glaucoma often follows.



Fig. 148.—Sclerosing keratitis. (Nettleship.)

Apart from these complications scleritis nearly always extends to the cornea, causing *sclerosing keratitis* (Fig. 148). An opacity develops at the margin of the cornea near the scleritic nodule. It is approximately triangular or tongue-

shaped, the rounded apex being towards the centre of the cornea. Similar opacities may develop farther from the margin and even at the centre. The opacities are grey or greyish yellow, becoming denser until they exactly resemble the sclerotic—hence the name. They are indeed due to changes in the substantia propria, which is embryologically a specially differentiated part of the sclerotic. There is little or no vascularisation, and ulceration never occurs. Some clearing from the centre towards the periphery, as well as near the marginal corneal loops, occurs, but the densest parts usually persist as bluish clouds. The whole margin of the cornea may become opaque like the sclerotic, but the pupillary area almost invariably escapes.

Treatment is the same as for episcleritis. When tubercle is suspected injections of tuberculin may be employed. Dionin and subconjunctival injections of saline, &c., have been recommended. Uveal complications must be treated with atropin, &c. (*See* Chap. XIII.)

Gumma of the Sclerotic is uncommon. It may be indistinguishable in appearance from scleritis, or it may take the form of nodules of various sizes, situated near the limbus, extending backwards to the equator, or even giving rise to an annular scleritis. Gumma may spread to the interior of the eye or a gumma of the ciliary body may spread outwards and involve the sclerotic. Unless active antisyphilitic treatment is adopted early and carried out thoroughly the eye is very likely to be lost from uveal complications, ciliary staphyloma, or phthisis bulbi.

The diagnosis depends upon the history, co-existing signs of syphilis, and the application of the Wassermann test.

Tubercle of the Sclerotic may take the form of a scleritis, may be an extension from the conjunctiva, iris, ciliary body, or choroid, or may be primary, forming a localised nodule which caseates and ulcerates. It should be excised or scraped and the tissue examined for tubercle bacilli.

Annular Scleritis, as already mentioned, may be a form of ordinary scleritis or of gummatous scleritis. A severer type is sometimes known as brawny scleritis and is characterised by a brownish-red, gelatinous-looking swelling surrounding the cornea and extending back towards the equator. It generally occurs in old people and is fortunately rare, for it is little amenable to treatment. Some cases, but not all, give a positive Wassermann reaction.

Ulceration of the sclerotic is always secondary, either from without or from within. Extension from the conjunctiva is almost

always due to tuberculous ulceration, rarely syphilitic. Extension from within is almost always from the iris or ciliary body and is usually tuberculous in the case of the iris, syphilitic in that of the ciliary body. Ulceration of malignant growths which have perforated the sclerotic also occurs –sarcoma of the iris or ciliary body, glioma of the retina. All these conditions are rare.

Blue Sclerotics. The sclerotic is bluish in babies, but a much more pronounced blue coloration is sometimes seen in several members of the same family, is an hereditary condition, and persists throughout life. A curious and hitherto unexplained feature of these cases is that the patients in many of the families also suffered. The sexes are about equally affected ; only those affected can transmit the disease. In a case examined microscopically the sclerotic was about one-third the usual thickness ; the cornea was also thin and Bowman's membrane was absent.

CHAPTER XIII.

DISEASES OF THE IRIS AND CILIARY BODY.

The uncouth term *uveitis* has the merit of emphasising an important fact, viz., the close relationship which exists between the anatomically distinct parts of the uveal tract. It draws attention to the frequency with which inflammatory processes involve the tract as a whole, and are not strictly limited to a single part. This feature is particularly well exemplified in inflammation of the iris and ciliary body. Probably iritis never occurs without some cyclitis, nor, *a fortiori*, cyclitis without some iritis. The disease is called iritis or cyclitis according as the iris or ciliary body appears clinically to be the more affected. The same disposition is also seen with regard to the choroid, though in less degree. General uveitis is commonest in the more chronic types of inflammation, but it is probable that the ciliary body is often involved in many cases which we are accustomed to regard as pure choroiditis.

For convenience of description it is best to consider diseases of the various parts of the uveal tract separately, but the anatomical, physiological, and pathological continuity of the parts can be scarcely too forcibly insisted upon.

Inflammation of the Iris and Ciliary Body.

Iritis.—In order that iritis and the special dangers which attend it may be thoroughly understood, it is necessary to remember the anatomical arrangements of the iris and the pathological changes which occur in it. The iris is practically a diaphragm of blood vessels and unstriped muscle fibres held together by a very loose, spongy stroma. In its perpetual movements the pupillary margin slides to and fro upon the lens capsule. The more the pupil is constricted the more of the posterior surface of the iris is in contact with the lens capsule; when fully dilated the iris probably does not touch the lens at all.

Inflammation of the iris is fundamentally the same process as occurs in other connective tissues: it consists in dilatation

of the blood vessels, impairment of the capillary walls, exudation of a highly albuminous lymph into the tissue spaces, and leuco- or lympho-cytosis. Owing to the extreme vascularity of the iris and the peculiar distribution of the vessels, and to the looseness of the stroma, these generic features of inflammation produce special results. Thus, simple hyperæmia tends to cause the pupil to contract mechanically, on account of the radial disposition of the vessels. This is to some extent physiological, but is greatly increased under pathological conditions. The extreme vascularity and the looseness of the tissues causes an unusually large amount of exudation on the one hand and of swelling on the other. Owing to the greater albuminous content of the fluid its viscosity is increased so that it escapes into the anterior chamber and out of the anterior chamber by way of the filtration angle (*vide* p. 18) with greater difficulty. The iris, from being a partially wrung-out flat sponge, becomes a sponge full of sticky fluid. Hence its freedom of movement is greatly impaired, and the normal reactions become very sluggish or completely abolished. The fluid, too, contains deleterious substances which act as irritants; the nerve endings are stimulated so that the muscle fibres contract. In any case in which the sphincter and dilatator fibres are equally and uniformly stimulated the sphincter overcomes the dilatator, so that constriction of the pupil follows.

It is easy now to understand the chief signs of iritis. The pupil is constricted, partly owing to hyperæmia, partly to irritation; the edge of the pupil is markedly irregular. The reactions of the pupil are sluggish, partly owing to the same causes which induce constriction, partly to what may be termed water-logging. The latter condition causes an alteration in the appearance of the iris. The delicate iris pattern, instead of being clear and sharply defined, becomes blurred and indistinct ("muddy" iris). The colour undergoes considerable change, varying according to the condition of normal pigmentation. In fair people with little pigmentation, the blue iris becomes bluish or yellowish green; brown irides show less difference, but become greyish or yellowish brown. In any case comparison of the colour of the two irides will usually reveal some slight difference, for iritis is generally unilateral during the acute attack.

As a result of the change in colour and blurring of the iris pattern, the hyperæmia of the iris itself is not very obvious, but it manifests itself in circumcorneal ciliary congestion (*vide*

p. 78). This is most marked if the ciliary body is seriously involved, but the conjunctival vessels are also frequently somewhat engorged, so that care is necessary in distinguishing the condition from conjunctivitis. The secondary nature of the conjunctival congestion is shown by the relatively slight discharge : what discharge there is is chiefly lacrymal, never mucopurulent in the absence of actual conjunctivitis as a complication.

The iris is richly supplied with sensory nerves from the ophthalmic division of the fifth nerve. It is not surprising, therefore, that pain is a prominent symptom of acute iritis. It is not confined to the eye, though severe neuralgic pain is felt here, but is also referred to other branches of the first division of the fifth nerve, especially to the brow and parts supplied by the supraorbital and trochlear branches, but also to the cheeks and malar bone, and sometimes to the nose and teeth. It is worse at night.

The albuminous exudates escape slowly into the anterior chamber and mix with the normal aqueous. If the ciliary body is much involved the aqueous itself is abnormally albuminous. The aqueous often contains leucocytes and minute flakes of coagulated proteid, seldom fibrinous networks except in severe cases. It therefore becomes hazy, further interfering with a clear view of the iris and easily mistaken for haziness of the cornea, which is usually not involved. In very intense cases, especially of traumatic iritis with septic infection, large numbers of polymorphonuclear leucocytes are poured out : these sink to the bottom of the anterior chamber and form an hypopyon. Hypopyon is rare in simple iritis without perforation of the globe. Hyphæma, or blood in the anterior chamber, may also occur, but is rare in simple iritis.

The abnormal condition of the aqueous impairs the nutrition of the endothelium which covers the back of the cornea. The cells become sticky and may desquamate in places. The exudates tend to stick to the more affected spots, forming *keratic precipitates* (" keratitis punctata "). These are seldom present in simple iritis, but form a prominent feature of cyclitis, varying roughly with the amount of cyclitis present.

The more albuminous the aqueous the more viscous it becomes. This viscous fluid filters out of the anterior chamber by the filtration angle with difficulty. Hence there is a tendency for the fluid to be retained, so that the intraocular tension rises. The rise is minimal and of no serious import in

cases of simple iritis - in fact, it is scarcely appreciable by clinical methods. If, however, the ciliary body is much involved, the albuminous constituents of the aqueous are very markedly increased, and rise of tension may be so great as to endanger the sight, requiring special attention.

The exudates which are poured out by the iris and ciliary body are naturally most concentrated in their immediate neighbourhood. They cover the surface of the iris as a thin film and spread into, and often completely over, the pupillary area. In this manner the pupil may become " blocked " with exudates, a condition which very seriously impairs the sight of the eye. Moreover, the exudates tend to stick the iris down to the lens capsule, so that it becomes immovably fixed.

If the patient is seen in the early stages and atropin is instilled the pupil dilates, though not so readily as a normal pupil, on account of the water-logging of the iris. By continuous treatment the iris may be freed from the lens capsule and the pupil become completely dilated and circular.

FIG. 149.—Diagram of seclusion and occlusion of the pupil, with bowing forwards of the iris (" iris bombé "). (Nettleship.)

The exudates, however, show a great tendency to become quickly organised. This is particularly seen in most cases of iritis, less frequently in cyclitis ; when it is a very marked feature of the case the inflammation is often described as *plastic*. If the exudates which bind the iris to the lens capsule have not been already broken down they become converted into fibrous bands which atropin is wholly unable to rupture. Such firm adhesions of the pupillary margin to the lens capsule are called *posterior synechiæ* (συνέχειν, to hold together). When they are present a mydriatic causes only the intervening portions of the circle of the pupil to dilate and the pupil assumes a festooned appearance (Plate VI., Fig. 2 ; Fig. 153). Even in the absence of a mydriatic minute inspection will generally show irregularities of the pupillary margin due to the synechiæ. Such an irregular pupil is a sign of present or past iritis. For diagnostic purposes homatropin should be instilled and the result confirmed by the appearance of the dilated pupil. Owing to the contraction of organising exudates upon the surface of the iris the retinal pigment epithelium may be pulled outwards over the pupillary margin. Patches of pigment are then seen in this situation (ectropion of the

uveal pigment). Posterior synechiæ show some predilection for the lower part of the pupil in the early stages, probably owing to gravitation of the plastic exudates.

In very severe cases of plastic iritis, or after recurrent attacks, the whole circle of the pupillary margin may become tied down to the lens capsule. The condition is called *annular* or *ring synechia* or *seclusio pupillæ* (Figs. 149, 152); it is one

Fig. 150.—Total posterior synechia (× 7), from a case of plastic irido-cyclitis commencing to cause phthisis bulbi. The iris is completely adherent to the lens capsule, and the periphery is retracted. There is a delicate inflammatory pupillary membrane (occlusio pupillæ). There is also an anterior capsular cataract, due to inflammation; it contains calcareous deposits, as shown by the patch of dark staining. The ciliary body is degenerated and detached from the sclerotic at the posterior part. The retina is completely detached and folded behind the lens.

of great danger to the eye (*vide* p. 265). In similar cases, especially if the case has been neglected and the pupil not well dilated by atropin at an early stage, the exudates in the pupillary area may also organise. A film of opaque fibrous tissue then fills the pupil; this condition is called blocked pupil, or *occlusio pupillæ* (Figs. 149, 152). If there has been much cyclitis the posterior chamber (Fig. 1) also becomes filled with exudates which may organise. They then tie down the

whole of the back of the iris to the lens capsule : this condition is called *total posterior synechia* (Fig. 150). It leads to retraction of the peripheral part of the iris, so that the anterior chamber becomes abnormally deep at the periphery, sometimes much deeper than in the centre.

Iritis is most frequently mistaken either for conjunctivitis or for acute glaucoma. The points which distinguish it from conjunctivitis will be gathered from what has already been described. The error of mistaking iritis for glaucoma is the most serious which can be made, more particularly because the treatment of the two conditions is diametrically opposed. Dilatation of the pupil with atropin, which is urgently necessary in iritis, is the worst possible treatment of glaucoma (*q.v.*). At the cost of some repetition, the distinguishing features will be given here.

(1) In iritis the pupil is smaller than normal and irregular ; in glaucoma it is larger and oval, usually with the long axis vertical.

(2) In iritis the intraocular tension is scarcely appreciably raised unless much cyclitis is present ; in glaucoma it is always appreciably raised, and is often very high.

These are the two chief objective signs. Cases occasionally arise which are doubtful even to the most experienced. A useful and harmless procedure which will almost invariably settle the question is the following : A drop of 2 per cent. homatropin (*not* atropin) solution is instilled. As the pupil dilates, in iritis the irregularities are emphasised and synechiæ are almost always revealed ; the tension is not appreciably affected by the mydriatic ; in glaucoma the pupil probably dilates slowly but quite evenly, retaining its roundness ; the tension is appreciably increased by the mydriatic. As soon as glaucoma is definitely diagnosed by this test eserin (1 per cent.) must be *immediately* instilled and repeated at intervals (*vide* p. 282). Atropin is never to be used for diagnostic purposes, since eserin is incapable of counteracting its mydriatic effect, and if the case is found to be one of glaucoma immediate iridectomy would be imperative.

(3) The subjective symptoms differ in the two diseases. Vision is usually more impaired in glaucoma than in iritis. In acute glaucoma the onset of the pain is sudden, and it is so severe that it is frequently accompanied by vomiting.

It will be advisable again to enumerate the chief symptoms and signs of iritis. The subjective symptoms are : pain, of a neuralgic character, referred not only to the eye but also to the

supra-orbital region; dimness of vision, due to cloudiness of the aqueous, exudates in the pupillary area, &c. The objective signs are: constriction and irregularity of the pupil, unless a mydriatic has already been instilled, in which case the irregularity is emphasised by the presence of the posterior synechiæ. If the mydriatic is instilled early these synechiæ may break down so that the pupil again becomes quite round; in such cases spots of lymph or pigment upon the anterior capsule of the lens often leave permanent marks of old synechiæ, and form most valuable evidence of previous iritis (Fig. 151). It has already been pointed out (p. 4) that the posterior layer of the retinal pigment epithelium on the back of the iris is less firmly attached to the iris than the anterior. When a synechia breaks down some of the posterior layer often tears away and remains attached to the lens capsule: these pigmented spots never disappear entirely. Discoloration or muddiness of the iris, whereby the iris pattern is masked; hyperæmia, manifesting itself chiefly as circumcorneal ciliary congestion; exudations, manifesting themselves either as more or less cloudiness of the aqueous or as solid deposits in the pupillary area and upon the iris: these are conspicuous signs of inflammation of the iris.

Fig. 151.—Spots of pigment on the lens capsule, left by posterior synechiæ which have broken down. (Nettleship.)

The course of iritis varies with its intensity. Even the slighter acute cases take three or more weeks before inflammation entirely subsides. The best sign is the prompt action of atropin, for in the worst cases it has little or no effect. Improvement is shown by good dilatation, diminution of injection and pain. In the chronic cases the ciliary body is almost always more seriously involved; the condition is one of iridocyclitis. The inflammatory signs are less, but diminution of vision is progressive, and the disease not infrequently lasts for years.

One of the most characteristic features of iritis and cyclitis is the great tendency to relapse. It depends, not upon the synechiæ, as was once taught, but upon the constitutional cause of the disease. Each fresh attack runs a similar, though usually less severe, course, often leaving fresh traces and increased impairment of vision.

Complete resolution may occur in slight cases treated early and suitably. The exudates become absorbed; the synechiæ break down, leaving only such slight traces that vision may be perfect. Comparatively slight cases may, however, leave very

serious results if they are improperly treated, and in severe cases these are the rule. Most of the evil results are attributable to neglect of or impossibility of early dilatation of the pupil, which causes permanence of the posterior synechiæ. If these are few, no special injury or impairment of sight follows, but future attacks are more likely to result in an increased number or in ring synechia.

Ring synechia, or seclusio pupillæ, is one of the worst sequels of the disease, since, if unrelieved, it inevitably leads to



FIG. 152.—Iris bombé, with very broad peripheral anterior synechia, annular posterior synechia (seclusio pupillæ), and inflammatory pupillary membrane (occlusio pupillæ). There is also an anterior capsular cataract.

secondary glaucoma and destruction of sight. Owing to the complete shutting off of the pupil the aqueous which is secreted by the ciliary body (*vide* p. 17) is unable to pass forwards into the anterior chamber, the iris being a non-permeable membrane. It therefore collects behind the iris, which becomes bowed forwards like a sail, a condition which is called *iris bombé* (Fig. 152). Regarded from in front, the anterior chamber is seen to be funnel-shaped, deepest in the centre and shallowest at the periphery. The filtration angle is obliterated by the adhesion of the iris to the cornea and sclera at

the periphery (peripheral anterior synechia). Hence the fluid is retained within the eye and the intraocular tension rises (*vide* p. 265).

Organisation of the exudates in the pupillary area leads to the formation of an inflammatory pupillary membrane or occlusio pupillæ. This interferes directly with the transmission of light rays and is often associated with ring synechia, with or without total posterior synechia (*vide* p. 245). In such cases there is generally plastic cyclitis and the eye is irretrievably affected.

Repeated attacks of iritis lead to atrophy of the iris, which becomes dirty grey, like felt or blotting paper. Red streaks often mark the site of permanently dilated vessels, usually of new formation and therefore not necessarily radial in direction. The pupillary margin is thin and frayed: the reactions are diminished.



Fig. 153. — Plastic iritis with nodules in the angle of the anterior chamber. (Nettleship, from a drawing by Holmes Spicer.) The patient was not syphilitic.

Varieties of Iritis.—The varieties of iritis are usually divided into primary and secondary, the latter being those which are due to extension of inflammation from some other part of the eye, usually the cornea. The primary iritides are due to some general dyscrasia, though it is not always possible to determine its exact nature (so-called idiopathic iritis). The most undoubted cause is certainly syphilis; other causes are gonorrhœa, tubercle, and diabetes; alleged causes are gout, "rheumatism," acute exanthemata, &c. Finally, iritis is usually an important element of sympathetic ophthalmia (*q.v.*).

Syphilitic Iritis manifests itself in two forms. Syphilis is the commonest cause of simple plastic iritis, which occurs in the secondary stage, soon after the skin eruptions, usually within the first year after infection, but not before the third month. There may be nothing characteristic about this form of iritis, or nodules may occur upon the iris (*vide infra*). Syphilitic iritis lasts two to eight weeks and does not usually recur, thus differing from the "rheumatic" form.

Iritis occurs in at least 3—4 per cent. of syphilitics, and of cases of iritis syphilis accounts for at least 25—30 per cent. Syphilitic iritis is generally unilateral, but the second eye becomes affected sooner or later in about a quarter of the

cases. In iritis due to other causes it is unilateral in only about 10 per cent. Syphilitic iritis attacks males more than females, and three-fourths of the cases are between twenty and forty years of age. The Wassermann reaction is of great value in settling the diagnosis. The spirochæte has been found in the aqueous.

Simple plastic iritis also occurs in congenital syphilis, usually as an accompaniment of interstitial keratitis (*q.v.*). It also occurs in very young babies with congenital syphilis, without any corneal complication, but usually with large nodules or gummata on the iris. This is not common, but is probably the only cause of iritis in very young children apart from direct injury. It is sometimes seen so soon after birth that almost certainly in these cases it commences as an intrauterine inflammation. The average age of onset is five to six months. It is commoner in females than in males, and is unilateral in about half the cases. (Hutchinson.)

Fig. 154.—Nodules occurring in the secondary stage of syphilis, situated at the pupillary border of the iris. (Nettleship, from a drawing by W. G. Laws.)

The iris also becomes inflamed in some cases of acquired syphilis late in the secondary or very rarely during the tertiary stage. These cases are characterised by the formation of yellowish red nodules near the pupillary and ciliary borders of the iris, but not in the intermediate region. The nodules are usually multiple and vary in size from that of a pin's head upwards (Fig. 154). It has been customary to consider these nodules either condylomata or gummata, according to the stage of the complaint. There is no good ground, either clinical or pathological, for the distinction, and the term *gummatous iritis* may be used for all these cases. There is generally much exudation in gummatous iritis, and broad synechiæ are formed. The nodules are liable to be mistaken for tubercle (*q.v.*) or sarcoma (*q.v.*); the absence of iritis and the presence of only a single nodule usually distinguishes the latter condition, which, moreover, is extremely rare. Gummatous iritis may rarely extend to the corneo-sclera at the angle of the anterior chamber and lead to perforation of the globe.

The sites of previous gummatous deposits in the iris are marked by depigmentation of the stroma, probably owing to

stretching. Whitish spots in the ciliary region of the iris, especially near the crypts, occur in the early stages of syphilitic infection without previous iritis. Syphilitic degeneration of the vessels causes thickening of the coats, sometimes making them appear as white lines. Atrophy of the muscle fibres, especially of the sphincter, also occurs, and a circle of atrophic patches near the pupillary border is strong evidence of syphilis.

Vision is usually impaired permanently in syphilitic iritis, and the prognosis is worse in the gummatous type. Specific iritis is indeed a sign of severe syphilis ; Trousseau found that thirty-four out of forty patients ultimately developed grave sequelæ such as tabes and general paralysis.

Gonorrhœal Iritis is probably more common than is generally supposed. It occurs especially in those cases which have gonorrhœal "rheumatism," and seldom supervenes until after one attack of arthritis, usually in the knees. It exhibits no special signs ; it tends to recur, often during the onset of a gleet or arthritic attack. There is little doubt that it is a metastatic infection. The patients are almost always men, and as a rule both eyes are affected, though not at the same time. Extension to the ciliary body may be indicated by fine vitreous opacities, but involvement of the choroid is very rare. Another more characteristic form may occur during the acute attack. The exudates into the anterior chamber then have a peculiar gelatinous appearance.

"*Rheumatic*" *Iritis*.—In patients with iritis, in whom no specific or gonorrhœal taint can be discovered, a history of rheumatic pains in the muscles and joints can often be elicited. Iritis seldom accompanies an attack of acute rheumatism, and only rarely can a history of such an attack be obtained. The patients are often gouty or have rheumatoid arthritis. What the pathological relationship of the iritis is to these complaints must remain a matter of conjecture until their ætiology is placed beyond dispute. It is most probable that both the "muscular rheumatism" and the iritis are due to a common cause, toxins derived from a septic focus in some part of the body, *e.g.*, mouth, nasal sinuses, intestinal tract, etc. The iritis in these cases is usually a plastic iritis of moderate severity. It often attacks both eyes and shows a very marked tendency to recur, and the recurrence seems to bear a direct relationship to the recurrent attacks of pain or arthritis. There is often an unusual amount of conjunctival and episcleral congestion. Iritis in an elderly patient is likely

to be gouty, often starting suddenly in the night and sometimes ushering in an acute attack of gouty arthritis.

Diabetic Iritis is rare : it is probably a gouty iritis occurring in a diabetic subject. It shows a marked liability to the formation of new or enlarged vessels in the iris, with the formation of plastic exudates and occasionally an hypopyon. On the whole it runs a favourable course.

Tuberculous Iritis occurs in a miliary and a conglomerate or solitary form. In the former there is usually a yellowish white nodule surrounded by numerous smaller satellites ; there is the same tendency as in gummatous iritis for the nodules to be near the pupillary or ciliary margins. In the earliest stages the nodules are minute, greyish, and translucent. There are often spots of " k.p." on the back of the cornea, indicating involvement of the ciliary body. In conglomerate tubercle there is a yellowish white tumour, though smaller satellites may be present. The nodules contain giant cells. There is usually less general iritis than in the gummatous form, but there is almost always some. The condition may be mistaken for gummatous iritis or for sarcoma. The absence of specific history, a negative Wassermann reaction, the failure of anti-syphilitic treatment, and the age of the patients children or young adults –are features distinguishing it from gummatous iritis. The presence of satellites, the usual, but not invariable, absence of visible vessels upon the surface of the nodules, the age of the patient, and the presence of iritis distinguish it from sarcoma. The diagnosis may be extremely difficult, but the great rarity of sarcoma of the iris should be borne in mind. Simple iritis is said to be sometimes tuberculous ; this form is very rare.

In conglomerate tubercle of the iris the corneo-sclera at the angle of the anterior chamber almost invariably becomes ultimately eroded and involved. The wall of the globe is thus weakened and eventually gives way. The tuberculous mass then grows rapidly through the perforation, and a large portion of the iris may become prolapsed. In this manner the eye is inevitably lost.

von Pirquet's cutaneous reaction may be applied to doubtful cases. A positive result is of little value except in children, but a negative result eliminates the diagnosis of tubercle with a fair degree of certainty. Wolff-Eisner or Calmette's conjunctival test should not be used on account of the danger of severe reaction. Subcutaneous injection of Koch's old tuberculin gives a characteristic rise in temperature, &c., in the

presence of tubercle, but there is no proof that the ocular lesion is the cause of the reaction, and the test is not free from danger.

Sympathetic Iritis.—See Sympathetic Ophthalmia, Chap. XXI.

"*Serous*" *Iritis.*—See p. 257.

Secondary Iritis: Purulent Iritis.—See Chap. XXI.

Treatment.—Dilatation of the pupil with atropin and hot applications are the essentials of local treatment. Atropin acts in three ways: (1) by keeping the iris and ciliary body at rest; (2) by diminishing hyperæmia; (3) by breaking down posterior synechiæ and preventing the formation of fresh ones. It may be used in the form of drops of a 1 per cent. solution or as an ointment of the same strength. I prefer the ointment for the following reasons: (1) its action is more continuous; (2) it is easier to apply, since it usually works into the eye even if only rubbed along the lashes; (3) it is less likely to cause symptoms of poisoning, which are not uncommon with the drops in children. Symptoms of poisoning—dryness of the throat, flushing of the face, delirium, &c.—are due to the excess of solution—often considerable in unskilful hands—passing down the nasal duct into the nose and throat. The dose administered in this manner is never lethal.

Atropin should be pushed in the early stages, best by frequency of application rather than increased strength. Every four hours is usually sufficient. When the pupil is well dilated, two or three times a day suffices. If atropin irritation ensues, hyoscin, scopolamin, or duboisin should be substituted. Dionin, 5–10 per cent., may be used in conjunction with the mydriatic.

Hot applications are extremely grateful to the patient, diminishing the pain, and are of therapeutic service in encouraging a more vigorous blood and lymph flow. Hot fomentations and bathings may be used, but dry heat applied to the surface of the closed lids has the same effect. This may be done by means of a Japanese muff-warmer or an electric heater, which is bandaged over a pad of wool, previously well heated and applied to the eye. By this means the heat is considerable and continuous, with a minimum of trouble and discomfort.

In very severe cases, or when the pupil does not readily respond to atropin, depletion of blood from the temple should be resorted to. The best method is by two or three leeches applied a short distance outside the external canthus. They

should not be too far from the eye, nor too near, for in the latter case much œdema of the lids may follow. Heurteloup's artificial leech may be used, but is not so efficacious.

If the pain is very intense a hypodermic injection of morphia may be given. Aspirin is very useful in relieving pain.

General treatment should be commenced by a smart saline purge, and the bowels must be kept freely open throughout the acute stage.

In other respects the general treatment depends upon the cause. In syphilitic iritis the patient is rapidly got under the influence of mercury, best by inunctions or the intramuscular injection of Lambkin's cream. Salvarsan, gallyl, &c., cause rapid improvement, but these cases also respond well to mercury. These drugs are most effectual in the cases occurring during the secondary stage, but should also be used in the gummatous form. Here they should be supplemented by iodides, but these must not be given simultaneously with injections of metallic mercury. The infantile form of acute syphilitic iritis responds rapidly to mercury, but neither drug is very efficacious in the congenital type, and in this, as well as in the later stages of the other forms, general tonic treatment is indicated.

In gonorrhœal iritis gonococcic vaccine sometimes produces good results.

In other forms of iritis, where a rheumatic taint is suspected or where no satisfactory cause can be found, it is usual to order salicylic preparations, and they appear to do good, not only in these, but also in gonorrhœal and diabetic iritis. An exhaustive search should be made for any septic focus in the body.

In the convalescent stage smoked glasses are ordered—for both eyes, especially on account of the consensual reaction to light. Atropin, or its equivalent, should be continued for at least ten days or a fortnight after the eye appears to be quiet, otherwise a relapse is very likely to occur.

Tuberculous iritis is treated by the same local applications as other forms. The usual constitutional treatment must be pushed. Improvement and even complete resolution have been recorded from the use of tuberculin injections, but they should be used cautiously, commencing with very small doses, since a violent reaction may have a serious effect upon the eye. The dose is very gradually increased.

Some authors consider that tuberculous iritis is generally the primary manifestation of the disease in the body. They

therefore advocate the removal of the eye as soon as the diagnosis is made, in order that the danger of extension of the disease to other parts of the body may be minimised. If perforation of the globe has taken place and the eye is irretrievably lost immediately excision should be urged, but in other cases ordinary treatment, supplemented by tuberculin, should be persisted in. Good results have undoubtedly been obtained by the use of tuberculin, and it should be given a persevering trial; nevertheless it often fails to ameliorate the condition.

Treatment of Sequelæ and Complications.—For the treatment of coincident cyclitis see p. 259.

Firm posterior synechiæ can sometimes be broken down by placing a small particle of solid atropin in the conjunctival sac. Care must be taken to prevent the dissolved atropin from passing down the nasal duct by pressure with the finger upon the lacrymal sac by the patient himself, but the surgeon must see that the pressure is rightly applied and kept up.

Annular synechia demands an iridectomy in all cases in order to restore communication between the anterior and posterior chambers, and thus avoid the supervention of secondary glaucoma (*vide* p. 265). In some cases it is necessary or advisable to be content with making a puncture or transfixion of the iris by a broad needle. No operative procedure of this kind must be undertaken during an acute or even the slightest attack of iritis if it can be possibly avoided, since the traumatic iritis set up will frustrate the object of the operation by filling the coloboma with exudates, and may even cause the loss of the eye. It is best, if possible, to forestall a ring synechia by performing the iridectomy before the adhesion extends round the whole circle. This can often be done, because operable ring synechia, *i.e.*, ring synechia without total synechia, is frequently the result of recurrent attacks of quiet or almost painless iritis, during each of which more of the circle is involved. The iridectomy is performed during a quiescent interval. It is often difficult to get a good coloboma owing to the atrophy and friability of the iris and the firmness of the adhesions. Hæmorrhage is common, and the hyphæma takes longer than usual to be absorbed. Iridectomy sometimes has a beneficial effect on recurrent iritis, but should not be done without special indications until all other non-operative measures have failed. The presence of "k.p." should generally be regarded as a contra-indication to intra-ocular operation.

Total posterior synechia can seldom be operated upon with success. Iridectomy is seldom possible, and the only procedure which can be adopted is extraction of the lens by a specially devised operation. Iridectomy or iridotomy may be possible if the lens is shrunken.

Cyclitis has already been referred to incidentally. The exudates from the ciliary body pass into the anterior chamber directly from that part which forms a boundary of the chamber (Fig. 2), and by passing forwards through the pupil. When they organise in the plastic form they not only cause total posterior synechia, but also surround the lens and extend



Fig. 155.—Phthisis bulbi, due to irido-cyclitis (× 3). *C*, cornea; *Ir.*, iris; *L*, lens; *C.b.*, ciliary body; *Ch.*, choroid; *R*, retina, detached and folded up behind lens, separated from choroid by albuminous coagulum; *Scl.*, sclerotic.

throughout the vitreous. Behind the lens they form a transverse membrane or cyclitic membrane. Strands of fibrous tissue are formed in the vitreous. They become anchored to the retina in various places, and by their subsequent contraction often lead to detachment of the retina. The exudates which organise upon the surface of the ciliary body cause the destruction of the ciliary processes, which results in abolishing or seriously diminishing the secretion of aqueous. Hence the intraocular tension becomes lowered and the eye is quite soft to the touch. The walls of the globe fall in, and the eye becomes shrunken—*phthisis bulbi* (φθίνω, to waste) (Fig. 155). After this has occurred degenerative changes supervene, and

the choroid becomes converted after months or years into bone.

Chronic Irido-cyclitis (*Syns.—Simple Cyclitis*, "*Serous Iritis*").—This is an extremely insidious and chronic disease, characterised by diminution of vision with slightly marked physical signs. In severe cases there is ciliary congestion, tenderness on pressure over the ciliary region, deep anterior chamber, precipitates ("keratitis punctata") on the back of the cornea, and dust-like opacities in the vitreous. Posterior synechiæ are not a conspicuous feature, but are liable to be formed slowly and insidiously. The tension is usually slightly raised in the earlier stages, lowered in the later. There may be œdema of the upper lid and neuralgic pain in the eye and brow. There is sometimes myopia, owing to irritation of the ciliary muscle.



Fig. 156. — Typical arrangement of the spots on the back of the cornea in "keratitis punctata" (k.p.). (Nettleship, from a sketch by Dr. Herringham.)

The *keratic precipitates* ("k.p.") (Fig. 156) consist of leucocytes which are deposited from the aqueous upon the back of the cornea and stick there. They may contain pigment granules, showing their origin from the uveal tract (pigmented k.p.). In the most characteristic form they are scattered over a triangular area of the lower part of the cornea, the smaller spots being above, the larger below (Fig. 156). This arrangement is due to gravitation of the particles towards the bottom of the anterior chamber, combined with the perpetual movements of the eye, which are mostly in the horizontal direction. The typical arrangement is often but not always observed. More commonly a few isolated spots are seen scattered irregularly over the lower part of the cornea. They require great care in examination for their discovery (*vide* p. 82), and their importance cannot be over-estimated. The smaller spots frequently coalesce, forming small plaques, which gradually become more translucent ("mutton-fat k.p."). Precipitates are generally sharply defined and can thus be distinguished from opacities in the deeper layers of the cornea. They are more likely to contain pigment when the iris is brown, but "pigmented k.p." also occurs with blue or grey irides if the inflammation lasts a long time. Pigmentation therefore may give some indication of the duration of the disease. The pigment persists almost indefinitely. Similar precipitates are

rarely seen upon the anterior capsule of the lens, but the leucocytes do not readily stick here owing to the smooth surface, devoid of endothelium.

The vitreous opacities are of the same nature, viz., wandering leucocytes, but many are also probably minute particles of albuminous exudate. Their mobility in the vitreous shows that the consistency of this substance has undergone change, sometimes amounting to complete fluidity, due to defective nutrition.

The depth of the anterior chamber is an important sign, not easily explained. It is undoubtedly connected with the deficient filtration of the albuminous aqueous through the angle, which also accounts for the rise in tension. The peripheral part of the anterior chamber is often particularly deep, even deeper than the central: this is however more marked in the later stages of plastic cyclitis, when it is due to mechanical retraction of the iris from organisation of exudates in the posterior chamber.

In severe or prolonged cases the deeper layers of the cornea may become infiltrated, as in interstitial keratitis, though seldom to the same extent. This is specially liable to occur in tuberculous cases, and in these there are often minute greyish nodules on the surface of the iris. These should be carefully looked for, as they are of considerable diagnostic significance, and the prognosis is correspondingly grave.

In the slightest and most insidious cases of irido-cyclitis the symptoms and physical signs are minimal. Considerable diminution of vision without obvious cause should always excite apprehension, and the cornea should be most carefully explored by oblique illumination with magnification by the loupe, as well as by the direct method with a strong convex lens. A few spots of k.p. are decisive proof of cyclitis, and may be the sole physical sign. Change in the colour of the iris, due to atrophy, is an important sign, since it may at once attract attention, especially if the normal eye has a brown iris.

Chronic irido-cyclitis occurs under similar conditions to simple iritis and is commoner in women than in men. Syphilis or tubercle may be the cause. It is probably always due to some form of toxæmia or bacterial metastasis, and a careful search should be made for a septic focus in some part of the body. It is most frequently found in the mouth—pyorrhœa alveolaris—in the nose and accessory nasal sinuses, or, especially in women, in the genital tract. Very often no such source of toxins can be demonstrated, but the patients are usually

of a debilitated type. There is often constipation, and it is not improbable that the intestinal tract is a frequent source of the toxins.

The disease is generally very chronic and liable to exacerbations with gradual and insidious formation of posterior synechiæ. Vision is greatly diminished during the more acute stage, but recovers considerably in the intervals, but each recurrent attack leaves more and more permanent defect. The eye may finally become soft and tender, and enter into the condition of phthisis bulbi, but this occurs only after many years in simple cases of irido-cyclitis. Less serious cases, however, not uncommonly occur, especially when the septic focus, *e.g.*, pyorrhœa aveolaris, is discovered and is amenable to radical treatment.

Treatment of irido-cyclitis is essentially the same as that of iritis, but special attention must be directed to any septic foci which may be found and to the general health. Septic foci such as pyorrhœa alveolaris, nasal, genital, or urinary sepsis, furunculosis, &c., must be radically treated when possible. In some cases an autogenous vaccine has produced rapid cure, but it more often fails. The cases often drag on indefinitely, with occasional exacerbations. During the more acute phases energetic treatment with atropin, hot bathings or the muff-warmer, and if necessary blisters or leeches, is indicated. Small doses of calomel (gr. $\frac{1}{4}$, three times a day) or salol and salicylates do good in any cases of obscure ætiology, probably by acting as intestinal disinfectants. Iodides help to cause absorption of vitreous and other exudates. In severe cases the patient should be kept in bed and submitted to mercurial inunctions or baths, which often do good in cases in which there is no specific history. Diaphoresis by vapour baths and hypodermic injections of pilocarpin may be tried in intractable cases, which form the majority. Cases of syphilitic or tuberculous origin, of course, require the appropriate courses of treatment, but tuberculin should be used with great caution (*vide* p. 255).

If the intraocular tension is raised seriously, so that there is danger to the sight of the eye from this cause, it must be relieved by paracentesis (*vide* p. 199). It usually has only a transitory effect, and may require to be repeated every two or three days. It is theoretically unsatisfactory, since the comparatively sudden reduction of the intraocular pressure to zero causes dilatation of the ciliary vessels and allows the passage of a lymph which is even more albuminous than that

which has been let out. It should therefore not be resorted to unless imperatively indicated. On the other hand, the final result is often very satisfactory, probably because the rapid flow of lymph flushes out the secretory channels and carries away endothelial and epithelial débris and stagnant toxins. If repeated paracentesis fails to relieve the tension, it may be necessary to do an anterior sclerectomy (trephining), but this should only be done as a last resource.

During the intervals between exacerbations the pupils should be kept moderately dilated with weak (0·5 per cent.) atropin. The patient should have plenty of fresh air, good diet, and tonics.

If the eye becomes useless, shrunken, and painful it may be necessary to excise it.

Plastic Irido-cyclitis.—The pathology of this condition has already been described incidentally. In it the signs of irido-cyclitis in general are increased. The cyclitic membrane behind the lens may be seen with the ophthalmoscope or even by oblique illumination. In young children the condition forms one type of pseudo-glioma (Chap. XIX.). In the later stages the degenerative changes in the ciliary body prevent it from fulfilling its functions of supplying the eye with lymph and nutriment. The vitreous suffers first, becoming fluid, and later the lens, which becomes opaque. Finally the eye shrinks (phthisis bulbi).

Treatment is the same as for chronic irido-cyclitis. The blind, shrunken globe is often painful and a continual source of annoyance to the patient. It should, under these circumstances, be excised.

Gumma of the Ciliary Body causes an intense acute plastic irido-cyclitis, with severe iritis, much exudation into the anterior chamber, often deep infiltration of the cornea and usually great pain. It is a rare complication of syphilis, not confined to the tertiary stage. It varies in the severity of the symptoms and the rate of progress. It is only to be diagnosed clinically with certainty when the inflammation extends into the sclerotic (*vide* p. 239). If it does not respond to active anti-syphilitic treatment the eye eventually shrinks.

Tubercle of the Ciliary Body occurs with tubercle of the iris and of the choroid, and is usually only to be inferred clinically rather than definitely diagnosed. As already mentioned the tubercle bacillus may account for some cases of chronic cyclitis.

Purulent Irido-cyclitis.—See Chap. XXI.

Sympathetic Irido-cyclitis.—See Chap. XXI.

Congenital Abnormalities of the Iris.

One iris may have a different colour from the other, or parts of the same iris may differ in colour. Both conditions are known as *heterochromia iridis*. The blue iris is due to the absence of pigment in the iris stroma, the pigment in the retinal epithelium being seen through the translucent stroma. Light irides often have patches of brown pigmentation. These benign *melanomata* are due to abnormal groups of retinal pigment epithelium lying in the posterior layers of the stroma.

The pupil is normally slightly to the inner side of the centre of the cornea. In some cases it is considerably displaced, usually also to the nasal side *corectopia* (κόρη, pupil, ἐκ, out of, τόπος, place). Rarely there are other holes in the iris besides the pupil—*polycoria*.

The iris may be apparently absent *-aniridia* or *irideremia*. Anatomical investigation has shown that there is always a narrow rim persistent at the ciliary border, but it is hidden from view during life by the sclerotic. On examination the ciliary processes and the suspensory ligament of the lens can be seen. Aniridia is usually bilateral. There is a tendency for secondary glaucoma to develop in these eyes.

Fig. 157. — Congenital coloboma of the iris.

There may be a gap in the iris, usually pear-shaped or like a Gothic arch, continuous with the pupil and extending towards but not always as far as, the ciliary border. It is called a *congenital coloboma* (κολόβωμα, mutilation) of the iris (Fig. 157). It is usually downwards or down and slightly in, corresponding with the position of the fœtal, so-called choroidal cleft: such a coloboma is called typical. Colobomata are found in other directions, and are then atypical. Typical coloboma of the iris is often associated with typical coloboma of the choroid (*q.v.*). It is one of the commonest congenital malformations of the eye.

Persistent Pupillary Membrane is due to persistence of part of the anterior vascular sheath of the lens, a fœtal structure which normally disappears shortly before birth. Fine threads stretch across the pupil, or may be anchored down to the lens capsule (Plate VI., Figs. 3, 4). They are distinguished from post-inflammatory synechiæ in always coming from the anterior surface of the iris just outside the pupillary margin, *i.e.*, from the position of the corona or circulus iridis minor.

Such tags are of frequent occurrence and are of no pathological importance. They are commonest in young children and probably undergo further absorption as age advances.

The fœtal pupillary membrane consists of a network of minute bloodvessels supported by a very delicate stroma which contains pigment cells. Sometimes the pigment is left on the lens surface and persists. It forms a multitude of very fine brown dots scattered over a circular area 5 or 6 mm. in diameter in the centre of the pupil. These spots are distinguished from pigment spots left by posterior synechiæ which have broken down (*vide* p. 247) in being much smaller, much more numerous and very regularly arranged, and also by the absence of any concomitant signs of iritis. They do not usually interfere appreciably with vision.

Tumours and Cysts of the Iris.

Tumours of the Iris and Ciliary Body.—See Chap. XX.

Cysts of the Iris.—*Serous cysts* of the iris sometimes occur, and are due to closure of iris crypts, with retention of fluid. *Cysts of the retinal epithelium* occur, due to accumulation of fluid between the two layers of retinal epithelium. They look like a bombé iris limited to parts of the iris—a limitation which is impossible in the case of true bombé iris (*q.v.*). In these cases the posterior layer of epithelium is often adherent to the lens. *Implantation of epithelium* on the iris sometimes occurs after perforation wounds or operations, giving rise to pearl cysts or cholesteatomata. The epithelium may spread over the iris and line the whole anterior chamber, causing glaucoma. Eyelashes are sometimes carried into the anterior chamber in perforating wounds and lodge upon the iris.

Fig. 1.



Fig. 2.

PLATE VIII. (*To face p.* 263.)

Fig. 1.—Inferior crescent ("Fuchs's coloboma"). Note the distortion of the disc, which is unusually well marked in this case.

Fig. 2.—Cupped disc in glaucoma. R. and L. discs from a case of primary glaucoma. Note the bending of the vessels in the R. eye, the scleral rings, and the exposure of the laminæ cribrosæ.

CHAPTER XIV.

GLAUCOMA.

Glaucoma is a symptomatic condition, not a disease *sui generis*. The characteristic symptom, or rather physical sign, is increased intraocular pressure. It will be clear from the description of the mechanism whereby the normal intraocular pressure is maintained (*vide* p. 14) that increase may be due either to (1) increased production of lymph associated with normal or diminished outflow, or to (2) diminished outflow associated with normal or increased inflow. Though the factors which bring about increased production of aqueous cannot be eliminated, yet it may be taken as certain that pathological increase of tension is almost invariably due to defective outflow.

Two great classes of cases in which the tension is pathologically increased can be distinguished, viz., (1) those in which the tension is only moderately increased, in which the anterior chamber is deep, and in which there are more or less definite signs of inflammation of the ciliary body (Chap. XIII.); and (2) those in which all grades of increased tension are met with, in which the anterior chamber is shallow, and in which, though there may be very evident signs of congestion and irritation, any definite signs of ciliary inflammation are either absent or secondary in onset. It is well to keep these two groups quite separate, since their pathogenesis is different and the differences in clinical course and treatment are marked. The term glaucoma should be limited to the second group.

True glaucoma may be conveniently divided into two forms, primary and secondary. Primary glaucoma, acute and chronic, has been called idiopathic, since it comes on without any apparent fundamental cause. Since the pathology of secondary glaucoma has been fairly well elucidated and throws some light upon that of primary, the former will be considered first.

Secondary Glaucoma.—We have seen that the increased tension of irido-cyclitis is due to defective filtration of the

aqueous at the angle of the anterior chamber owing to the viscous nature of the fluid. Secondary glaucoma is almost invariably due to mechanical obstruction at the same spot. Usually the obstruction takes the form of adhesion of the iris to the back of the corneo-sclera. This peripheral anterior synechia causes the obliteration of the network of the ligamentum pectinatum iridis, and prevents the fluid from gaining access to the canal of Schlemm. It is therefore imprisoned within the eye and the intraocular pressure rises. In order that this may occur it is necessary that the angle should be obliterated over a considerable part of its circumference, but there is a great tendency in most cases for the cause which has produced partial peripheral anterior synechia eventually to complete the circuit. In some cases in which secondary glaucoma supervenes there is no actual synechia, but the meshes of the ligamentum pectinatum are choked with leucocytes, pigment granules, fibrin, &c., so that filtration is inefficient.

Though peripheral anterior synechia is undoubtedly the immediate cause of secondary glaucoma, it is itself produced by a variety of conditions, most of which are various forms of anterior or posterior synechia.

When an anterior synechia is formed the plane of the iris is advanced, so that the angle of the anterior chamber on this side is made more acute than normal. The cause which led to the formation of the synechia also causes iritis, usually of the traumatic type. Traumatic iritis is merely a plastic iritis due to injury. In it much exudation is poured out, possessing great tendency to organise. It collects in the diminished angle and becomes transformed into fibrous tissue, which welds the iris and corneo-sclera together, thus producing a peripheral anterior synechia, which may be strictly localised, so that no secondary glaucoma supervenes. Such eyes however are liable to fresh attacks of iritis, often of an insidious character. Each attack is followed by further occlusion of the angle, until finally the amount remaining open is insufficient to carry out efficient filtration and the tension rises.

The chief causes of secondary glaucoma are the following.

(1) *Perforation of the cornea with anterior synechia.*—The perforation may be due to an ordinary perforating wound of the cornea with incarceration of iris in the scar, or it may be due to a perforating corneal ulcer. The wound may be due to an operation, *e.g.*, extraction of cataract, for a peripheral section through the corneo-scleral margin or actually in the sclero-

tic near the margin has a similar effect to a wound in the cornea. The synechia need not necessarily be of iris, but after cataract extraction is often one of the lens capsule, which has the same effect of advancing the contiguous parts of the iris and obliterating the angle. Secondary glaucoma after operations may also be due to other causes (*vide* p. 458).

(2) *Annular posterior synechia* (*vide* p. 245).—This acts by interposing an impermeable barrier between the posterior and anterior chambers. The lymph secreted by the ciliary body is thus prevented from passing forwards into the anterior chamber. The iris becomes bowed forwards—bombé—and the periphery becomes apposed to the corneo-sclera, where it later becomes adherent. The aqueous is thus unable to escape from the eye and the tension rises. If the condition is not relieved by operation secondary glaucoma causes blindness. The prolonged high tension then causes degeneration of the ciliary processes, which cease to produce so much fluid, so that finally the tension may be normal or even sub-normal, and the eye may shrink. The condition is relieved by iridectomy, or if this is impossible, by iridotomy, communication between the posterior and anterior chambers being thus restored.

(3) *Wounds of the lens.*—When the lens is wounded it swells (*vide* p. 407), and pushes the iris forwards into contact with the corneo-sclera. Moreover the swollen lens matter in the aqueous impedes filtration through such part of the angle of the anterior chamber as remains open, both mechanically and also by increasing the albuminous content of the aqueous. Mere apposition is sufficient to produce permanent secondary glaucoma, which should at once be relieved by operation (*vide* p. 408). If it is not performed the iris becomes adherent to the corneo-sclera and the glaucoma becomes permanent, although the lens may be eventually absorbed.

(4) *Dislocation of the lens.*—This may be complete through the pupil into the anterior chamber. It then blocks the angle, especially if the iris is firmly contracted against its posterior surface. Partial lateral dislocation of the lens causes it to push forwards the iris on the side towards which it is dislocated. Since the circle of the equator of the lens is not much smaller than that of the angle a considerable portion of the latter is blocked, and secondary glaucoma supervenes.

(5) *Intraocular tumour.*—The mechanism whereby this produces secondary glaucoma will be described later (*vide* p. 387).

(6) *Intraocular hæmorrhage.*—Severe intra-vitreous or subchoroidal hæmorrhage forces forwards the vitreous and lens,

so that the iris is pushed into contact with the cornea. It also acts by filling the eye with highly albuminous fluid which filters with difficulty. If the vessel which has ruptured is large the tension may be raised to that of blood pressure.

A special type of glaucoma is sometimes met with after retinal hæmorrhage, which may be due to some unknown cause or to thrombosis of the central vein (*q.v.*). It is probably caused by mixture of the lymph with albuminous fluids. Such cases are sometimes grouped together under the designation *hæmor-*

Fig. 158.—Normal angle of anterior chamber.

rhagic glaucoma, a term which is however best avoided. They may be indistinguishable from primary glaucoma if not seen until the media are too opaque for ophthalmoscopic examination. Iridectomy is likely to be accompanied by severe hæmorrhage, and is therefore contra-indicated.

Primary Glaucoma.—The cause of primary glaucoma is unknown. The most probable theory is that of Priestley Smith, who attributes the preponderant *rôle* to the lens. It has already been pointed out (*vide* p. 9) that the lens continues to grow throughout life. The space between the equator of the lens and the ciliary processes, the circumlental space, will

therefore become smaller as the patient becomes older. If the eye is small the space may become so diminished that slight congestion of the ciliary processes may bring them in contact with the lens. The effect will be to prevent the fluid which is secreted by the ciliary body from passing forwards through the pupil. The lens will therefore be forced forwards, and will push the iris in front of it, making the anterior chamber very shallow, and bringing the periphery of the iris in contact with the corneo-sclera. In this manner the filtration angle will be



Fig. 159.—Peripheral anterior synechia (*S*), causing blockage of the filtration angle. *A*, cornea; *B*, canal of Schlemm; *E*, iris.

occluded and glaucoma will supervene. If the condition persists an acute attack of glaucoma is caused. If the occlusion of the angle is not complete a subacute attack is caused, and the spontaneous diminution of the congestion of the ciliary body may relieve the condition. The onset of fresh congestion leads to another attack, and thus chronic glaucoma is brought about. It will be seen from what follows that the facts fit in very well with this theory.

Primary glaucoma is essentially a disease of late adult or advanced life. It usually occurs after fifty years of age and is very common in Jews. It is sometimes hereditary, and in

these cases affords examples of "anticipation" (Nettleship), *i.e.*, it occurs at an earlier age in each succeeding generation. It is commonest in women, who are more liable to venous congestion in various parts of the body. Hypermetropic eyes are more susceptible than those with normal or myopic refraction; in fact, primary glaucoma is rare in myopic eyes. This has been shown to be due not to the hypermetropia *per se*, but to the smallness of the eyes. In order that an eye may be hypermetropic it is not necessary for it to be small, but as a matter of fact hypermetropic eyes usually are small. It has been found that the size of the cornea is a good criterion of the size of the ring formed by the ciliary processes, *i.e.*, of the circumlental space. The size of the lens varies little in different eyes of patients of the same age. It follows, therefore, that an eye with a small cornea will probably have a small circumlental space, and will be very liable to glaucoma.

As age advances the anterior chamber becomes shallower. If the cornea is small this will have the effect of diminishing the size of the angle of the anterior chamber. Filtration is carried out less easily when the meshes of the ligamentum pectinatum iridis are crowded together than when they are widely separated. Moreover the fibres of the ligamentum pectinatum tend to become thickened and sclerosed in elderly people. In these eyes a very slight further diminution of the angle may bring on an attack of glaucoma. Thus, the mere dilatation of the pupil with a mydriatic, by folding up the iris so that it is crowded into the angle, may suffice to occlude it entirely: *hence the extreme danger of instilling a mydriatic into the eyes of elderly people*, especially if they are hypermetropic or have small corneæ and shallow anterior chambers. Mere swelling of the lens in the early stage of cataract (*vide* p. 287) may induce glaucoma in eyes predisposed to the disease.

Since both eyes usually have a similar structure, glaucoma is likely to be bilateral, but one eye is often affected before the other.

The anatomical effects of pathological increased intraocular pressure are as follows. The already shallow anterior chamber is made still more shallow. In an early acute attack the periphery of the iris is merely apposed to the corneo-sclera. In the later stages and in chronic glaucoma of some standing it is firmly adherent. The longer the condition has lasted the firmer is the union. The iris is first bound down by organised exudate; later, the iris stroma atrophies and the inner wall of Schlemm's canal, which may be almost obliterated, is covered only by

degenerated retinal pigment epithelium. Anterior to this a "false" angle is formed where the iris leaves the cornea; no filtration can take place either through the peripheral anterior synechia or through the false angle.

The part of the eye which suffers earliest and most from the increased pressure is the head of the optic nerve. The lamina cribrosa, which is more resistant than the nerve tissue, is less resistant than the sclerotic. Hence it becomes pushed backwards, the nerve fibres being depressed also. The first manifestation of the effects of pressure is a bowing backwards of the connective tissue which forms the lamina cribrosa, so that it becomes concave anteriorly instead of passing straight across the porus opticus. This effect continually increases, until the lamina cribrosa is displaced backwards as a whole. Meanwhile the nerve fibres have been pressed together, so that the papilla becomes flat or depressed. The pressure causes the nerve fibres to atrophy, so that finally the lamina cribrosa is exposed upon the surface. In the final stage a deep cup is formed, generally having overhanging edges.



A B

FIG. 160. A, diagram of meridional section of normal disc; B, diagram of meridional section of glaucomatous cupped disc. Note the displacement backwards of the lamina cribrosa.

Pulsation of the arteries at the edge of the disc is often seen in glaucoma. While venous pulsation is of little importance spontaneous arterial pulsation is always pathological (*vide* p. 121). It is not always spontaneous in glaucoma if the tension is not very high, but even then it is induced by very slight pressure of the finger through the lid. The arterial pulsation is due to the increased pressure upon the walls of the vessels, so that the intravascular pressure is only able to force blood through at the height of the cardiac systole.

Other parts of the eye show less change. The pressure causes degeneration of the nerve fibre layer of the retina. The choroid becomes degenerated and thinned, only the larger vessels remaining. The ciliary body becomes degenerated in the last stages, after which the tension may cease to be raised owing to defective secretion of lymph.

The subjective effects of pathological increased intraocular pressure are manifold. Pain is complained of, due to stretching

of the sensory nerves of the eye. The patient sees coloured haloes round lights; these are due to alteration in the refractive conditions of the corneal lamellæ. The pupil becomes slightly dilated and immobile, owing probably to œdema and pressure on the ciliary nerves as they run through the choroid. Rapid diminution in the amplitude of accommodation may be a prominent feature, so that there is an apparent increase in presbyopia. It is attributable to pressure on the ciliary nerves and on the ciliary muscle. Diminution of vision is due to cloudiness of the media, retardation of the blood flow, and pressure on the nerve fibres in the retina and optic papilla.



Fig. 161.—Horizontal meridional section of a glaucomatous eye.

Cloudiness of the media affects the cornea principally, and is due in the early stages to altered refractive conditions, in the later to œdema. Pressure on the nerve fibres first affects the temporal side of the retina, and therefore the nasal side of the field. Later, the field becomes contracted in all directions, and central vision is depressed. Finally, vision is abolished, owing to total atrophy of all the fibres.

Acute Glaucoma.—In even the most acute cases of glaucoma careful inquiry will often elicit prodromal symptoms. Transient attacks of obscuration of vision, as if a cloud were in front of the eye, have occurred. Bright lights have appeared to be surrounded by rainbow haloes. A feeling of discomfort in the eye and neuralgic headache accompany these symptoms. Such attacks become more frequent, come on especially at night and after excitement or worry. It is noticed that stronger glasses are required for near work. This condition may extend over months or years.

If the patient is examined while the symptoms are present it will be found that there is a slight haziness of the cornea, so

that it looks bluish, like glass that has been breathed upon—hence the term glaucoma (γλαυκός, sea green). If the field of vision is taken between the attacks some slight contraction of the nasal field will be found, but central vision may be perfect.

The acute attack sets in suddenly. It may be due to some condition inducing venous congestion, *e.g.*, constipation, menstruation, over-feeding, alcohol, &c., or to worry, fatigue, a recent illness, or to the instillation of a mydriatic. The pain and anxiety of acute glaucoma in one eye may induce an attack in the other. Intense pain is felt in the eye and over the distribution of the fifth nerve. The pain is frequently so bad that it causes vomiting, and the attack is liable to be mistaken for a severe "bilious attack." The temperature may be raised. The constitutional disturbance is often so great that the patient is prostrated, the pulse becoming irregular and intermittent. The vision rapidly diminishes, so that in a few hours only hand movements can be recognised. In a considerable number of cases both eyes are affected almost simultaneously.

Objective examination shows some œdema of the lids and conjunctiva; the latter is intensely congested and looks dusky red, owing to the dilatation of the veins. Ciliary congestion is marked. The cornea is cloudy and insensitive to the touch. The anterior chamber is very shallow. The iris is discoloured, the pupil moderately dilated and oval, generally with the long axis vertical. The reactions to light and accommodation are abolished. Ophthalmoscopic examination is impossible owing to the cloudiness of the cornea. The tension of the eye is considerably raised.

There is no true inflammation in the early stages, so that the term inflammatory glaucoma, frequently used, is inadvisable; it should be replaced by *congestive glaucoma*.

If the condition is not relieved by operation, the amount of permanent diminution of vision depends upon the severity and duration of the acute attack. Total abolition of vision may result. More frequently improvement occurs, ushered in by diminution of pain. Considerable lowering of the visual acuity, and, still more, contraction of the field follows every acute attack. All grades, indeed, may be met with, from the mild prodromal attacks to the severest, with complete blindness. The tension remains permanently slightly elevated. Some congestion and irritability persist. The pupil reacts sluggishly, and the iris shows signs of atrophy, usually first in one or more sectors. Ophthalmoscopic examination now becomes possible. Cupping of the optic disc may or may not be found, according

to the duration of the raised tension before and after the acute attack. A single acute attack is not followed by cupping immediately, for this demands more or less prolonged high tension.

In every disease of one eye the other should be thoroughly examined. In acute glaucoma it may be found that chronic glaucoma has existed long unobserved in the other eye, and well marked cupping of the disc may be present. The same causes which induced the acute attack in one eye may rapidly induce a similar attack in the other. The pain and worry associated with preparations for operation and so on increase the danger. This eye should therefore be carefully watched, and prophylactic measures adopted. It is usually sufficient to instil a drop of 1 per cent. solution of eserin in the sound eye every day so as to keep the pupil contracted.

It is of the utmost importance that pathological cupping of the disc should always be recognised when present (Plate VIII., Fig. 2). It differs in ophthalmoscopic appearance from a deep physiological cup, with which it is most likely to be confounded, in that the excavation reaches to the edges of the disc and the sides are steep, not shelving. The retinal vessels have the appearance of being broken off at the margin of the disc. If they are accurately focussed here their continuations upon the floor of the cup are slightly out of focus and look broader and paler. When the edges overhang, as is often the case, the course of the vessels as they climb the sides of the cup is hidden. By the indirect method slight lateral movement of the large lens causes a distinct parallax (*vide* p. 111), which is more marked the deeper the cup. By the direct method the difference in level between the vessels at the edge and on the floor can be measured (*vide* p. 115).

There is always some atrophy of the optic nerve when the disc is cupped by the glaucomatous process ; it is therefore not surprising that there may be great difficulty in distinguishing a shallow glaucoma cup from the slight depression which follows simple atrophy of the nerve without increase of tension (*vide* p. 365). If the cup is deep and total it is certain to be glaucomatous. In shallow glaucomatous cups the disc has a pink colour, whereas the atrophic cup is white. In many early cases all the conditions have to be weighed carefully before it is possible to come to a definite conclusion ; the field of vision usually affords the most important criterion, the contraction being chiefly nasal in early glaucoma, concentric in optic atrophy.

The final stage of the untreated disease is *absolute glaucoma*.

The eye is completely blind. The anterior ciliary veins are dilated, and a reddish blue zone surrounds the cornea. The cornea is clear, but insensitive: it may have vesicles (bullous keratitis) or filaments (filamentary keratitis) upon it. The anterior chamber is very shallow. The iris is dilated, atrophic, and may have a broad zone of pigment around the pupil (ectropion of the uveal pigment). The pupil is grey or greenish, instead of jet black. The optic disc is deeply cupped. The tension is high; usually the eyeball is as hard as stone.



FIG. 162. —Diagram showing intercalary region.

Such an eye is generally painful, with temporary exacerbations, though patients often prefer to bear the pain rather than submit to excision. If it is still retained degenerative changes occur. The more important are due to giving way of the sclerotic before the continued high intraocular pressure. In this manner scleral staphylomata are produced. They may be in the neighbourhood of the ciliary body—ciliary staphylomata, or at the equator—equatorial staphylomata.



FIG. 163.—Diagram showing intercalary staphyloma.

Anatomical investigation shows that *ciliary staphylomata* are of two kinds. In one, the region where the iris is adherent to the corneo-sclera gives way (Fig. 162). These are called *intercalary staphylomata* (Fig. 163). In them the iris projects into the anterior chamber from an attachment at the anterior margin of the staphyloma, while the ciliary body, little altered, forms the posterior margin. The other form is the true *ciliary staphyloma* (Fig. 164). In them the region of the ciliary body itself gives way, so that it becomes spread out over the inner surface of the ectasia. In many cases both parts become ectatic.

Clinically it is impossible to distinguish between these forms.

Equatorial staphylomata can only be seen clinically when the eye is turned well to one side and the lids separated. The

thinning and bulging of the sclerotic occurs principally at the spots which are weakened by the perforation of the vortex veins and are unsupported by the recti muscles. Such globes may become enormous, with walls as thin as paper. There is considerable danger of rupture from slight injury.

Sooner or later the tension becomes normal or diminished in eyes with absolute glaucoma. This may be due either to stretching of the walls as already explained, or to degeneration of the ciliary body, whereby its secretory functions are diminished or abolished. Usually both factors play a part, varying according to the particular case. Such an eye may even shrink, but more commonly ulceration of the cornea occurs, owing to the defective resistance of the degenerated tissues. Hypopyon ulcer, panophthalmitis, phthisis bulbi then form the sequence of events.



Fig. 164.—Diagram showing true ciliary staphyloma.

Diagnosis.—Acute glaucoma is more likely to be mistaken for iritis than any other disease. The differential diagnosis has already been discussed (p. 246).

Treatment.—Acute glaucoma demands immediate energetic treatment. It is imperative that the tension shall be reduced as soon as possible. Theoretically this is best and most permanently effected by immediate iridectomy, and in most cases other measures fail. In any case they should not be dallied with too long. The moment is, however, an unfavourable one for operation. The conjunctiva is chemosed, the anterior chamber is extremely shallow, there is no time for exhaustive preparations, and a general anæsthetic will be necessary owing to the impermeability of the stretched cornea to cocain. Owing to the constitutional disturbance and the irregular action of the heart, one may feel diffident about giving a general anæsthetic in these cases. The danger is liable to be over-estimated, but a general anæsthetic may be replaced by deep infiltration of the orbit with novocain.

It is permissible to try other remedies for a short time first. Eserin (1 per cent.) should be instilled every five minutes for the first half hour, *one drop being placed in the other eye at the commencement* (*vide* p. 271); the eserin drops are then con-

tinued every half hour until the tension falls or operation is decided upon. Meanwhile the eye is continually bathed with water as hot as can be borne. This generally relieves some of the pain. If after an hour or two there is no sign of improvement, as indicated by distinct diminution of tension and good constriction of the pupil, iridectomy must be at once performed. Posterior sclerotomy may be employed as a temporary measure for the relief of tension while preparations for the iridectomy are being made (*vide* Chap. XXII.). The anterior chamber becomes deeper, thus facilitating the performance of iridectomy.

Even if the results are satisfactory iridectomy must be performed as soon as the eye is quiet. There is, indeed, one objection to non-operative treatment, even when successful, and to posterior sclerotomy, viz., that the patient may refuse the radical operation when the acute stage has passed off. In cases in which this is to be feared, it is advisable in the patient's interest to perform the operation at once. If there should be a lacrymal mucocele (*vide* p. 594) present both canaliculi should be ligatured, or the puncta cauterised with the actual cautery. If both eyes are affected, both should be operated upon at the same sitting. In no other disease is this procedure indicated or even justified.

The prognosis of iridectomy in acute glaucoma is so good, if it be performed early, that there is no valid reason for substituting any other operation for it. Some surgeons trephine for acute, as well as for chronic, glaucoma (*vide* p. 439). In my experience the technical difficulties of trephining under these circumstances are as great as, and the results are less favourable than those of iridectomy.

Iridectomy was first tried for glaucoma by von Graefe (1856) on the erroneous theory that as it lowered the tension of the normal eye (which is not true) it would be beneficial in glaucoma. The true explanation of the efficacy of iridectomy in glaucoma is that it reopens the angle of the anterior chamber and thus restores filtration. In acute glaucoma the iris is at first merely apposed to the corneo-sclera at the periphery. The manipulations draw it away. Part of the iris is then torn away at its ciliary attachment, and the angle of the anterior chamber is reopened in this situation. There is some reason to think that fluid is absorbed by the cut edges of the coloboma, for wounds of the iris do not heal by cicatrisation, the lymph-spaces of the stroma always remaining continuous with the anterior chamber at the cut margins.

In absolute glaucoma pain is best relieved by hot bathing and internal administration of aspirin. If possible consent should be obtained to excise the eye. It is rarely justifiable to trephine or perform any other operation merely for the relief of tension in these eyes, since there is nearly always a risk that the cause of the glaucoma may be an intraocular malignant growth, usually sarcoma of the choroid.

Chronic Glaucoma, sometimes called simple glaucoma, is fundamentally the same disease as acute primary glaucoma: every grade of severity is met with, but the more chronic



FIG. 165.—Field of vision in commencing glaucoma, showing contraction of nasal side.

forms are so insidious that special attention must be directed towards their discovery.

The patient usually complains of transient attacks of obscuration of sight, and of gradually diminishing acuity of vision; but in some cases there may be no history of haloes and the diminution of vision is continuous and very insidious.

The eyes may appear perfectly normal at the first examination, though sometimes the anterior ciliary veins are congested, and the pupil is somewhat dilated and sluggish. An abnormally small cornea should draw attention to the possibility of glaucoma, and hypermetropia increases the probability. The tension may be quite normal, and is found to be elevated only during an attack of cloudy vision. Hence it may be necessary

to examine the patient frequently and at various times in the day, especially during such attacks.

Subjective examination will often reveal no diminution of central vision. Hence it is of the utmost importance to take a careful chart of the field of vision. The commonest change is partial loss of the nasal field, often accompanied by some general contraction (Fig. 165). The change in the nasal field may be regular in outline, or there may be indentations with the apex directed towards the fixation spot. Such sectional defects may be above or below. In later stages the general contraction is



Fig. 166.—Normal fields with tests as follows:—10/300, 6/2000, 3/2000 (*vide* p. 135). (A. H. H. Sinclair.)

more marked, and eventually only a paracentral patch of the temporal field persists, central vision being abolished. Partial scotomata are to be found by special means before nasal constriction occurs, or when it is only slightly developed. If the central area of the field is mapped out on a Bjerrum's screen (*vide* p. 137), or with a suitable scotometer, an area of relative defect can frequently be traced in direct continuity with the blind spot (*Bjerrum's scotoma*). The scotoma may pass in an arc from the blind spot above or below the fixation point, or may form a complete annular scotoma. It is due to injury of bundles of nerve fibres at or near the edge of the disc. The destruction of these fibres is also said to account for a charac-

teristic sharply defined horizontal edge to the lost portion of the field on the nasal side (*Ronne's step*) (Figs. 165, 167). The earliest sign of all is said to be a sickle-shaped extension of the blind spot above or below, or both, with the concavity of the sickle directed towards the fixation point (*Seidel's sign*) ; this is of more doubtful significance. A relative central scotoma sometimes follows rapidly on the development of the paracentral scotomata : the prognosis is worse in these cases.

Retinal adaptation is usually slower in people with chronic



Fig. 167.—Glaucoma fields with 3/250 and 1/2000. A = Position of blind spot. (A. H. H. Sinclair.)

glaucoma, so that they take longer to get used to the lower degree of illumination on passing into a dimly lighted room.

Ophthalmoscopic examination will often show some cupping of the disc ; frequently it is far advanced, though the symptoms have been so slight as to have passed almost unnoticed. The field in these cases will be found to be much damaged.

Primary glaucoma invariably attacks both eyes sooner or later ; usually one is considerably more advanced than the other. The chronic form sometimes occurs in young people, and seems to attack men almost as frequently as women. It also occasionally occurs in myopic eyes.

There is no evidence that chronic glaucoma is due to increased blood pressure, though, so frequently occurring in

elderly patients, it is often associated with it. Although the intraocular pressure responds passively to rapid changes in the general blood pressure (*vide* p. 15), slow changes are compensated and a fresh equilibrium of secretion and excretion is established.

Diagnosis.—Chronic glaucoma has frequently been mistaken for cataract or optic atrophy. Cases occur in which the gradual loss of vision is attributed to cataract, and the patient is told that nothing can be done until the cataract is ripe for operation. Vision is thus irretrievably lost. As a rule diagnosis is very easy. The haze of the pupil is bluish, diffuse and uniform, unlike the usual appearance in cataract. The pupil may be slightly dilated and generally reacts less to the stimulation of light. Examination with the ophthalmoscopic mirror in most cases renders the diagnosis certain by showing a uniform red reflex and the absence of opacities in the lens.

Doubt as to the presence of glaucoma may arise in cases with senile striæ in the lens. In patients predisposed to glaucoma the swelling of the lens which occurs in the early stages of cataract (*vide infra*) may lead to increase of the intraocular tension. The disc is not usually cupped in these cases, but there is generally more failure of vision than is accounted for by the lenticular opacity, and the field of vision may show contraction on the nasal side; the reaction of the pupil to light may be sluggish. In such cases it is advisable to do a preliminary iridectomy, which should be of the glaucoma type with a large and peripheral coloboma.

Cupping of the disc in glaucoma is accompanied by atrophy of the nerve fibres, and it may be difficult to distinguish this atrophy from optic atrophy due to other causes, especially those giving rise to what is known as "primary" optic atrophy. The latter condition shows some depression of the surface of the disc which is usually too slight to be actually measured by the ophthalmoscope, but is demonstrated by the bending of the vessels as they pass over the edge of the disc. The depression is greater in most cases of chronic glaucoma. In cases which give rise to difficulty in diagnosis little aid is afforded by the tension of the eye, which may be normal or inappreciably raised at the time of examination. Reliance must be placed upon the history of the case, the condition of the cornea and anterior chamber, and the record of the field of vision. The latter shows concentric constriction in primary optic atrophy, more marked contraction of the nasal side in glaucoma. Accurate mapping out of the blind spot and the central region of the

field on a large scale reveals changes in glaucoma (*vide supra*) which do not occur in optic atrophy.

Treatment.—An operation should be performed as soon as the condition is definitely diagnosed, the chief criterion in doubtful cases being the condition of the field of vision. If there is contraction of the nasal side the operation should be performed at once, even if central vision is quite normal.

Until a few years ago iridectomy was the operation invariably performed for the relief of chronic glaucoma. The prognosis of iridectomy, however, is not nearly so good as in the acute form, owing to the fact that the periphery of the iris is often firmly adherent to the corneo-sclera before the condition is diagnosed. Hence a special endeavour was made to open up the occluded angle. If the section is made at the apparent corneo-scleral margin, when the iris is torn away it is almost certain to tear at the false angle, and little or no good results. It was customary, therefore, to make the section as peripheral as is consistent with safety to the ciliary body, *i.e.*, 2 mm. behind the corneo-scleral margin, with the object of carrying the incision through the adherent part of the iris. It must be confessed that this object was seldom if ever attained.

It was found, however, that this method of performing iridectomy sometimes succeeded in spite of impossibility of restoring the normal method of filtration. It effected this by establishing a *filtering scar*. The new scar in these cases is composed of spongy tissue, through the interstices of which the intraocular fluid is able to make its way into the subconjunctival tissue, where it is absorbed.

Deliberate attempts have therefore been made to establish a safe filtering scar for the relief of chronic glaucoma. Such a scar is usually formed only if there is some impediment to proper cicatrisation, as, for example, when the iris is incarcerated in the wound. It is not justifiable deliberately to leave a knuckle of iris in the wound when doing an iridectomy, as has been advocated. Such a procedure involves the grave risks of iridocyclitis, secondary infection of the eye, and sympathetic ophthalmia. Even more unjustifiable is the insertion of a silk thread or other drain into the wound.

In an ordinary corneo-scleral section the lips of the wound are in good apposition and sound healing rapidly takes place. This is much less likely to occur if there is a gap beween the lips of the wound. Under these circumstances the gap becomes filled with loose scar tissue and a filtering cicatrix may result. Various operations have been based upon this principle. In

Lagrange's operation an ordinary iridectomy is performed, but before closing the wound a small piece of the anterior lip is snipped off without wounding the conjunctival flap. In *Herbert's operation* a small rectangular trap-door is cut in the sclerotic just outside the limbus, the hinge being towards the cornea. A circular wound in the sclerotic offers the best chance of success theoretically. This can be effected by means of a punch or a trephine.

Trephining is the operation which is now generally performed for chronic glaucoma (*vide* p. 439). By it a disc 1·5 mm. in diameter is removed from the wall of the globe just inside the limbus, so that part lies in the cornea and part in the sclerotic. The knuckle of iris which prolapses into the wound should be excised, so that a small peripheral iridectomy is performed. If a larger disc is removed there is danger of the tension becoming permanently too low, with the risk of malnutrition of the eye. The operation not infrequently fails from blockage of the wound with iris or too dense scar tissue. Filtration is encouraged by gentle massage of the eyeball through the upper lid. Secondary infection of the eye is liable to occur from injury of the epithelium covering the bulging conjunctiva, but it is comparatively rare and is a risk which may be taken justifiably in treating so grave a disease as glaucoma. If trephining fails it can be repeated at some other part of the limbus, but the wound should always, if possible, be placed where it is covered and supported by the lid, though it should not be too close to the previous site.

Trephining is uncertain in its results and is liable to be complicated with serious dangers, both immediate and remote, yet in my opinion it affords the best means on the whole yet devised for dealing with chronic glaucoma.

In very advanced cases the field of vision may be found reduced almost to the fixation point. Experience teaches that in these cases operation may not only do no good, but may result in the sudden complete loss of all vision. After explaining this risk to the patient it is still advisable in most cases to perform the operation, because this unfortunate result is rare, and in any case the eye is likely to become painful unless the tension is relieved.

Since glaucoma always occurs in the other eye sooner or later the question of a prophylactic operation in this eye arises. Since it is attended by some danger, both immediate and remote, and since the advent of glaucoma may be long delayed, it is inadmissible to operate until some slight contraction of the

nasal field can be demonstrated. A broad iridectomy performed at this early stage has a much better prognosis than when done later. It has the disadvantage of slightly lowering the vision by producing some corneal astigmatism, whereas trephining does not usually produce this result. The greatest care must be taken to warn the patient of the danger of the disease attacking the other eye, and of the earliest symptoms. He should be examined thoroughly and the field of vision taken every three months, and he should be instructed at once to consult the surgeon if any signs of the disease occur. Weak pilocarpin or eserin drops may be used every other day as a prophylactic measure, and the general régime should be ordered so as to avoid cerebral congestion. He should also be warned against putting drops, lotions, or ointment of any kind into his eyes without the advice of an ophthalmic surgeon.



Fig. 168.—Diagram of eye with infantile glaucoma (buphthalmia). Note the stretching of the corneo-sclera at the periphery, the flattening and displacement backwards of the lens, the cupping of the disc, and the general enlargement of the globe.

When one eye has been almost or quite lost at the time of the first visit it becomes a serious question whether the better eye should not be operated upon rather than the worse. Many such difficulties arise in the treatment of glaucoma, and can only be decided by the conditions of the individual case.

It has been stated that operation is unconditionally indicated when chronic glaucoma is definitely diagnosed. Cases arise in which it is impossible to be certain that the disease is present, and others in which operation is refused. In such cases the pupils should be kept contracted by miotics. Eserin or pilocarpin often succeed in reducing the tension of chronic glaucoma and postponing the evil day when useful vision is lost. *They never cure chronic glaucoma.* Hence they must be adopted only as a temporary means of alleviation or as useful adjuncts. If the tension is definitely raised and operation is refused, 1 per cent. eserin should be ordered three times a day until the tension is normal. If the tension is not definitely raised the pupils should be kept moderately constricted with 0·5 per cent. pilocarpin or 0·25 per cent.

eserin once or twice a day for a prolonged period. These cause conjunctival and ciliary irritation if used constantly, but pilocarpin is less irritating than eserin, though perhaps rather less efficient. Gentle massage with the finger-tips is certainly useful, promoting lymph-flow, and temporarily reducing the tension.

Infantile Glaucoma (*Syns.—Buphthalmia, Hydrophthalmia*). —Glaucoma in rare cases attacks children, when it assumes a quite different clinical appearance. It arises from congenital blockage of the angle of the anterior chamber, due either to a congenital defect whereby the root of the iris does not become normally separated from the corneo-sclera or becomes adherent to it through intra-uterine or infantile inflammation. In many cases examined microscopically Schlemm's canal has been found defective or absent. The fundamental condition is therefore the same as in glaucoma in adults, viz., defective filtration of lymph from the eye. The reason why it assumes so different a clinical picture is dependent entirely upon the greater plasticity and extensibility of the walls of the young eye. Instead of offering an insuperable resistance to increased internal pressure the sclerotic gives way more or less uniformly, so that the globe becomes very large.

The thinned sclerotic of the ciliary region is bluish in colour, owing to the uveal pigment showing through. The junction of the cornea and sclerotic also stretches, so that the cornea is forced forwards and assumes a globular shape (keratoglobus). The anterior chamber is therefore extremely deep (Fig. 168). There are often slight opacities in the cornea, some appearing as lines with double contour; these are due to ruptures in Descemet's membrane. The lens does not participate in the general enlargement; owing to the expansion of the ciliary region the suspensory ligament is stretched so that the lens is flattened and displaced slightly backwards. This removes some support to the iris, which becomes tremulous (iridodonesis). The optic disc is deeply cupped if the condition has lasted long.

The intraocular tension is raised, but often scarcely appreciably as determined by clinical methods owing to the expansion of the globe. This fact for a long period prevented the true pathology of the disease from being recognised.

As a result of the expansion the eyes are usually myopic, though less than might be anticipated from their length. This is due to the flattening of the lens and its displacement backwards, as well as to some flattening of the cornea, all of which factors tend to counteract the axial myopia.

Treatment is unsatisfactory. Iridectomy and anterior sclerotomy have been employed with success, but more frequently with failure. The best results have been obtained from anterior sclerectomy with the trephine, and this method is also less dangerous than either iridectomy or anterior sclerotomy. Miotics are useless.

CHAPTER XV

THE LENS

The lens is composed entirely of epithelium, which is surrounded by a cuticular envelope or capsule (*vide* p. 9). It is therefore, subject only to degenerative changes, and is incapable of becoming inflamed. Degenerative changes in the lens invariably result in loss of transparency in the parts affected. This condition of partial or complete opacification is called *cataract*.

The degenerative changes commence by the accumulation of fluid between the lens fibres; so that spindle-shaped spaces are formed. They, in the first instance, cause opacity merely by the irregular refraction which they induce. Later, the fluid contains albuminous deposits which are actually opaque: they form globular masses, called Morgagnian globules. At a later stage the fibres break down into rounded masses which are indistinguishable from the Morgagnian globules.

Cataracts are classified according to the position and extent of the opacity or opacities in the lens, and it is found that the situation and distribution correspond with various combinations of clinical conditions—age, general disease, &c. In some cataracts the opacities spread and fuse until the whole lens becomes opaque; such are called progressive cataracts: in others they remain stationary. Senile cataract, which is the commonest form of all, is a progressive cataract.

Senile Cataract.—This, as its name implies, rarely occurs in persons under fifty years of age.

In the *incipient* senile cataract radial spokes or sectors of opacity are seen, with clear areas between them (Figs. 169—170). They are difficult to see in daylight or by oblique illumination (*vide* p. 91), and cataract should not be diagnosed without confirmation with the ophthalmoscope. With the undilated pupil only the ends of the spokes are seen, but when the pupil is dilated with cocain or homatropin (*never* with atropin, *vide* p. 268), the linear opacities are often found to be the apices of sectors, with their bases towards the periphery. They generally begin in the lower part of the lens. Careful

examination with oblique illumination and the ophthalmoscope will show that the opacities are in the superficial parts or cortex of the lens, some in front of the nucleus, others behind. They start from the region of the equator and

Fig. 169.—Senile cataract. 1, section, showing opacities in the cortex; 2, appearance by reflected light—dark striæ on a red background; 3, appearance by oblique illumination—grey striæ on a dark background. (Nettleship.)

extend towards the axis of the eye, more and more spokes and sectors developing as time goes on.

Seen by oblique illumination the opacities are grey (*vide* p. 91); seen with the ophthalmoscopic mirror at reading distance they appear black against a red background. At the very earliest stage the opacities shift with the incidence of the light, showing that they are merely differences of refractive index (*vide* p. 285).



Fig. 170.—Commencing senile cataract, the striæ confined to the lower part of the lens, a very common mode of commencement.

The pupil in old people is seldom so black as in the young, and is sometimes distinctly grey. If the greyness is uniform, cataract should not be diagnosed unless definite opacity is shown on examination with the ophthalmoscope. This greyness without opacity is caused by increase in the refractive index of the cortex of the lens in old people (*vide* p. 49), and is due to increase of reflection and scattering of light.

The above description applies to the commonest arrangement of the opacities in senile cataract; it may be distinguished as the *subcapsular* type. Two other types occur less frequently. In one group the opacities are *supranuclear*. They are more variable in appearance and consist of concentric lines, radial streaks and cloudy patches. This irregularity distinguishes them from lamellar cataract (*q.v.*). In the third group the opacities are *intranuclear*, by which term these cases may be distinguished from the congenital nuclear cataract. In this group the nucleus of the lens is diffusely

cloudy, gradually clearing towards the peripheral cortex. Combinations of the various types are not uncommon, subcapsular sectors being associated with intranuclear cloudiness, or the latter with supranuclear opacities. Moreover dots of opacity occur. They are larger and denser than the dots of congenital punctate cataract, and are unlike that form in being slowly progressive. Punctate senile cataract may appear as early as twenty to thirty years of age. The intra-



FIG. 171.—A, shadow of iris by oblique illumination in immature cataract; B, absence of shadow in mature cataract.

nuclear form is also earlier (about forty) than the ordinary subcapsular.

In the progressive stage of senile cataract the lens contains more water than normal, and this is associated with swelling of the fibres. When the opacity has become considerable the swelling causes an appreciable *intumescence* of the whole lens, shown by the increasing shallowness of the anterior chamber. This may be the cause of increased intraocular pressure in eyes predisposed to glaucoma (*vide* p. 268).

The nucleus in the common subcapsular type undergoes little change and does not become opaque, but eventually the

whole of the cortex is cataractous. The cataract is then said to be ripe or *mature*. The whole of the pupillary area may appear to be opaque before the cataract is mature, since the most superficial layers of the cortex are the last to degenerate. As long as there is any clear lens substance between the pupillary margin of the iris and the opacity the iris throws a shadow upon the grey opacity when light is cast upon the eye from one side (Fig. 171, A). When the cortex is completely opaque the pupillary margin lies almost in contact with the opacity, separated only by the capsule; the iris then throws no shadow, and the cataract is known to be mature (Fig. 171, B). This is an important guide to the most favourable time for operation.



Fig. 172. Diagram of Morgagnian cataract. N, nucleus of lens.

At this stage it will be found that the anterior chamber has regained its normal depth. The watery fluid has been absorbed from the lens, which has again returned to its normal volume.

If the process is allowed to go on uninterrupted the stage of *hypermaturity* sets in. The cortex is then completely disintegrated and transformed into a pultaceous mass. Usually the loss of water continues so that the lens becomes more and more inspissated and shrunken. The lens is then flat and yellow, often with cretaceous deposits and bright specks due to crystals of cholesterin. The anterior capsule becomes thickened by proliferation of the anterior cubical cells, so that a dense white capsular cataract is formed at the anterior pole in the pupillary area. Owing to shrinkage the lens and iris become tremulous, the anterior chamber being much deepened. Degeneration of the suspensory ligament may lead to luxation of the lens.

Sometimes the absorption of water ceases at the stage of maturity. The cortex then becomes quite fluid, and the nucleus sinks to the bottom of the capsule. The liquefied cortex is milky, the nucleus appearing as a brown shading, limited above by a semicircular line. The nucleus alters its position with changes in position of the head. Such a cataract is called a *Morgagnian cataract* (Fig. 172).

In rare cases the sclerosis of the lens fibres which causes the development of the nucleus extends beyond the usual limits.

so that the whole of the lens becomes converted into nucleus. Such a lens is hard, dark brown, and semi-transparent. The pupil looks black, the brownish colour being revealed only by oblique illumination. The condition is called *black cataract*, though strictly speaking it is not a true cataract. Central intra-nuclear senile cataracts often show a brownish colour by oblique illumination, due to coincident sclerosis.

The mode of treatment of cataract depends largely upon the amount of central sclerosis, *i.e.*, upon the size of the nucleus. Up to about thirty years of age the nuclear fibres are still fairly soft, and capable of becoming absorbed if the aqueous gains access to them. After this age absorption is very slow and incomplete, and if cataract occurs the nucleus must be removed from the eye. The size of the nucleus then determines the size of the incision which is necessary. The cases in which the nucleus is very small are called *soft cataracts*, since they consist chiefly of soft cortical matter. In most patients over fifty the nucleus is large, and these cataracts are called *hard cataracts*, although the lens is by no means hard throughout. This only occurs in black cataract, in which the nucleus reaches its maximum size, viz., that of the whole lens. Such cataracts require a very large section for their removal. In mature cataracts the brownish appearance of the nucleus by oblique illumination gives some idea of its size and an indication of the size of the section necessary for its removal.

Symptoms.—The appearance of black spots in front of the eyes is usually the first symptom complained of. They differ from the ordinary muscæ volitantes occasionally complained of in normal eyes, and much exaggerated in cyclitis, &c., in that they are stationary, retaining their relative position in the field of vision in different positions of the eye. Uniocular polyopia, another symptom, is the doubling, trebling, &c., of the objects seen with the eye. It is due to the irregular refraction of the degenerating lens, so that several images are formed of each object. It is often worst on looking at bright lights, and is therefore noticed most in the evening.

As the opacity extends and becomes denser, the acuity of central vision suffers, especially when there is much central opacity. In the latter cases vision is often better in a dull light, owing to the dilatation of the pupil. In most cases of senile cataract the pupillary region suffers latest, so that a bright light is grateful to the patient, both on account of the better illumination and also because the rays which pass through the irregularly refracting peripheral parts of

the lens are cut off, but the patients seldom like to face the light.

Eventually the central area becomes affected and vision steadily diminishes until only quantitative perception of light remains. In many cases of mature senile cataract fingers can still be counted at a few feet, or at least hand movements discerned. In all cases light should be perceived readily and the direction of incidence accurately indicated. The detection of the *projection of light* is of the utmost importance, as it affords important evidence as to the probabilities of a good result from operation. It is tested as follows. The opposite eye is covered securely by the palm of the patient's hand. Light is then reflected from the ophthalmoscopic mirror into the cataractous eye from various directions, the patient looking straight forwards. He is told to point with his other hand in the direction from which the light seems to come. He ought to do this readily and accurately. If he does not, we suspect some disease of the retina, *e.g.*, patches of retino-choroidal atrophy, &c., and a less favourable prognosis is given. Relatively poor projection is not an absolute contraindication to operation, and each case must be determined on its merits.

It will be seen that cases of advanced cataract in which the fundus cannot be satisfactorily explored by the ophthalmoscope at the first visit, and in which projection is relatively bad, offer considerable difficulties in deciding the treatment to be adopted. It is therefore of the greatest importance that every case of incipient cataract should be most carefully explored and exhaustive notes of the ophthalmoscopic condition taken, so that at a later stage, when the fundus can no longer be observed, its previous condition is already on record. Every case of incipient cataract should therefore have the pupil dilated so that a thorough examination of the eye may be recorded. Homatropin may be used with impunity in most cases, but a drop of 1 per cent. eserin must invariably be instilled before the patient leaves, otherwise there is danger that an attack of glaucoma may be induced by the administration of the mydriatic. If the eye is definitely of glaucomatous type, with small cornea, shallow anterior chamber, &c., cocain should be used instead of homatropin. The field of vision should also be taken at this stage.

The rate of development of senile cataract varies greatly, sometimes occupying many years, or, indeed, the cataract may never reach maturity. The progress is usually more rapid in very old people. Very rapid maturation in younger patients

usually indicates some complication, *e.g.*, cyclitis, diabetes, &c. The forms with fine radial lines are slower than those with cloudy opacities. It is best to examine every case at stated intervals, a careful drawing of the opacities being recorded at each visit. The length of the intervals must be determined by the individual case.

Cataract occurs equally in men and women. It is usually bilateral, but develops earlier in one eye than the other. Cases of hereditary predisposition have been recorded, and in some of these the cataract develops at an earlier age in successive generations ("anticipation," Nettleship). Cataract can be produced in rabbits by administration of naphthalin and other poisons. It occurs in cases of ergotism in human subjects, and autotoxins may account for its occurrence in tetany and myotonia atrophica.

Pathology.—The cubical cells lining the anterior capsule of the lens (*vide* p. 9) undergo vacuolation in senile cataract. Changes in these cells can be made out clinically by using a strong binocular loupe. The cortical opacities are due to the formation of Morgagnian globules and the breaking up of the lens fibres (*vide* p. 285). The hydrolysis of the lens protein leads to the formation of tyrosin both in the lens and aqueous: tyrosin is not found in the normal lens (Burdon Cooper). Cholesterin crystals are not infrequently seen in cataracts. Senile cataract has been attributed to changes in the cortex produced by shrinkage of the nucleus, but the evidence is unsatisfactory. Many other theories have been advanced, but the most probable is that the nutrition of the lens suffers owing to metabolic changes in the general organism, possibly owing to the formation or want of destruction of specific autocytotoxins.

Treatment.—No treatment by drugs, &c., has any effect upon the progress of uncomplicated senile cataract.

In *incipient cataract* the condition of the patient may be much ameliorated during the tedious process of maturation. A low degree of myopia (1 D to 4 D) may develop during this stage; it is due to relative increase in the index of refraction of the nucleus of the lens, a change in the opposite direction to that which usually occurs (*vide* p. 49). Astigmatism may develop or undergo change. These errors of refraction should be corrected; but often the astigmatism is irregular and glasses afford little help. Considerable loss of vision may be associated with the refractive changes which precede the definite formation of opacities. Tinted glasses may be found beneficial, the tint varying with the circumstances of the case.

Amber tinted glasses are most generally useful. They cause an extraordinary increase of definition even in normal persons, due to absorption of rays of short wave-length. Blue glasses, which allow the chemically active ultra-violet rays to pass, are not contraindicated in this case, since the cutting off of the more luminous rays is restful. These or smoked glasses are indicated, especially when there is a considerable degree of central opacity, since the pupils are kept slightly dilated. This effect may be obtained with greater certainty by instilling a very weak mydriatic. Atropin, $\frac{1}{16}$ to $\frac{1}{8}$ gr. to $\overline{3}$ i., one drop every other day, may be ordered, if homatropin is found not to raise the tension. The slightest predisposition to glaucoma, *e.g.*, high hypermetropia, small cornea, very shallow anterior chamber, &c., contraindicates this treatment, and it is wise to observe the tension carefully during the treatment in all cases. Sometimes weak atropin causes more blurring, in which case it must be abandoned. Central opacities often cause diminution of central vision apparently out of proportion to the amount of opacity observed.

There is no reason to restrict the use of the eyes in incipient uncomplicated senile cataract, but the patient may be much assisted by instructions as to the arrangement of illumination and so on. If the pupillary area is free brilliant illumination will be found best; if the opacities are largely central, a dull light beside and slightly behind the patient's head will give the best result.

In *mature cataract* the lens must be extracted. Before deciding to operate, attention must be paid to details other than those connected with vision, previously described. The pupil should react promptly and normally to light. Careful search must be made for precipitates on the back of the cornea ("k.p."), for the cataract may be a mature complicated cataract (*vide* p. 302). The urine must be tested to eliminate albuminuria and glycosuria, though these do not necessarily contraindicate operation. The state of the conjunctival sac must be thoroughly examined. If there is the slightest conjunctivitis, and above all, if there is dacryocystitis, a course of preliminary treatment is necessary. Old people frequently suffer from chronic conjunctivitis induced by senile ectropion, &c. It is best treated by relieving the cause as far as possible, and the use of astringent lotions. An occasional painting with silver nitrate is the most potent means which we possess of removing infective organisms from the conjunctival sac in these cases, since they are carried away mechanically with the

desquamated epithelium, &c. Often the process is long and tedious, and doubt as to the safety of operating still persists. In such cases a bacteriological examination should be made, and if pathogenic organisms are found, especially virulent pneumococci, operation must still be postponed. In cases of doubt the eye may be tied up with a pad and bandage for one night. The pad is examined the following morning, and if there is any discharge upon it, or if the lids are gummed together with inspissated secretion, further treatment and repeated bacteriological examinations are indicated. It is inadvisable to tie up the eye the night previous to operation, for it is found that this prodecure favours the growth of bacteria in the conjunctival sac.

The presence of a mucocele is an absolute contraindication to operation. It must be cured (Chap. XXXII.), or the lacrymal sac must be excised, or the canaliculi must be temporarily obliterated. The last may be effected by tying a ligature round each canaliculus or by cauterising each punctum with the actual cautery. The best treatment is usually excision of the sac.

The teeth, nose, and throat, and any other likely focus of sepsis should be examined, and any pyorrhœa, &c., eliminated before the cataract operation is undertaken.

The treatment of *unilateral* and of *immature cataract* offers some difficulty (cf. p. 301). When the cataract is mature in one eye while the other retains good vision little advantage is gained by operating upon the cataract. The difference in refraction between the two eyes after operation will be so great that it will be impossible for the patient to see well if the refraction is corrected, and if uncorrected the large blurred images formed by the eye may be a positive disadvantage. The sole advantage which is gained is an increase of the field of vision on the affected side. This may be a matter of great importance, as in people who work amidst machinery or have to go about where there is much traffic ; in these exceptional cases extraction is indicated. There is also the slight advantage that the eye is prepared for the time when vision fails in the less affected eye, but this may be long delayed. These slight advantages do not as a rule justify operation, which, it must be remembered, is attended with some, if usually trivial, danger, not only to the eye operated upon, but also to the other eye (*vide* p. 425). On the other hand, the cataract must not be allowed to progress to too advanced a condition of hypermaturity. Operation is then more difficult and more dangerous.

The case should be watched, and if signs of thickening of the capsule, calcareous deposits, &c., appear, extraction should be performed even though the vision in the other eye is still good.

Cases of immature cataract with loss of useful vision require even more skill in the determination of the best time for operation. The difficulties and dangers of extraction are undoubtedly increased by operating while there is still a large amount of clear soft cortex. It is difficult to remove from the eye, tends to the production of iritis and other complications, and leads to the formation of dense secondary cataract (*vide* p. 301). Immaturity, however, is not an absolute bar to operation; extraction under these circumstances may be followed by perfect results. It must be remembered that the patients are old, and, if not operated upon, are doomed to practical blindness, which in the lower classes entails the loss of all wage-earning capacity. Operation will probably be attended by at least the recovery of useful vision, whereby the con-

Fig. 173.—Lamellar cataract. 1, 2, 3, as in Fig. 169. (Nettleship.)

ditions of existence are much ameliorated. It is not necessary, therefore, to wait indefinitely for complete maturity if useful vision has already been lost.

Some surgeons temporise in these cases, performing a preliminary iridectomy, on the grounds that the operation accelerates the ripening of the cataract. It certainly occurs in rare cases, but is by no means constant. Preliminary iridectomy has the advantage of facilitating the subsequent extraction, but has the considerable disadvantage of subjecting the patient to the discomforts and dangers of two operations instead of one. It may be employed in complicated cases in which it is desired to test the reaction of the eye to operative interference, for it is a less severe operation than extraction, and will afford indications as to the advisability of further procedures. It is also to be advocated in cases with much diffuse opacity, often more concentrated in the central part of the posterior cortex, for in them the diminution in visual acuity is very marked, maturation is indefinitely delayed and the complications produced by a large amount of sticky cortex

are much to be feared. The extraction of the cataract should not be performed until at least four weeks after the preliminary iridectomy.

The correction of the refraction after extraction of cataract is dealt with elsewhere (*see* Aphakia, p. 483).

Fig. 174.—Lamellar cataract.

Cataracts of Congenital or Infantile Origin.—These are almost always partial and stationary. The commonest forms are lamellar and anterior capsular; less common are the various forms of congenital cataract, mostly of lamellar type, sometimes central or total.

Lamellar Cataract (*Syns.*—*Zonular Perinuclear Cataract*).—This usually occurs so early in infancy that it is doubtful if it is not congenital. Although there is no true nucleus at this early age the central parts of the lens are conveniently termed the nucleus. The opacity in lamellar cataract is situated in the layers surrounding this central core, which itself usually contains punctate opacities; the superficial cortex is quite clear (Figs. 173—175). When the pupil is dilated a grey discoid opacity is seen, surrounded by a perfectly transparent marginal area. The diameter of the disc varies, that of the clear peripheral area varying inversely. With the mirror the disc appears black and sharply defined at the outer edge, diminishing in density towards the centre; the peripheral area shows a normal red reflex. Along the outer edge spokes of opacity, resembling the handles of a steering wheel, often extend slightly into the clear area (Fig. 174). They are called riders, and are due to spindle-shaped opacities between the lens fibres in layers a short distance outside the main opacity. Occasionally two concentric rings of opacity are seen. The cataract is usually stationary, but cases occur in which total opacity gradually develops. Both eyes are



Fig. 175.—Lamellar cataract with very slight opacity. $V = \frac{6}{12}$.

almost always affected, though not always to the same degree.

The opacity is always sufficiently large to fill the area of the undilated pupil. The diminution of vision is therefore entirely dependent upon the density. The patient is brought for examination on account of defective vision. He usually holds objects very close to the eyes and is thought to be "short sighted." Myopia is indeed not uncommon in these cases, but the approximation of objects is usually for the purpose of obtaining larger retinal images (*vide* p. 34).

There is no doubt that lamellar cataract is due to a period of malnutrition at some stage of late intra-uterine or early infantile life. Epithelial structures are most affected. Epithelium as a rule covers surfaces, and the oldest cells are cast off, being replaced by young cells derived from the basal layer. There are two sites in which the epithelium persists: (1) the lens, where, owing to its formation as an invagination of epiblast, the oldest cells are central and cannot be cast off; (2) the enamel of the teeth, where the cells become calcified and thus retained. Hence lamellar cataract is almost invariably accompanied by defective enamel in certain of the permanent teeth. The hypoplasia differs essentially from the condition of the teeth in congenital syphilis. The teeth have an eroded appearance, with transverse lines across them, the incisors and canines being most affected (Fig. 179).

FIG. 176. — Hypoplasia of teeth.

The time at which the pathological process took place is indicated by the size of the diameter of the opacity and the particular teeth affected with hypoplasia. Only those teeth are affected whose enamel germs are being formed at the time. As regards the lens the youngest fibres are the most superficial (*vide* p. 9), so that the diameter of the opacity indicates the size of the lens at the time. Concentric rings of opacity are accounted for by successive periods of malnutrition.

The cause of the malnutrition is probably to be found in errors of feeding and possibly exanthemata. There is some reason to think that rickets is a cause, and congenital syphilis has been indicted, but on insufficient grounds. A history of convulsions is very common.

Treatment depends upon the density and the diameter of the opacity. In cases with dense opacity and very poor vision with undilated pupils the treatment depends upon the diameter

of the opacity. If it is small with a wide area of clear cortex, and if distant vision is much improved when the pupil is dilated and the refraction corrected so far as possible, an optical iridectomy may be performed. In some cases, however, the opacity increases, so that iridectomy should not be done until there is fair certainty that the condition is stationary. Suitable cases for this operation are therefore quite uncommon. Usually the opacity is large, and it is then necessary to remove the lens, which has the grave disadvantage that it abolishes accommodation. Since the patients are almost always seen when quite young, the central core of the lens does not yet form a hard nucleus. Non-sclerosed lens fibres become absorbed if the aqueous gains access to them. Hence lamellar cataract can be treated by discission or needling, whereby an aperture is made in the anterior capsule through which the aqueous enters. This is the ordinary treatment of lamellar cataract, but it should not be employed unless the vision is seriously impaired or the other methods of treatment are impossible. As all varieties of density are met with the advisability of needling in cases with fair vision has to be considered. The decision of this question depends upon whether vision with corrected refraction and retained accommodation is to be preferred to probably improved vision after operation without accommodation. I am of the opinion that vision of 6/12, or even 6/18, with retained accommodation, is more valuable than a problematic 6/9, or even 6/6 without accommodation, but with the necessity of wearing constantly very strong convex glasses. I do not therefore operate in such cases.



Fig. 177.—Anterior capsular cataract. (Nettleship.)

It is not advisable to needle lamellar cataracts until the child is 9 or 10 months old. During the waiting period the pupils should be kept dilated with 0·5 per cent. atropin once every other day, so that the retina may be stimulated by light passing through the clear zone. This procedure diminishes the risk of nystagmus or squint developing.

Anterior Capsular Cataract (*Syn.—Anterior Polar Cataract*).—This form of cataract is commonly known as anterior polar. It is best to reserve this term for any cataract at or near the anterior pole of the lens, since there are two forms of anterior polar cataract, viz., anterior capsular and anterior cortical. Similarly the term posterior polar should be used in the same manner, though there is more ambiguity here. There are two

forms of opacity which are known as posterior polar cataract. The posterior cortical cataract is the commonest form of complicated cataract (*vide* p. 302) (Fig. 181). The other posterior polar "cataract" is not strictly a cataract at all, since it is due to persistence of part of the posterior vascular sheath of the lens, and is therefore situated upon the posterior surface of the lens, *i.e.*, it is not a true lenticular opacity (Fig. 183), though there is always some opacity in the adjacent lens fibres.

Anterior capsular cataract is due to abnormal proliferation of the cubical cells which line the anterior capsule (Fig. 178), and is usually limited in uncomplicated cases to a small area in the centre of the pupil (Fig. 177). The stimulus to proliferation is caused by contact with the normal or inflamed cornea. Contact of the lens with the normal cornea causes opacity in the lens only if it occurs at an early age. This is indeed fortunate, for if it were not so most intraocular operations, *e.g.*, iridectomy,



Fig. 178.—Diagram of section of anterior capsular cataract.

would be impossible. In very young children it is probable that a very short time of contact is all that is necessary. The older the patient the longer is the time required. Contact with an inflamed cornea is more liable to produce an anterior capsular cataract than with the normal cornea. In most cases it is due to perforation of a corneal ulcer, usually caused by ophthalmia neonatorum; more rarely to a perforating wound.

Anterior capsular cataract is sometimes congenital, when it is also probably due to contact with the cornea, possibly without actual perforation, or to delayed formation of the anterior chamber. In these cases it is almost always bilateral, whereas the acquired form is generally unilateral.

When an ulcer perforates the aqueous escapes and the lens and iris are driven forwards into contact with the back of the cornea. If the perforation becomes blocked with iris the anterior chamber re-forms, the length of time of contact between the lens and cornea varying in different cases. If it

is short, no harm may be done to the lens unless the patient is very young. If it is more prolonged an anterior capsular cataract is formed and the lens adheres more or less to the wound. When the anterior chamber re-forms the lens usually separates completely from the cornea: less frequently the adhesion stretches out into a fine filament, which may persist or break. Occasionally the adhesion is so firm that the lens is permanently anchored to the cornea; the eye is usually lost by panophthalmitis or secondary glaucoma in these cases.

The dragging upon the adhesion when the anterior chamber is re-formed may cause a conical protrusion of the cataract—*pyramidal cataract* (Figs. 114, 177).

The deleterious effects of contact may affect the underlying cortical fibres, so that an anterior cortical cataract may occur with a capsular one. In the absence of cortical degeneration the opacity is so small and sharply defined that vision is little impaired, and no treatment is required.

Congenital Cataract manifests itself in a variety of different forms. As already stated, *anterior capsular* and typical *lamellar cataracts* may be congenital. Many other forms are nearly allied to the lamellar type, those occurring early in fœtal life being small in diameter (*vide* p. 296). To this category belong the following:—

Central or nuclear cataract, a small spherical opacity in the centre of the lens, surrounded by clear cortex.

Fusiform cataract, also called spindle-shaped, axial, or coralliform, an antero-posterior spindle-shaped opacity, sometimes with offshoots giving an appearance much resembling coral: this form shows a great tendency to occur in families. *Discoid cataract* is also a familial form, showing a somewhat ill-defined disc of opacity just behind the nucleus in the posterior cortex.

In other congenital cataracts minute points of opacity are seen scattered throughout the lens, or limited to parts—*punctate cataract*. Many varieties of this type occur. Bluish spots, seen by oblique illumination near the anterior surface of the lens, and of congenital origin, are not uncommon. They are almost transparent when viewed with the ophthalmoscope, remain unchanged throughout life and require no treatment. A single minute round opaque spot, usually eccentric, and situated deeper in the lens substance, is not infrequently seen in the routine examination of patients with the ophthalmoscope. It is doubtless congenital and is stationary.

Most of these congenital cataracts are stationary. They may

be associated with other congenital stigmata, such as nystagmus (*vide* p. 514), congenital colobomata, &c. They may require no treatment, or optical iridectomy or discission may be indicated (*vide* p. 297). It is wise to wait until puberty in cases apparently suitable for optical iridectomy, the pupils being kept dilated with atropin in the meantime, since some are not stationary but gradually progress to the formation of total cataract.

Total cataract may be congenital or the result of progressive partial congenital cataract. The lens may be shrunken and much degenerated, and there are often other congenital defects in the fundus, &c. These cases should be treated by discission, but the prognosis given must be very guarded. Sometimes needling reveals persistence of the posterior vascular sheath of



Fig. 179.—Secondary cataract from a section by Treacher Collins.

the lens, with or without persistence of the hyaloid artery. In such cases violent attempts to remove the opacity by needling will result in the loss of the eye.

The pupil is often small with congenital total cataract and reacts very feebly, if at all, to light and on convergence. Neither does it dilate appreciably with atropin. Attempted discission of the shrunken cataract often causes rupture of suspensory ligament, and it may be necessary to remove the membranous lens with iris forceps through a keratome incision. Vitreous is usually lost and there may be severe reaction, but the result not infrequently justifies the heroic measures.

Congenital cataracts should not in general be needled until the child is 9 or 10 months old. If there is a clear peripheral zone they should be treated like lamellar cataracts (*vide* p. 297). In cases where the lens is completely opaque, or the pupil will not dilate, and when squint or nystagmus is developing, it is

advisable to needle at a much earlier age, though there is some risk in doing so.

The Treatment of Unilateral Cataract in Children.—When dense unilateral cataract occurs in a child, whether from lamellar or congenital cataract or from traumatism (*vide* "Traumatic Cataract"), so that the pupil becomes grey or white, it is advisable to needle early. In these cases the appearance of the eye militates against the individual obtaining employment. Moreover, needling is a less severe operation than extraction of cataract, which may become necessary at a later date if the other eye fails. The treatment of unilateral cataract in children differs therefore from that of a similar condition in adults (*vide* p. 293).

Secondary Cataract (*Syn.*—*After-cataract*) is the opacity which persists or follows after the extraction or discission of the lens, In both these operations the posterior and part of the anterior capsule remain *in situ*. If only the posterior capsule remains



FIG. 180.—Posterior cortical cataract. 1, 2, 3, as in Fig. 169. (Nettleship.)

in the pupillary area the corrected vision will probably be good, though it may be much impaired by wrinkling of the capsule and the consequent irregular refraction. In many cases, especially when the cataract is not quite mature, some soft, clear cortex sticks to the capsule. This becomes partially absorbed by the action of the aqueous, but it often becomes shut off from the aqueous by adhesion of the remains of the anterior capsule to the posterior capsule. In such cases the cubical cells which line the anterior capsule also persist; they continue to fulfil their function of forming new lens fibres, though those formed under the abnormal conditions are abortive and opaque (Fig. 179). If these remnants lie in the pupillary area a dense membrane is formed through which the rays of light penetrate with difficulty, so that vision is very imperfect. If the previous operation has been followed by iritis, exudates also adhere to the lens remnants and organise, thus contributing a fibrous membrane in addition.

Secondary cataract is demonstrated either by oblique illumination or by the ophthalmoscope. If fine, it may be

difficult to see, forming a grey film by the former method, a cobweb-like haze by the latter. The denser membranes are easily recognised. They vary in density, showing coarse opaque bands separated by more transparent areas.

Treatment.—Secondary cataract requires needling (*vide* p. 443).

Complicated Cataracts (*Syn.—Secondary cataracts*) are those forms which result from malnutrition of the lens, due to disease of other parts of the eye or of the general system. The lens is nourished by lymph which is supplied by the ciliary body. If, owing to disease of the ciliary body or to lymph secreted from abnormal blood, the nutrition of the lens suffers, opacities are formed. They usually commence in the centre of the posterior part of the cortex, and are therefore at first *posterior cortical cataracts* (often called posterior polar) (*vide* p. 298) (Figs. 180—1). The opacity seldom remains confined to this situation: it progresses, affecting first the periphery of the anterior cortex close to the nucleus, finally involving the whole cortex. In many cases the opacities are fine and dust-like, and are scattered throughout the cortex from the commencement, increasing in number and density as time goes on.



FIG. 181.—Posterior cortical cataract (posterior polar cataract), seen by reflected light.

The total cataract formed in this manner is usually soft and uniform in appearance. In still later stages the watery constituents become absorbed, the capsule becomes thickened, the whole lens shrinks, giving rise to tremulousness of the iris, and other degenerative changes—calcification, &c.—ensue.

Complicated cataracts occur in advanced cases of cyclitis, in absolute glaucoma, in choroido-retinitis—disseminated choroiditis, "retinitis" pigmentosa, &c.—in high myopia, in detachment of the retina, &c.; they also occur in suppurative inflammation of the cornea, especially that produced by ulcus serpens. The opacity in the posterior cortex, which is generally stellate in shape, is seen in its most characteristic form in "retinitis" pigmentosa, in which disease also its slow progress can be easily watched. The vision is already much diminished before complicated cataract makes its appearance. This fact is of the utmost importance from the prognostic point of view, since, even if the cataract is successfully removed, the pro-

gressive diminution in vision due to changes in the fundus is not thereby influenced. In every doubtful case not only must the central and peripheral vision be carefully investigated, but an exhaustive search must be made for precipitates upon the back of the cornea.

Treatment must be directed in the first case to the cause of the complication. This is often tedious and unsatisfactory, but must be persevered in as long as useful vision persists. If then the perception and projection of light appear to be fairly good, and the cataract is of a nature suitable for operation, it should be removed by discission or extraction, according to the age of the patient. Many cases are not suitable for operation, mostly on account of the very defective vision and projection of light. Even in these, if there is a possibility of success, operation may be undertaken after warning the patient of the doubtful issue, for the loss of such an eye weighs little against a possibility of improved vision. It is wise in these cases to do a preliminary iridectomy (*vide* p. 294).



FIG. 182.—Coloboma of the lens (Marcus Gunn). The direction of the coloboma, upwards, is somewhat unusual. Note the defective development of the suspensory ligament of the lens.

Diabetic Cataract should be regarded as a form of complicated cataract. Cataract in diabetic persons is by no means always diabetic in the proper sense of the term. Senile cataract of the usual type, following the usual course, often occurs, and should be treated in the ordinary manner, though in the early stages the general disease must invariably receive every attention, both in the matter of diet and drugs. True diabetic cataract is comparatively rare. It is always bilateral. Dusty opacities appear throughout the cortex and rapidly increase until total cataract supervenes. A uniform milky opacity of rapid onset should always suggest the possibility of diabetes, though of course the urine should be tested as a matter of routine in all cases of cataract.

Diabetic cataract, though usually occurring in patients with a large percentage of sugar in the urine, is not due to the mere presence of sugar in the aqueous, for it is never sufficiently concentrated to cause cataract *per se*. The cause is unknown,

but is probably some deleterious agent in the lymph secreted by the ciliary body.

Treatment.—It is imperative in all cases of true diabetic cataract to treat the general condition before adopting operative measures. It is rare for the opacity to clear up under such treatment; but since cases do occur, and since operations upon the eyes of diabetic patients have special dangers of their own, this chance should always be afforded.

If general treatment is unsuccessful the cataract must be extracted. In spite of the special difficulties attending the operation the results are often quite satisfactory, though a guarded prognosis should be given. Contrary to what might be anticipated, the wound usually heals well. The special dangers are local and general. Of the former, the tendency to severe iritis and to hæmorrhage are the most important. The necessary manipulation of the iris is likely to set up traumatic iritis of a peculiarly violent type. Iridectomy in the course of the operation may be attended with much hæmorrhage, which not only obscures the view of the field of operation, but may endanger the eye: violent intra-vitreous or subchoroidal hæmorrhage may destroy the eye at the time of operation. Hence it is desirable to remove a diabetic cataract by simple extraction, *i.e.*, extraction without iridectomy, or at any rate with only a peripheral button-hole iridectomy (*vide* p. 456). The chief danger affecting the general system in these cases is the risk of the sudden onset of diabetic coma. It is comparatively slight, and must be guarded against as far as possible by a suitable course of anti-diabetic treatment before operating. Another grave danger is that of septic infection. If it occurs panophthalmitis rapidly supervenes, owing to the very defective resistance of the tissues.

Traumatic Cataract.—See p. 408.

Persistence of the Posterior Vascular Sheath (*Syn.*—*Posterior Polar Cataract*).—See p. 310.

Glass-workers' Cataract occurs chiefly in men who have long been engaged in glass manufacture, particularly beer bottles and plate glass. It does not affect those who make flint-glass bottles or pressed glass articles. The cataract is very characteristic. In the early stages there is a small disc of opacity in the posterior cortex of the lens, but it may extend throughout the cortex in the later stages. The radiations from the molten glass contain few ultra-violet rays, which are therefore probably not the cause. It is most likely that heat is the cause, acting, not directly on the lens, but upon the iris and ciliary body, thus influencing the

nutrition of the lens. As already stated (p. 19), heat radiation is absorbed by the pigment epithelium of the iris, ciliary body, and retina.

It has recently been shown that certain *iron-workers*, especially tin-plate millmen and chain-makers, suffer from an identical condition. It is apparently rare in other iron-workers, probably because they seldom look at the hot metal, and when doing so protect their eyes with coloured glass. It does not occur among acetylene welders, &c., who are unable to carry out their work without efficient protection of the eyes by suitable deeply tinted protective goggles.

Congenital Abnormalities of the Lens.

Besides the various forms of *congenital cataract* (*vide* p. 299), abnormalities in the shape and position of the lens occur, often associated with other malformations of the eye.

Coloboma of the lens is the condition in which there is a defect in the inferior margin, usually notch-shaped; less frequently it occurs in some other part of the margin (Fig. 182). It is due to defective development of part of the suspensory ligament.

Ectopia lentis or congenital dislocation is a subluxation of the lens, usually upwards or up and in, and bilateral. The condition is often hereditary. The lens is small, but the edge is generally invisible until the pupil is dilated. The usual signs of subluxation (*vide* p. 401) are then seen.

Lenticonus, which may be anterior or posterior, is an abnormal curvature of the corresponding surface of the lens, so that it is somewhat conical instead of spherical.

CHAPTER XVI.

DISEASES OF THE VITREOUS.

The vitreous humour is an inert, jelly-like structure which subserves optical functions. It possesses no blood vessels in post-natal life, and is incapable of becoming inflamed: the old term "hyalitis" rests upon misconception, and should be avoided. We have, therefore, to deal only with symptomatic conditions.

Opacities.—Black specks, floating before the eyes, are seen by normal persons under favourable conditions. These *muscæ volitantes* are opacities of various kinds, viewed entoptically, *i.e.*, they throw a shadow upon the sentient elements of the retina, thus appearing as dark spots in the field of vision. Any relatively intransparent bodies situated anterior to the rods and cones are therefore able to produce muscæ. To this category belong the corpuscles circulating in the retinal blood vessels; if it were not for the fact that the retina is normally adapted for red light the entoptic images of the circulating corpuscles would be a serious impediment to clear vision. Other muscæ are due to minute specks in the vitreous, so small and so slightly intransparent that they cannot be seen objectively by the ophthalmoscope.

Under abnormal conditions muscæ may be so increased as to interfere with vision and to become visible by the ophthalmoscope. They then indicate some disease of the uveal tract, particularly of the ciliary body: they are found in cyclitis, retino-choroiditis, myopia, &c. In their slightest manifestation they are dust-like opacities, which may permeate the whole vitreous or be limited to the anterior part. When very fine a plane mirror and magnification by a convex lens are necessary in order to distinguish them (*vide* p. 104). They are due to minute albuminous coagula and aggregations of leucocytes, the former derived from the ciliary body and choroid, the latter only from the ciliary body and possibly the retina. In the more severe cases flakes and threads are seen, and the entoptic images may be so sharply defined that an intelligent

patient is able to draw them accurately. The larger opacities are often found after hæmorrhage into the vitreous. They almost invariably float about, showing that the vitreous is fluid (*vide infra*), though they may be more or less anchored to the retina.

Dense vitreous opacities obscure the view of the fundus with the ophthalmoscope. In moderate cases the disc and vessels may be made out, as if seen through a dense haze. The disc looks redder than usual, and it may be difficult to decide whether there is papillitis or not.

Very frequently in the slighter cases no objective signs of disease can be made out in the fundus; the foci are either too fine to be appreciated or are anterior to the field of ophthalmoscopic vision, *i.e.*, in the anterior part of the choroid or in the ciliary body.

Treatment.—Slight cases of muscæ volitantes, without objective signs, require no direct treatment. Patients should be advised to ignore the spots as much as possible, as they are often only visible when attention is specially directed to them. Any error of refraction should be corrected, and special indications as to the amount and conditions of near work should be given. In many cases the patients suffer from gastro-intestinal disorders, which should be suitably treated.

Treatment of the more severe cases of vitreous opacity depends upon the cause. When this is known, as in iridocyclitis, syphilitic retinitis, retino-choroiditis, tubercle of the uveal tract and so on, attention must be specially directed to the treatment of the primary foci. The prognosis is best in the syphilitic and the milder iridocyclitic cases.

When the cause cannot be discovered iodides are generally given, as they are supposed to promote absorption. They may be combined with mercury, even in non-syphilitic cases.

The eyes should be kept at rest with atropin and dark glasses. Hot bathings and leeches or dry cupping may be tried; and subconjunctival injections have been beneficial in some cases, but must not be used if there is any active inflammation. Dionin has also been advocated.

Fluidity of the Vitreous (*Syn.*—*Synchisis*, συγχέω, to pour together, disturb) is due to absorption of the fibrile and degeneration of the jelly-like tissue, caused by the digestive action of leucocytes and their secretions. It is therefore a common feature in the cases in which opacities are present, and is due to the same causes. It cannot be diagnosed with certainty in the absence of opacities, and is indicated by their

free movements when the eye is rapidly moved. Fluidity of the vitreous may in some cases be simply a senile degeneration. The tension of the eye may be normal, but soft eyes nearly always contain fluid vitreous.

The degeneration of the vitreous which leads to fluidity often causes the deposition of crystals of cholesterin, which sink to the bottom of the vitreous chamber, but are stirred up by movements of the eye. They then appear as a very beautiful shower of golden rain—*synchisis scintillans*. There may be relatively little interference with vision.

Fluidity of the vitreous requires no treatment in itself, but it is a serious complication if any intraocular operation is contemplated. In such eyes the suspensory ligament of the lens is often weak, so that prolapse of the vitreous, dislocation of the lens, and so on, may occur. Even when the vitreous pours out under these circumstances useful vision may yet be retained, the vitreous being replaced by lymph.

Blood in the Vitreous.—Hæmorrhage into the vitreous may result from arteriosclerosis or inflammation of the retina, from contusions or wounds of the eye, pernicious anæmia, malaria, &c., or without apparent cause. Small hæmorrhages are seen ophthalmoscopically and can be accurately watched, though the surrounding vitreous always contains exudates, unless the blood is subhyaloid (*vide* p. 336). Larger hæmorrhages, filling the vitreous with blood, are suspected when no reflex can be obtained on throwing in light with the mirror. It may then be possible to see a red mass behind the lens by oblique illumination.

There is one form of severe vitreous hæmorrhage which occurs in apparently healthy young adults, usually males: it is probably due to a blood condition associated with defective coagulability. Some of the patients suffer from epistaxis. It has a great tendency to attack both eyes in succession and to recur, so that though absorption may be complete in the early attacks permanent defect or complete loss of sight may ultimately follow from damage to the retina, retinitis proliferans, or dense vitreous opacities. It is characteristic of vitreous hæmorrhage that absorption usually takes place without organisation, owing to the absence of fibroblasts in the vitreous. When organisation occurs, as in so-called retinitis proliferans (*q.v.*), it is most marked near the disc, from which membranes and strands stretch forwards. This is due to the presence of mesoblastic tissue, containing potential fibroblasts, around the central vessels. Even then organisation rarely

occurs in the absence of some general dyscrasia, *e.g.*, syphilis nephritis, &c. In other cases, therefore, the prognosis is good, bearing always in mind the tendency to recurrent hæmorrhage.

Treatment.—Rest is imperative in all cases of vitreous hæmorrhage, and in the early stages the patient should be kept in bed. All straining must be avoided.

In cases of recurrent hæmorrhage attempts may be made to increase the coagulability of the blood by the administration of calcium salts, the treatment being controlled by repeated estimations of the coagulation time.

Fibrous Tissue in the Vitreous is found in cases of plastic iridocyclitis, usually in the form of membranes stretching across behind the lens. They cannot as a rule be seen clinically, as the lens is opaque.

A subacute form of plastic endophthalmitis occurs in children and gives rise to one form of so-called *pseudoglioma*. It is often first noticed by the mother as a whitish reflex in the pupil (amaurotic cat's eye). Generally there is no knowledge of inflammation having occurred in the eye, but a history may be obtained of fits, ear disease, an acute specific fever, post-basic meningitis or some serious illness. Several cases have been proved by pathological examination to have been due to the meningococcus. There are usually signs of past iritis or iridocyclitis—posterior synchiæ, &c. The diagnosis from glioma is of great importance (*vide* p. 391).

As already mentioned, recurrent hæmorrhages are liable to give rise to retinitis proliferans, which is really a formation of fibrous tissue in the vitreous.

In some cases of this type, especially in syphilitic, gouty, and possibly tuberculous subjects, minute bunches of new-formed blood vessels project from the disc or retina unusually near the disc into the vitreous (*vide* p. 339).

Pus in the Vitreous.—This is found only in panophthalmitis, which is almost invariably due to a perforating wound or ulcer, though cases of metastatic inflammation of the retina and choroid also occur and lead to a similar result (*vide* p. 318). The reflex with the ophthalmoscopic mirror is poor or absent. Oblique illumination shows a yellow mass behind the lens. The eye is always intensely inflamed, and little difficulty is usually experienced in arriving at a correct diagnosis. The *treatment* in cases of exogenous infection is that of panophthalmitis due to perforating injury (*vide* p. 424). In the rare cases of endogenous infection it is that of the primary cause.

Persistent Hyaloid Artery.—The hyaloid artery, which in

fœtal life nourishes the vitreous and supplies blood to the posterior vascular sheath of the lens, becomes obliterated during the sixth and seventh months, and usually disappears completely before birth. A remnant, stretching forwards from its origin on the disc, is normal in oxen. Similar remnants are not very rare in man : they appear as a strand projecting from the disc into the vitreous. There may be membranes veiling the disc from view in such cases. In other cases only the anterior part of the hyaloid artery persists, appearing as a circular spot on the back of the lens, often inaccurately termed a *posterior polar cataract* (Fig. 183). More or less of the posterior vascular sheath may persist in such cases, giving rise



Fig. 183.—Opacity due to persistence of part of the posterior vascular sheath of the lens, often called posterior polar cataract. Note that the opacity is usually more circumscribed than in posterior cortical cataract. (Fig. 181.)



Fig. 184.—Diagram of persistent hyaloid artery, with persistence of the posterior vascular sheath of the lens. (Treacher Collins.)

to a larger central opacity on the back of the lens. In these cases there is a gap in the posterior capsule of the lens and the adjacent posterior cortical lens fibres are cataractous. The hyaloid artery may persist in its entirety with more or less of the vascular sheath (Fig. 184). It may contain blood, and blood vessels may be seen upon the back of the lens ; it is then liable to be mistaken for a glioma of the retina, and constitutes one form of pseudoglioma (*vide* p. 391).

The appearance of the posterior part of the hyaloid artery, when persistent, is very characteristic. A filmy, greyish cord, sometimes containing blood, passes forwards from the disc towards the lens. It undergoes serpentine writhings when the eye is moved.

When the hyaloid artery is persistent there is often defective or atypical development of the vitreous, part of which resembles fibrous tissue in structure. Every stage may be met with, from that interfering little or not at all with vision to complete opacity, usually with maldevelopment of the whole eye—microphthalmia.

Foreign Bodies in the Vitreous.—See p. 414.

Parasites in the Vitreous.—*Cysticercus* is excessively rarely found in the vitreous in England, though it is less uncommon in some other countries. The actual parasite may be seen ophthalmoscopically as a pearly translucent mass with peristaltic movements. The treatment—removal—is very difficult.

CHAPTER XVII.

DISEASES OF THE CHOROID AND RETINA.

It has already been pointed out that different parts of the uveal tract rarely suffer alone. This intimate connection is most marked in the anterior parts, and clinically the evidence of cyclitis when the choroid is inflamed is slight, though not entirely wanting. The outer layers of the retina are dependent for their nutrition upon the choroid, so that when the latter suffers the former is always involved secondarily. Primary affections of the retina occur without involvement of the choroid: primary affections of the choroid invariably involve the retina secondarily in greater or less degree.

This profound relationship between the retina and choroid, so indisputably manifested clinically, renders it advisable to consider their diseases in close connection with each other. It will be found that some diseases commonly designated as "retinitis" are in reality secondary to a primary choroiditis, whilst others also included under the same term are not inflammatory, as is suggested by the word. It is well, therefore, to bear in mind that "retinitis" is used in a broad sense. The same ambiguity is noticed in the use of the term "choroiditis," which frequently designates a purely degenerative condition, without any evidence of inflammation.

Primary Affections of the Choroid.

Inflammation affecting the choroid primarily—*choroiditis*—occurs in two forms, exudative and suppurative. The former appears in the form of isolated foci of inflammation scattered over certain areas of the fundus, and is conveniently classified according to the position of the areas involved. The latter spreads over the whole choroid and retina, and the primary seat may be in the retina: it leads ultimately to panophthalmitis.

Exudative choroiditis is generally syphilitic in origin, though certainly not so generally as was formerly thought; it affects chiefly either the posterior part of the fundus—disseminated choroiditis—or the anterior part—anterior choroiditis.



FIG. 1.

FIG. 2.

PLATE IX. (*To face p.* 313.)

Fig. 1.—Disseminated choroiditis. Note that the retinal vessels pass anterior to the atrophic spots.

Fig. 2.—Ruptured choroid (R. eye, indirect method), due to blow with a fist eighteen months previously.

Disseminated Choroiditis may be taken as a type of the disease (Plate IX., Fig. 1). The recent foci are seen ophthalmoscopically as round yellowish spots; when near a retinal vessel they lie at a deeper level than the vessel. They are due to infiltration of the choroid, the exudates hiding the choroidal vessels which cause the normal red reflex. In the early stages the elastic membrane of Bruch is intact: under these circumstances only fluid exudates can pass through it, but these suffice to make the overlying retina cloudy and grey. Hence the edges of the spots are a little hazy and ill-defined. The exudates not only pass into, but also through the retina, so that punctate or diffuse opacities are seen in the vitreous. When the vitreous haze is excessive the ciliary body is also probably involved. In later stages the membrane of Bruch may be absorbed, though it offers considerable resistance in common with all elastic membranes. When this has occurred leucocytes are enabled to pass through into the retina and vitreous.

FIG. 185.—Atrophy after syphilitic choroiditis. (Nettleship, after Hutchinson.) *a*, atrophy of pigment epithelium; *b*, atrophy of epithelium and choriocapillaris, exposing the large choroidal vessels; *c*, spots of complete atrophy, many with pigment accumulation.

Owing to the fibroblastic activity of the choroidal stroma the exudates become organised, so that a small white mass of fibrous tissue is formed, which destroys the normal structures of the choroid and retina, and fuses the two membranes firmly together. The colour of the spots therefore gradually changes to white, partly due to the fibrous tissue deposited, partly to thinning and atrophy, whereby the white reflex from the sclerotic is permitted to shine through (Fig. 185).

The pigment of the retinal pigment epithelium is extremely resistant, even though the cells which contain it be destroyed. It tends to become heaped up into masses, partly intra-, partly extra-cellular: moreover the pigment cells are stimulated to proliferate. Isolated masses of black pigment are thus

formed in the white areas, but more particularly at the edges, so that in the atrophic stage white spots surrounded by a black zone of pigment are seen (Fig. 186). The process has then reached its natural termination, and these spots remain permanently almost unaltered. They are much more sharply defined than the actively inflammatory spots.

Meanwhile fresh foci arise and pass through the same stages, until finally the whole fundus may be covered with atrophic



FIG. 186. Section of a patch of disseminated choroiditis, showing the fusion of the degenerated retina (*R*) and choroid (*Ch*) (× 60). Note the heaping up of the retinal pigment epithelium (*P*) at the edges of the adhesion. *Scl.*, sclerotic.

spots. In the milder cases only a few spots are formed and the exudates in the vitreous become absorbed. In the more severe the spots are very numerous, the vitreous opacities increase, and finally the nutrition of the lens suffers and a complicated cataract (*q.v.*) results. Owing to the transiency of the acute stage the atrophic stage naturally comes much more frequently under observation.

The symptoms in the early stages are principally the defects of vision due to the retinal lesions and to cloudiness of the vitreous. The spots are slightly raised, so that the contour

of the retina is altered. This causes distortion of the images, giving rise to similar appearance of distortion of the objects seen—metamorphopsia : thus straight lines appear to be bent in various directions. Frequently objects appear smaller than they are—micropsia ; sometimes larger—macropsia : these results are probably due to separation or crowding together respectively of the rods and cones. They are not likely to be noticed unless the macular region is involved. Subjective symptoms of light—photopsiæ—occur, such as flashes of light, due to retinal irritability. These subjective symptoms are often accompanied by the perception of a black spot in front of the eye, corresponding with the lesion—*positive scotoma.*

In the later stages the affected spots are incapable of giving rise to visual impulses, so that *negative scotomata* exist in the field of vision, *i.e.*, though there is no perception of a spot in front of the eye there is a hiatus in the field of vision of the same nature as the normal blind spot. Their relative importance depends upon their situation. Peripheral scotomata may pass unnoticed, but central scotoma destroys direct vision. Peripheral vision still permits the patient to get about well, but all fine work is impossible.

The disease is chronic, organisation of the exudates taking several weeks. The occurrence of fresh spots may extend the acute stage over months, and the ultimate defects are permanent.

As already mentioned, the disease is usually due to syphilis, generally acquired, sometimes congenital, and hence associated frequently with interstitial keratitis ; but in many cases the cause is obscure, sepsis, anæmia and disorders of nutrition being assigned. The changes produced by myopia cause similar signs and symptoms ; they are not inflammatory, but degenerative from the commencement (Chap. XXIV.).

Treatment is primarily that of the ætiological factor—syphilis or such cause as can be discovered. Iodide of potassium may assist absorption in all cases, and should be administered. The general régime advised for cyclitis (*q.v.*) is suitable in these cases. Marked irritative symptoms indicate the use of dark glasses, the abandonment of all near work, and sometimes the application of leeches.

Anterior Choroiditis is also usually syphilitic, and manifests itself in the same form as disseminated, but is confined to the peripheral parts of the fundus. On this account it is frequently discovered only by the ophthalmoscope. Similar changes are

also sometimes found in high myopia. Simple pigmentary changes at the periphery occur in old people as a senile degeneration.

The periphery of the fundus is often peppered over with minute spots of pigment in congenital syphilitis; this is possibly a purely retinal affection. It can only be distinguished by degree from a similar pigmentation which may be a mere idiosyncrasy.

Central Choroiditis occurs in disseminated choroiditis, and in certain rare forms. In Förster's areolar central choroiditis the spots are said to behave in exactly the opposite manner to those of disseminated; they are first black, then enlarge, becoming white in the central, and finally quite white. The disease extends outwards, the peripheral spots being always the most recent.

Juxtapapillary Choroiditis (*Retinochoroiditis juxtapapillaris*) (Jensen) occurs in young persons, as an exudation close to the disc, oval in shape and about the same size as the disc. The exudates cover the retinal vessels, and there are vitreous opacities and sometimes "k.p." There is a sector-shaped defect in the field of vision. The cause is unknown. The inflammation slowly subsides, leaving a patch of atrophy. Recurrence may take place.

Diffuse Choroiditis is characterised in the early acute stage by one or more plaques of yellowish white or grey areas, shading off at the edges into normal fundus. The patches spread and coalesce so that the greater part of the fundus may be finally involved. The exudates gradually organise, leaving white areas in which the larger choroidal vessels persist, as a characteristic band-like network. The retinal pigment becomes heaped up into dense black, irregular spots, variously grouped. The retinal vessels course over the patches little changed in appearance. The coalesced areas leave islands and spaces of normal fundus between them when organisation is complete. Fresh exudates occur simultaneously with the organisation of older ones, and sometimes the edge of a patch appears to creep over the fundus like the advance of a myxomycetes.

Some of these cases are syphilitic, some tuberculous (*q.v.*). In others the cause cannot be traced, but probably many are due to metastatic bacterial invasion (*vide infra*, Metastatic Endophthalmitis).

Purulent Choroiditis.—See p. 423.

Tubercle of the Choroid occurs in acute or miliary and chronic forms. Miliary tubercles are found in the late stages of acute

miliary tuberculosis, especially tuberculous meningitis. Occasionally they may be seen before there is any evidence of meningitis or generalised tubercle. They are very common in the late stages of tuberculous meningitis in children, though they may appear only a day or two before death. My own observations, clinical and post-mortem, lead me to think that they are nearly always present in this disease. Ophthalmoscopically they appear as round, pale yellow spots, most frequently observed in the neighbourhood of the disc, though any part of the choroid may be attacked. Generally only three or four spots are seen, but as many as sixty or seventy have



FIG. 187.—Section of miliary tubercle of the choroid (× 60), from a child with hip disease who died of meningitis. Note the giant cells and the small round cells at the periphery.

been found. They vary in size from pin-point specks to 1 or 2 mm. in diameter. They usually project slightly, so as to raise the retina, but the inner surface is often quite flat, while the outer surface projects into the sclera. They afford most important diagnostic evidence of tubercle in cases of meningitis and obscure general disease. Microscopically they consist of typical giant-cell systems, containing a variable number of tubercle bacilli (Fig. 187).

Chronic tubercle of the choroid may occur as a diffuse inflammation affecting large areas or the whole choroid and characterised by the extensive development of granulation tissue, or, more rarely, as a solitary or conglomerate mass, simulating sarcoma. In the former case the choroid becomes

replaced by granulation tissue containing giant cells, which spreads until it involves the retina and finally fills the posterior part of the globe. The sclerotic becomes eroded ; perforation takes place, usually near the anterior part of the eye, a fungating mass appearing under the conjunctiva. The chronic forms of tubercle of the choroid may nearly simulate glioma of the retina (*q.v.*), constituting one form of pseudoglioma (*q.v.*).

They are commoner in children than in adults, but in all cases certainty as to their tuberculous nature can rarely be obtained clinically. Similar ophthalmoscopic appearances are met with resulting from metastatic choroiditis (*q.v.*), and from changes following localised hæmorrhage into the deep layers of the retina (so-called " massive exudation " in the retina). von Pirquet's test or injection of tuberculin may afford help in the diagnosis.

Treatment.—No local treatment is indicated in miliary tubercle of the choroid. The patient quickly succumbs to the general disease. In diffuse and conglomerate tubercle treatment with tuberculin should be instituted in the early stages, but if the eye is extensively involved it is best to enucleate it, and thus remove a dangerous focus from which the organism may be disseminated into the system.

Metastatic Endophthalmitis.—Endogenous bacterial infection of the eye manifests itself in various ways. Most commonly the uveal tract is affected, either as a whole (metastatic uveitis) or in its individual parts, as for example in gonorrhœal iritis. The milder forms of iridocyclitis (*q.v.*) and uveitis are probably due to toxins circulating in the blood-stream, and derived from bacterial foci in other parts of the body, *e.g.*, the mouth, the generative apparatus, especially in women, the intestinal tract and so on. It is certain, however, from anatomical examination that actual bacterial embolism occurs, and in these cases two facts stand out prominently. First, various organisms show a special selectivity for the various structures of the eye. Thus the tubercle bacillus never attacks the retina primarily, though this structure is often involved secondarily in tuberculous disease of the choroid. Second, except in the case of extremely virulent organisms such as the streptococcus the inflammation set up by endogenous infection is usually less severe than when the organism is introduced directly into the eye from without. Thus, if the eye is infected with pneumococci by a perforating wound an exogenous panophthalmitis (*q.v.*) is likely to be set up, leading to the complete destruction of the organ. So much is this the case that saprophytic organisms, such as the bacillus

subtilis, which are non pathogenic in other parts of the body, often cause panophthalmitis. If, however, an organism such as the pneumococcus invades the eye by way of the blood-stream, though an intense inflammation results it tends to subside more rapidly than in the exogenous cases. Probably the organism becomes attenuated in the blood-stream and tissues through the controlling effect of specific anti-bodies.

FIG. 188.—Metastatic choroiditis.

In this manner virulent bacteria may set up endophthalmitis of every grade of severity. In the days when puerperal fever was prevalent, intense metastatic panophthalmitis, often attacking both eyes, was not uncommon. It was characterised by the appearance of an hypopyon and the rapid development of a yellow pupillary reflex due to pus in the vitreous. An intense metastatic uveitis, involving iris, ciliary body and choroid, and due to bacterial invasion, was set up. Such cases progressed exactly like ordinary exogenous panophthalmitis (*q.v.*).

They are now seldom seen, but cases of metastatic bacterial invasion often of obscure origin still occur in which the early stage of uveitis with hypopyon is present. They occur in the course of infectious diseases, especially pneumonia, influenza, measles and scarlet fever, and of meningitis, dysentery, &c. Ophthalmoscopically the media are hazy, so that the yellow œdematous retina is only dimly seen. Under treatment the condition gradually subsides, not infrequently with the restoration of useful vision. In more severe cases the inflammation causes destruction of the ciliary processes, the intraocular tension falls, and the eye gradually shrinks. In children the inflammation is probably often due to the diplococcus intracellularis of Weichselbaum and is followed by an exuberant development of new fibrous tissue of cyclitic origin. These cases are not easily distinguished clinically from glioma, and form the largest group of so-called pseudoglioma (*q.v.*). In the milder cases there is less development of cyclitic fibrous tissue in the vitreous and less scarring of the retina, but whitish atrophic spots and areas accompanied by pigment disturbance persist as evidence of the previous inflammation. Yet milder cases give rise to ophthalmoscopic appearances which may be mistaken for diffuse or conglomerate tuberculous diseases of the choroid (*q.v.*) (Fig. 188), or thrombosis of a branch of the central vein of the retina. In some cases the optic papilla has been the chief local focus, and an appearance of intense papillitis with much exudation extending into the neighbouring retina is seen. Some of the cases have been associated with furunculosis, and there is little doubt that they are due to metastatic infection with staphylococci. Others have been proved to be due to the pneumococcus, diplococcus intracellularis of Weichselbaum, &c. Other obscure cases of retinitis with white spots or oval areas, usually near the disc, with or without hæmorrhages, are probably due to the same cause.

The presence of precipitates ("k.p.") on the back of the cornea and inconspicuous posterior synechiæ shows that in many cases of localised endogenous choroiditis the whole uveal tract is really involved (uveitis).

The *treatment* depends upon the severity of the attack, and the possibility of determining the primary focus and the specific organism. In the worst cases the patient should remain in bed, and purgatives and the drugs for the relief of pain should be administered. Hot applications should be made and atropin instilled. If the eye becomes full of pus it should be enucleated or eviscerated according to the principles which

govern the treatment of panophthalmitis of exogenous origin (*q.v.*).

In the milder cases purgatives and general tonic treatment are indicated. Atropin should be instilled to keep the eye at rest, and dark glasses worn. Counter-irritation with leeches or blisters to the temple may be indicated. If the primary source of infection can be discovered it must be treated radically. If the organism can be isolated vaccine treatment is indicated, and in the more obscure cases treatment with a polyvalent vaccine may be of some avail.

Suppurative or Purulent Choroiditis.—See Metastatic Endophthalmitis (p. 318) and Panophthalmitis (p. 423).

Degenerative Changes may be post-inflammatory or primary. The former, culminating in localised spots of complete atrophy, have already been considered. Generalised atrophy, more or less complete, is found in the later stages of glaucoma. The loss of nourishment to the retina causes atrophy of the outer layers and of the nerve fibre layer in these cases. Degenerative changes in the choroid often cause migration of pigment from the pigment epithelium into the more superficial parts of the retina. The pigment tends to become deposited in the perivascular lymph spaces of the veins, so that the retinal veins may be mapped out here and there by pigment. More noticeable ophthalmoscopically are jet-black branched spots of pigment, resembling bone corpuscles, and standing out in sharp relief. This condition is seen in its most typical form in "retinitis" pigmentosa, which is in reality due to primary choroidal atrophy. An almost identical picture, though usually without the characteristic distribution of the pigmented spots, may result from choroidal atrophy due to other causes, *e.g.*, syphilis. This migration of pigment into the retina is shown to be due to interference with the choroidal circulation by the fact that it occurs after division of the short ciliary vessels in rabbits.

Primary choroidal degeneration may be localised or general: the best example of the latter is so-called retinitis pigmentosa, which will therefore be considered here.

Localised choroidal atrophy, apart from the post-inflammatory forms, is usually either central or circumpapillary.

Central Choroiditis, or more properly *central choroidal atrophy*, is most commonly the result of myopia (*q.v.*), syphilis, contusion (Chap. XXI.), or old age.

Senile central choroidal atrophy assumes two chief forms. In *central guttate choroiditis* (Tay's choroiditis) there are numerous

minute yellowish white spots in the macular region (Plate X., Fig. 1). They may increase in numbers, but otherwise they remain stationary. They are always small, usually round, but the larger spots may have cremated edges, thus showing signs of fusion. There may be indefinite signs of greyish pigmentation of the edges of the spots, due to the fact that the pigment epithelium is stretched over them (Fig. 189). The condition is bilateral. They have been mistaken for albuminuric retinitis (*q.v.*) and retinitis punctata albescens (*q.v.*). They are easily distinguished from the former, in which the spots are much brighter and more glistening, and are usually arranged in the form of a fan or star ; and from the latter, in which the spots are not limited to the macular region and are

Fig. 189.—Section of " colloid bodies," seated upon the membrane of Bruch. They are undergoing calcification, as shown by the deeper staining in the inner parts. They are covered with stretched pigment epithelial cells.

much whiter. Central guttate choroiditis causes *per se* little impairment of vision, but other senile changes are often present and account for defective sight. The spots are due to peculiar hyaline excrescences on the surface of the choroid, commonly known as *colloid bodies* (Fig. 189). They are of the same nature as Bruch's membrane, and like it are secreted by the pigmented epithelial cells.

Central areolar choroidal atrophy—to be carefully distinguished from Förster's areolar choroiditis (*vide* p. 316)—appears as a large circular or oval patch of degeneration in the macular region through which the choroidal vessels are visible, owing to atrophy of the retinal pigment epithelium (Plate X., Fig. 2). As a result of atrophy of the choroid the sclerotic shines through and the patch is white, though traversed by choroidal vessels. Only the larger choroidal vessels are seen, the smaller ones having disappeared, and even the large ones



PLATE X. (*To face p.* 322.)

Fig. 1. Central guttate choroidal deposits ("Tay's choroiditis"), from a woman, æt. 41, without failure of sight.

Fig. 2.—Central senile areolar choroidal atrophy, from a woman, æt. 60, L. eye; V. J. 20. Note the exposure of the choroidal vessels, the walls of which are thickened by perivascular changes.



FIG. 1.



FIG. 2.

appear smaller than usual owing to degeneration of the walls. There is an absolute central scotoma. The condition is to be feared in cases of cataract in which perception of light seems defective: hence the great importance of investigating the fundus thoroughly in cases of immature cataract. It is possible that this form of central choroidal atrophy is due to extravasation of exudates.

Besides these conditions minute changes limited to the area at and immediately around the fovea centralis occur not infrequently in old people and lead to grave disturbance or abolition of central vision. It is generally necessary to dilate the pupil with cocain or homatropin in order that they may be seen, care being taken to instil eserin when the examination is finished (*vide* p. 290). When central vision is very poor in an elderly patient and no cause can be found to account for the defect, such as cataract, glaucoma, &c., pathological changes will probably be found at the macula. In the early stage the fovea is surrounded by a ring of very fine pigment spots. The stippling is more sharply defined on the foveal side which usually has a circular or crenated edge; it diminishes rapidly peripherally, where the fundus becomes normal. The fovea gradually becomes paler in colour and the stippling denser, the change being associated with rapidly increasing failure of vision, until eventually the small central scotoma becomes absolute. Both eyes are affected, but one is usually attacked before the other, and many months may intervene.

Sometimes a round yellowish patch, about the size of the disc, is seen at the macula and may be bilateral. The patient is usually old with obvious disease of the retinal vessels. Some of these patches undoubtedly follow hæmorrhages.

Treatment.—The treatment of these degenerative changes is very unsatisfactory. When central senile changes are seen early the treatment of any source of sepsis, *e.g.*, pyorrhœa alveolaris, should be carried out. Strychin and general tonic treatment is usually applied, but without much success. The patient's mind may be relieved to the extent that the rest of the field of vision is not likely to become affected.

Circumpapillary Choroidal Atrophy is found in myopia (*q.v.*) and in late stages of glaucoma.

"Retinitis" pigmentosa is an extremely chronic, progressive degeneration of the choroid in both eyes, commencing in childhood and often resulting in blindness in middle or advanced life. The choriocapillaris suffers first, so that the nutrition of the outer layers of the retina fails early. The degeneration

commences in a zone near the equator of the eye, and gradually spreads both anteriorly and posteriorly. The macular region is not affected until very late in the disease. The condition is bilateral. The peculiar site of origin has been attributed to the relatively weak choroidal circulation in this zone, which is the meeting place of the short ciliary and the recurrent ciliary arteries (*vide* p. 11).

The symptoms of the disease are very characteristic, the most prominent being defective vision in the dusk (night blind-



Fig. 190.—Ring scotoma, from a case of retinitis pigmentosa.

ness). This symptom may be present several years before pigment is visible in the retina (*vide* p. 325). Vision under low illumination is carried on essentially by the rods (*vide* p. 61), and night-blindness is a sign of defective nutrition of these structures, which, from their situation, suffer first in disease of the choroid.

Examination of the vision may show perfect central visual acuity. Investigation of the field of vision, seldom satisfactory in children, but more reliable in young adults, shows concentric contraction of the fields, specially marked if the illumination is reduced. In early cases a partial or complete

annular or ring scotoma is found (Fig. 190), corresponding to the degenerated zone of retina. As the case progresses the field becomes continually slowly smaller, until at last it is reduced to a small area round the fixation point. Central vision may even now be normal, but the patients are quite incapacitated for getting about, for they are in much the same condition as a person looking down two long cylinders—they see only the thing they are actually looking at and nothing around. They therefore grope about helplessly. Loss of central vision does not usually occur until fifty or sixty, but vision may be lost earlier through cataractous changes occurring in the lens.

Ophthalmoscopic examination shows also a very characteristic picture (Plate XI.). In the zone affected the retina is studded with small jet-black spots, resembling bone corpuscles or spiders. The retinal veins, never the arteries, often have a sheath of pigment for part of their course (*vide* p. 321 ; Plate XI., Fig. 2). As the pigment from the retinal pigment epithelium migrates into the superficial layers the epithelium itself becomes decolorised, so that the choroidal vessels are now visible and the fundus appears tesselated. The pigment spots which lie near the retinal vessels are seen to be anterior to them, so that they hide the course of the vessels. There is no question, therefore, as to their exact position in the retina itself. In this respect they differ from the pigment around spots of choroidal atrophy (*vide* p. 313), in which the retinal vessels can be traced over the spots. The number of pigment spots differs much in individual cases, and they are often very scanty in the early stages.

In the earliest stages it can be proved that it is a zone of the retina which is affected and not the most anterior part, for normal retina may be seen at the peripheral limits of the ophthalmoscopic field. In the later stages this area diminishes *pari passu* with extension of the disease towards the centre.

The retinal blood vessels, both arteries and veins, become extremely attentuated and thread-like.

As the pigmentation increases and the retina becomes more and more atrophic the ganglion cells are destroyed, thus leading to degeneration of their axis cylinders, viz., the fibres in the nerve fibre layer of the retina and the fibres in the optic nerve. Optic atrophy, therefore, sets in and gradually increases. The disc exhibits the characteristics of primary optic atrophy (*vide* p. 366), but is not quite typical of this condition. The disc is pale, but seldom more than yellowish

white, like wax: there is no excess of fibrous tissue, but the vessels are very small and thread-like.

In the later stages the nutrition of the lens suffers. The complicated cataract which is formed is a very typical form of progressive posterior cortical cataract, going on to complete opacification of the cortex.

Only a few late cases have been examined microscopically. In all it has been found that the choroidal vessels show very extensive endovascular degeneration. The same applies to the retinal vessels, but probably these are less affected in the early stages.

The cause of retinitis pigmentosa is unknown. Several members of the same family are often affected, consanguinity of the parents is not infrequent, and the disease may affect several generations. Nettleship found heredity without consanguinity in 23 per cent., consanguinity without heredity in 23 per cent., and heredity combined with consanguinity in 3 to 4 per cent. The patients or other members of the family are found not infrequently to suffer from insanity, epilepsy, or other signs of mental debility. About one-third of the patients are deaf, and 4 per cent. of deaf mutes have retinitis pigmentosa (Nettleship). Evidences of cerebral disease or of congenital anomalies in the eyes or other parts of the body (harelip, supernumerary digits, &c.) may be present. Congenital syphilis may produce similar results, though these cases are seldom quite typical (*vide* p. 321).

In syphilitic choroido-retinitis exactly the same ophthalmoscopic appearances associated with night-blindness, may occur and may be present in both eyes. These cases are rare and can be distinguished by the later onset of night-blindness and the absence of the disease in other members of the family. Much more commonly in syphilis the patches are limited to certain areas of one or both retinæ. It is best to restrict the term "retinitis pigmentosa" to the very typical congenital and familial disease.

Pathology.—The opportunity to examine microscopically eyes with retinitis pigmentosa seldom occurs. In the cases on record the retinæ, especially their outer layers, are very atrophic and contain masses of pigment which have migrated from the pigment epithelium. The choriocapillaris of the choroid is absent under the affected areas: there is some doubt as to degenerative changes in the larger choroidal vessels, which sometimes appear to be little changed. It has been proved experimentally on rabbits that section of short posterior ciliary arteries is followed by

pigmentary degeneration of the retina, quite similar to retinitis pigmentosa, in the areas lying over the parts of choroid supplied by these vessels. This is strong evidence in favour of the view that the disease is primarily due to interference with the choroidal circulation. Treacher Collins, however, has advanced the theory that the disease is due to abiotrophy of the neuro-epithelium, analogous to abiotrophy (Gowers) of the central nervous system.

Treatment is eminently unsatisfactory, since nothing appears to have a decided influence upon the course of the disease. Strychnine by the mouth or by hypodermic injection should be employed, and the constant current has been advocated. Trephining at the corneo-scleral margin has been followed by temporary improvement (Mayou), probably by improving the intraocular circulation. Extraction of the lens may be followed by some improvement in vision, but should not be done until a late stage is reached.

Retinitis pigmentosa sine pigmento is a variety of the disease with the same symptoms, but without visible pigmentation of the retina. It is almost certainly only the early stage of the ordinary disease (Nettleship). It is progressive and leads to optic atrophy, therein differing from *congenital night-blindness*, which is a rare hereditary disease without ophthalmoscopic signs, remaining stationary throughout life.

Allied to these conditions is *retinitis punctata albescens*, in which, with the same history and symptoms, the retina shows hundreds of small white dots distributed fairly uniformly over the whole fundus. The condition is almost certainly an atypical variety of retinitis pigmentosa, and may occur in a family affected with this disease. It differs in the important fact that it is almost if not quite non-progressive, but one case is on record in which the white spots disappeared and typical pigment spots developed (Nettleship).

Detachment of the Choroid.—The choroid is often found detached from the sclerotic in eyes which have been lost by plastic iridocyclitis, glaucoma, &c. Detachment may also occur from severe hæmorrhage or new growth. These cases cannot be diagnosed clinically. Fuchs has shown that the condition is not uncommon after iridectomy for glaucoma and cataract extraction, and attributes it to slight separation of the ciliary body, so that the aqueous percolates from the anterior chamber into the suprachoroidal space. It occurs during the first days after the operation. The anterior chamber is shallow or abolished, and on ophthalmoscopic examination the detached choroid is seen as a dark mass

behind the lens. It may be visible as a dark brown mass by oblique illumination. The prognosis is usually good, the choroid becoming replaced and the anterior chamber re-established. Detachment of the choroid also occurs occasionally after trephining for glaucoma. These cases usually do badly. Probably the trephine hole is too far back and the attachment of the ciliary body is permanently injured.

Sarcoma of the Choroid.—See p. 385.

Primary Affections of the Retina.

Primary Retinitis, in its most severe manifestations, is almost always the result of some general disease, and should, therefore, be properly regarded as a symptomatic disease. It is probable that slight inflammation of the retina may be caused by overuse or misuse of the eyes, but though some of the symptoms may be ascribed to it, ophthalmoscopic signs are wanting or so little marked as to be ambiguous. Thus, *hyperæmia* of the retina is often described, but can never be diagnosed with certainty in the absence of definite pathological signs in the retinal vessels (*vide infra*).

Retinitis in general gives rise to the following symptoms and signs, only some of which need be present in individual cases. There is usually some change in visual acuity. Rarely it is increased in the early stages; more commonly it is diminished throughout. There may be concentric diminution of the field of vision, or scotomata may be present corresponding with the areas specially affected. There may be metamorphopsia, micropsia, or macropsia (*vide* p. 315). The light sense is diminished, and photophobia may be present. Pain is almost invariably absent, though discomfort may be complained of.

The ophthalmoscopic signs may be diffuse or localised. There may be general œdema, manifesting itself as a faint, diffuse haze, obscuring details, so that the normal bright red appearance is replaced by a paler cloudiness, often with definite white streaks, especially along the course of the vessels; or there may be circumscribed areas of exudation. The latter appear as white spots, discrete or confluent, or yellowish plaques, varying in size. They are not pigmented and the edges are ill-defined, so that there is little danger of mistaking them for patches of choroidal atrophy. The blood vessels usually show marked changes. The veins are distended, often irregular, darker than normal, and tortuous;

the arteries are less altered, but the finer branches are also tortuous. Hæmorrhages are common, though they are not in themselves evidence of primary retinitis. When occurring in the superficial layers they are flame-shaped, with feathery edges, situated particularly along the course of the vessels; when in the deeper layers they are round, with better defined contours.

The optic nerve, being in anatomical and physiological continuity with the retina, often suffers with it; when this occurs to a marked extent the condition is called *neuro-retinitis*. The margin of the disc is then obscured and often shows a radial striation. The disc is red, and may be measurably swollen, though seldom, if ever, so much as in the condition known as choked disc.

Retinal atrophy, which follows severe retinitis, is shown by permanent whitish or yellow opacity, with diminution in the size of the vessels, which are often bordered by white lines. The optic disc may show all the signs of advanced atrophy (*vide* p. 366).

It has already been pointed out that retinitis is most frequently the symptom of an internal disease. It is therefore usually bilateral. The diseases most commonly causing the condition are syphilis, vascular disease, albuminuria, diabetes, and leucæmia.

The *treatment* of retinitis consists in giving the eyes complete rest and in combating the general disease which is the cause. All near work is forbidden and the eyes are protected with smoked glasses, or even confinement in a darkened room. Whether due to syphilis or not mercury and iodides are given with a view to aiding the absorption of exudates and restoring the transparency of the vitreous, if this is affected. Mercury is contraindicated in renal retinitis. These means are supplemented by purgatives, diaphoretics, and tonics.

The blood vessels of the retina are peculiarly subject to disease in retinitis, partly as a factor in the inflammation, but more prominently as a concomitant of general disturbance or disease. It will be well, therefore, to discuss the commoner forms of vascular disease of the retina before passing on to describe the chief types of retinitis in greater detail.

Vascular Disorders of the Retina.

Anæmia may be part of general anæmia or due to local causes. It may be sudden or slow in onset. Sudden anæmia is seen in embolism of the central artery of the retina (*q.v.*)

and in quinine amblyopia (*q.v.*). Ophthalmoscopically there is great attenuation of the retinal vessels and the optic disc is pale. Spasmic constriction of the retinal arteries has been described in migraine (*vide* p. 378), but it is doubtful if it occurs. I have seen one case of spasm of the retinal arteries in one eye, giving rise to the symptoms of embolism of the retinal arteries: it passed off during the actual ophthalmoscopic examination. Anæmia of slow onset is seen in atrophy of the retina from any cause, such as previous retinitis, and in disease of the vessel walls as part of a general vasculitis. In both cases the vessels are attenuated, and sometimes in the former and commonly in the latter the walls become thickened and visible as white lines bordering the red blood stream; eventually the vessels may be transformed into white strands or may even disappear.

Embolism of the Central Artery of the Retina causes sudden and complete retinal anæmia. The eye, usually the left, becomes suddenly quite blind. Examination of the fundus reveals a very typical picture (Plate XII., Fig. 1). The larger arteries are reduced to threads, the smaller ones are invisible. The veins are little altered except on the disc, where they are contracted. Within a few hours the retina loses its transparency, becoming opaque milky-white, especially in the neighbourhood of the disc and macula. Owing to the opacity of the retina the outlines of the disc, which is abnormally pale, are obscured. At the fovea centralis, where the retina is extremely thin, the red reflex from the choroid is visible. It appears as a round "cherry-red" spot, presenting a strong contrast to the cloudy white background. The peculiar tint of the spot is due to this contrast. In the majority of cases there is no hæmorrhage here, as was once thought, though hæmorrhages here and in the immediate neighbourhood do occur rarely. The contrast sometimes brings into relief minute blood vessels near the macula which are otherwise invisible.

Sometimes the obstruction to the blood flow is not complete, or the flow may be partially restored in the course of a few days. Another peculiar phenomenon may be observed or may be induced by gentle pressure upon the globe. In some of the vessels, usually veins, the column of blood is broken up into red beads separated by clear interspaces. The beads move in a jerky fashion through the vessels, sometimes in the normal direction of blood flow, sometimes in the opposite direction. If the veins are easily emptied of blood or arterial pulsation is

PLATE XII. (*To face p.* 330.)

Fig. 1.—Embolism of the central artery of the retina. Note the œdema of the retina at the posterior pole and the "cherry-red" spot at the macula. The vessels are reduced in size. Subsequently the disc became atrophied, and the vessels still smaller.

Fig. 2.—Thrombosis of the central retinal vein. Note the distension, convolution, and irregularity of contour of the veins, and the retinal hæmorrhages and exudates.



FIG. 1.

FIG. 2.

produced by slight pressure on the eye-ball it is evidence of incomplete blockage.

The retinal œdema or possibly coagulation necrosis takes several weeks to clear up. The membrane regains its transparency, but is completely atrophic. The vessels are contracted or reduced to white threads. The disc is atrophic. If there have been hæmorrhages spots of degeneration replace them, and cholesterin crystals and pigmented spots may be seen in the papillo-macular region.

In some cases a certain degree of central vision persists in



FIG. 191.—"Silver-wire" arteries, degeneration of the walls of a vein, and white spots of degeneration.

spite of apparent complete occlusion of the central artery. It is due to the presence of cilio-retinal arteries (*vide* p. 119) which, when present, always supply the macular region, and naturally escape occlusion. The remainder of the field of vision is lost. In rare cases a cilio-retinal artery alone becomes blocked.

In other cases the embolus is arrested in a branch of the central artery. The area supplied by this branch is then affected alone. In the early stages the corresponding scotoma is usually somewhat indefinite, but later it settles down to a permanent sector-shaped defect.

When perception of light is lost the pupil is large and the

direct reaction to light fails. In some cases faint perception of light may be regained in the temporal region of the field : it is possibly due to partial establishment of collateral circulation. The intraocular tension is normal, as might be anticipated.

Embolism of the central artery, like cerebral embolism, occurs with mitral stenosis, especially if there has been fresh endocarditis. This factor is, however, very frequently absent, and it is probable that most cases are really due to thrombosis. There has been endarteritis, due to general arteriosclerosis from



Fig. 192.—Perivascular changes in the retinal vessels.

nephritis, &c., and the already narrowed lumen of the vessels has become suddenly occluded. The onset may be less rapid in such cases and preceded by premonitory symptoms, such as obscuration of vision, &c. Other cases render it probable that the condition may be due simply to spasm of the walls of the artery, thus accounting for some remarkable cures.

The condition has been observed at ages varying from fifteen to eighty. Rare cases of simultaneous bilateral blocking of the central arteries have been recorded.

Treatment is seldom of any avail, but attempts should be

made to drive the obstruction on into a less important branch if the case is seen early. Inhalation of amyl nitrite, massage of the globe, and paracentesis of the cornea have been employed for this purpose; such measures must be adopted without delay.

Amaurotic Family Idiocy shows ophthalmoscopic signs resembling those of embolism of the central artery, but of quite different origin. The disease occurs almost if not quite invariably in Jewish children, and commences during the first year of life. Several members of a family may be affected. The apparently



FIG. 193.—Retinal vein with enormously thickened wall and narrow lumen. (Coats.) The perivascular lymph space is dilated.

healthy child becomes gradually blind, with muscular weakness and wasting, and mental apathy passing into idiocy. Death follows in from one to two years. Ophthalmoscopically there is a round brilliantly white area at the macula, fading off peripherally into the normal fundus. In the centre of the patch is a brownish-red, circular spot at the fovea. In the later stages there is optic atrophy. It is always bilateral. The disease is a primary neuronic degeneration of the whole of the central nervous system, probably of toxic origin. The ganglion cells of the retina participate in the degeneration.

Maculo-cerebral Family Degeneration has some points of resemblance to amaurotic family idiocy, but should be carefully distinguished from it. It is a familial disease, occurring in other

than Jewish children, and commencing at a later age, usually at about six or eight years. Defective vision, with central scotoma, is accompanied by weak intellect. Ophthalmoscopically the discs are pale and the vessels small. At the macula there are yellowish grey spots and granular pigmentation. Similar macular degeneration has been seen as an hereditary disease without cerebral deterioration.

Degenerative Changes in the Retinal Vessels, apart from their interest as a local manifestation of disease, are of the utmost importance in general prognosis. They may be the first evidence of arteriosclerosis, and particularly of disease of the cerebral vessels, pointing to the danger of cerebral hæmorrhage, and indicating lines of treatment which may prolong life. Undue tortuosity of the vessels is of no significance unless accompanied by other abnormalities, such as irregularity in the size and breadth of the arteries, so that stretches of the vessel are much constricted alternating with normal or somewhat dilated portions. These changes in the lumen are due to endothelial proliferation in the intima. Minute miliary aneurysms are seen in rare cases. The normal light reflex from the vessel walls is often unusually bright and broad, especially in vessels at some distance from the disc (" silver-wire " arteries) (Fig. 191). Under normal conditions it is possible to see a vein through an artery at a point of crossing; in arteriosclerosis the artery loses its translucency so that the vein is obscured. Moreover, the artery exerts an abnormal pressure on the veins so that the blood flow is obstructed; the vein seems therefore to stop at the crossing and is more distended on the distal side than on the side towards the disc. Sometimes the vein appears to be pushed aside by the crossing artery; in severe cases the vein, whether crossing above or below the artery, is diverted so that it crosses at right angles, the shortest possible route. The veins may also exhibit a beaded appearance, with alternate constrictions and dilatations. More pronounced changes make the walls of the vessels visible, so that the blood column, often narrowed, is bounded by white lines, the thickened fibrous walls (Fig. 193); this may affect both arteries and veins, usually only individual vessels, in a portion of their course (Figs. 191, 192).

The changes indicated lead to increased permeability of the walls, and increased internal pressure, due to general disease, supplements this defect, and causes undue extravasation of lymph, and even hæmorrhage. Œdema of the retina thus arises, manifesting itself as a greyish opacity around the disc,

or in spots along the course of the vessels. Hæmorrhages occur as linear striated extravasations along the vessels, or as round spots scattered over the fundus.

These changes occur most frequently in elderly people and are seldom entirely absent in the aged. They are specially pronounced in cases of chronic nephritis, syphilis, some forms of poisoning, especially by lead and probably by auto-intoxication, notably that of intestinal origin. They are frequently associated with high blood pressure, and always indicate the necessity for exhaustive examination of the circulatory and excretory systems. They are more frequent in women than men, though the former seem to be more tolerant of high blood pressure than the latter (Foster Moore). They may be very marked and extensive in quite young people as the result of congenital syphilis ; rarely as an hereditary condition without syphilitic taint. Extensive disease of the retinal vessels, with much diminution in their calibre, so interferes with the nutrition of the retina that consecutive atrophy of the optic nerve not infrequently follows. The ophthalmoscopic appearances of the vessels and disc then closely resemble those found in advanced cases of retinitis pigmentosa. This condition may be accompanied by groups of sharply defined small white spots in the retina and even a fan- or star-shaped figure at the macula (*arterio-sclerotic retinitis*, Foster Moore). It is frequently unilateral, thus differing from renal retinitis.

Angioid Streaks. Dark brown or pigmented streaks, which anastomose with each other and resemble blood vessels in distribution, are sometimes seen ophthalmoscopically in retinæ which are undergoing degenerative changes. They differ in distribution from any normal set of vessels, are usually situated near the disc, at a deeper level than the retinal vessels, and are very irregular in contour. They are probably due to proliferation of new vessels into scar tissues (W. T. Lister).

Hyperæmia may be arterial or venous. Arterial hyperæmia, characterised by fulness and tortuosity of the arteries, accompanies not only inflammation of the retina, but also inflammation of neighbouring structures, especially the uveal tract. Venous hyperæmia, characterised by dilatation and great tortuosity of the veins, is the result of impeded return of blood to the heart. It may be due to general venous congestion, seen in its most extreme form in congenital malformation of the heart (*cyanosis retinæ*), or to local causes. The latter most commonly affect the veins in the porus opticus, as is seen

in moderate degree in glaucoma and optic neuritis, and in extreme form in thrombosis of the central vein of the retina. Increased intraorbital pressure, as from a tumour, may also impede the exit of blood from the eye. The veins are much enlarged and dark in colour in polycythæmia.

Hæmorrhages from the retinal vessels may be pre retinal or intra-retinal. *Pre-retinal or subhyaloid hæmorrhages* are extravasations of blood between the retina and the hyaloid membrane of the vitreous. They always occur in the neighbourhood of the macula, and are usually large. They are round at first, but quickly become hemispherical, the upper margin being straight; this is due to the effect of gravity (Plate XIII., Fig. 1). Occasionally two such hæmorrhages may be seen in the same eye. Retinal vessels are hidden from view in the affected area. The upper layers become lighter in colour, generally attributed to the sinking of the red corpuscles. The blood gradually becomes absorbed, usually in a patchy manner, but finally disappears, though numerous cholesterin crystals may often be left as bright glistening spots. Vision is restored, but recurrences are not uncommon, and other complications of vascular origin may modify the otherwise favourable prognosis.

Intra-retinal hæmorrhages, as already mentioned, are striate or flame-shaped when situated in the nerve fibre layer, rounded or irregular when in the deeper layers or between the retina and choroid. Intra-retinal hæmorrhages are absorbed very slowly, gradually becoming white, rarely pigmented.

Retinal hæmorrhages are due to many causes. Most frequently the vessel walls are weakened by general disease, which may be a vascular degeneration due to age or to altered composition of the blood, as in pernicious anæmia, leucæmia, scurvy, purpura, sepsis, albuminuria, diabetes, phosphorus poisoning, &c. Any of the causes leading to retinal hyperæmia may give rise, secondarily, to hæmorrhages. They may be due to pressure during birth in new-born infants, and are probably responsible for some so-called congenital retinal defects, *e.g.*, white and pigmented spots of atrophy at the macula and elsewhere. Similarly they occur in cases of severe compression of the thorax or neck in older people. Whooping cough may lead to retinal as well as the more common conjunctival hæmorrhages. Traumatism, *e.g.*, severe contusions and wounds, gunshot wounds, of the eye and of the orbit without direct injury to the eye, is another cause, and is often responsible for very extensive extravasations of blood, which

PLATE XIII. (*To face p.* 336.)

Fig. 1.—Subhyaloid hæmorrhage, 96 hours after probable time of onset. Four months later the patient was reported to have quite recovered his sight.

Fig. 2.—"Retinitis proliferans," showing the origin of the new-formed fibrous tissue in the favourite situation at or near the disc. The bands stretch forwards into the vitreous, a feature which cannot be adequately represented in a drawing.

FIG. 1.



FIG. 2.

may burst through into the vitreous. To this category belong the post-operative hæmorrhages. When the globe is opened in performing cataract extraction, or, still more, iridectomy for glaucoma, the normal or raised intraocular tension is suddenly reduced to zero. The intraocular vessels then dilate, and great



Fig. 194.—Thrombosis of a branch of the central retinal vein.

strain is thrown upon their walls. If these are diseased, there is much danger of their rupturing. Considering the age and condition of these patients, it is surprising that hæmorrhage is not more frequent.

Minute hæmorrhages, unless in the macular region, may cause little obscuration of vision. Subhyaloid hæmorrhage

usually abolishes central vision temporarily: it takes some weeks to clear up, the length of time varying with the size of the extravasation. The smaller spots may remain unaltered for months, though this is only apparent in some cases, old spots clearing up and being replaced by new ones. They may be absorbed without leaving any trace.

Venous Thrombosis may affect the central vein of the retina (Plate XII., Fig. 2) or one of its branches (Fig. 194). In the former case the obstruction is always just behind the lamina cribrosa. All the veins of the retina become enormously engorged with blood, and extremely tortuous. Blood escapes at innumerable spots, so that the retina is covered with hæmorrhages. Sight is much impaired, though not so rapidly as in obstruction of the central artery, but recurrent extravasations finally destroy it entirely. When a single branch is blocked the œdema and hæmorrhages are limited to the area supplied by the vein. In these cases the defect in vision is not accurately sectorial, as in the case of a branch of the artery, and the macular region generally escapes. The affected retina becomes atrophic, with fine pigmentary changes. Secondary glaucoma ensues in two to three months in a considerable number of the cases, probably owing to the increased albuminous constituents of the intraocular lymph. It does not occur when only a tributary is involved. In many cases branches of tortuous new vessels are formed upon the disc (Fig. 195); in others a collateral circulation is effected by similar tortuous new vessels in the retina. Vascular disease and hæmorrhages are not infrequently present in the other eye, and bilateral thrombosis of the central vein sometimes occurs.



Fig. 195. — New-formed vessels (Foster Moore).

The patients are usually elderly, with cardiac or vascular disease, often induced by nephritis; the blood pressure is usually high, and there is often albuminuria. The obscuration of vision is almost always noticed on waking in the morning. Probably the low blood pressure and sluggish flow during

sleep allows thrombosis to occur in the vessel where it is constricted normally at the lamina cribrosa and pathologically by endovascular changes. Thrombosis may, however, be due to local causes, such as orbital cellulitis, following facial erysipelas, &c. It occasionally occurs in young people following a febrile attack, and is then probably due to an infective phlebitis.

Treatment.—No treatment is of avail. Atropin should be avoided, as tending to induce glaucoma. Local treatment is indicated only if the eye becomes painful, when it should be removed. The condition is to be regarded as a danger signal, and constitutional investigation and treatment should be carried out assiduously.

Special Forms of Retinitis.

Syphilitic Retinitis. Syphilis is one of the commonest causes of retinitis, but it is usually a secondary retinitis, accompanying disease of the choroid. It however occurs as a primary retinitis also, and in this form syphilitic endarteritis is a prominent sign. There are dust-like opacities in the vitreous, especially in the posterior part; the retina is cloudy, particularly in the neighbourhood of the disc, which may be hyperæmic. White spots may be seen in the macular region, and yellowish or white spots, often bounded by pigment, at the periphery of the fundus. The vessels may be degenerated (*vide* p. 334), with whitish exudations along their course; hæmorrhages are rare.

Every transition is seen to a condition much resembling choroido-retinitis pigmentosa (*vide* p. 323), but seldom showing such a characteristic distribution of the pigment.

The subjective symptoms are defective central vision, night-blindness, irregular and concentric contraction of the field, with or without central, paracentral, or ring scotomata, and metamorphopsia.

In most cases the amount of organisation which takes place at the sites of the inflammatory deposits of syphilitic retinitis is very small, but in some cases there is a well-marked tendency to the formation of new blood vessels. These may not be limited to the retina itself, but may extend into the vitreous, forming convoluted coils. They are held together by a minimum of delicate connective tissue (cf. Retinitis proliferans). They are commonest on or near the disc.

In acquired syphilis the disease usually occurs one to two years after infection; usually both eyes are involved, but not

with simultaneous onset. As a rare manifestation the macular area is alone affected, showing a grey or yellow deposit, or numerous small yellow spots and dots of pigment. This form shows a great tendency to relapse.

Retinitis is not uncommon in congenital syphilis. Such patients often show a dusty or peppery discrete pigmentation of the retina at the periphery, associated with a tigroid condition of the fundus in this situation. It is only distinguishable from what is often seen as a normal condition by the greater aggregation of the pigment. There may be thickly strewn black and white spots, like a mixture of pepper and salt. In more definite forms there are yellowish red and black spots at the periphery (anterior retinitis), a condition often seen in interstitial keratitis (*vide* p. 225), or larger grey or white patches may be seen, or the condition observed in the acquired form may be fully developed.

If the diagnosis is doubtful the Wassermann test should be applied.

Treatment.—A prolonged course of anti-syphilitic treatment is indicated. Dark glasses should be worn and the eyes rested.

Albuminuric Neuro-retinitis, in its most typical form, presents an ophthalmoscopic picture which is almost pathognomonic, being simulated only in some cases of intracranial tumour (Plate XIV.). In addition to the general signs of retinitis—haziness of the retina and disc, hyperæmia and hæmorrhages—the distinguishing feature is the presence of brilliant white spots and patches in the retina. The earlier deposits are cloudy, with soft edges (" cotton-wool " patches) ; the later brighter, more sharply defined and punctate. The disc is surrounded by large white patches or by a continuous " snow-bank." Around the macula are smaller dots or round patches, also silvery white. Radiating from the fovea are spokes of white dots or fine lines, forming a star-shaped figure, which is extremely characteristic. The fovea itself escapes, and the star is often incomplete in some direction. The vessels generally show very definite degenerative changes (*vide* p. 334). In some cases, especially in the albuminuric retinitis of pregnancy, a flat detachment of the retina occurs, almost certainly due to the retina being raised from the choroid by exudates. The detachment is usually bilateral and involves the lower part of the fundus. Unlike most detachments of the retina, these frequently disappear, the exudates being absorbed.

Albuminuric retinitis by no means always displays the typical picture. Often there is a neuro-retinitis which exhibits

PLATE XIV. (*To face p.* 340.)

Albuminuric neuro-retinitis. Note the papillitis and macular star-shaped figure, specially marked in Fig. 1, and the masses of white exudate in the retina. The patient from whom Fig. 2 was taken, a man, *æt.* 21, died three months after the drawing was made.

FIG. 1.

FIG. 2.

no characteristic features specially associated with nephritis. In these cases there is moderate swelling of the disc—seldom so much as in the choked disc of intracranial disease—more or less widespread œdema and hæmorrhages. Irregularly scattered bright white spots and patches may be present or wanting. *The urine should be examined in every case of retinitis.*

The white spots of albuminuric retinitis are chiefly composed of exudates, which are often fibrinous (Fig. 196), later becoming hyaline. They are situated particularly in the inner layers, but may be present in all the layers. There is some leucocytic infiltration, and peculiar swollen nucleated structures—cytoid bodies—are found in the nerve fibre layer: they are probably varicose nerve fibres. The exudates and necrotic retinal elements undergo fatty degeneration. The peculiar arrangement of the spots in the macula is attributed to the radial disposition of Müller's fibres, but it is probable that when the œdema is considerable there may be actual radial folds in the retina, within which the exudates accumulate. The changes in the blood vessels are those common to vasculitis in other small vessels, consisting of endo- and perivasculitis and hyaline degeneration of the walls.



Fig. 196.—Albuminuric retinitis. Masses of fibrinous exudate in the retina.

There is no constant relationship between the retinitis and the vascular disease. The vessels may be quite normal, *e.g.*, in some puerperal cases and in the rare cases occurring with acute nephritis. In the cases with chronic interstitial nephritis, *i.e.*, the great majority, the vessels are much diseased. The retinitis is, therefore, not directly due to the vascular disease, but probably to toxins circulating in the blood stream.

Albuminuric retinitis occurs in about one-third of cases of nephritis and may occur in all forms of nephritis, including scarlatinal, puerperal and "trench" nephritis, but in by far the greatest number of cases the disease is chronic interstitial nephritis, and it is very rare in ordinary acute nephritis. The last-mentioned fact accounts for the small quantity or even total absence of albumin in the urine in some cases. Though the degree of retinitis bears no fixed relationship to the nature

or severity of the renal mischief, yet in all cases its presence is of grave significance. The retinal changes may be the first evidence of renal disease, hence the extreme importance of their discovery, which is accentuated by the fact that the majority of hospital patients die in from six months to two years. Under more favourable conditions patients may survive five years or even longer. The prognosis is equally grave in children. The risk to life is much less in the scarlatinal and puerperal cases. In the latter the prognosis is worse the earlier the onset of the retinitis, but fortunately it usually commences in the later months of pregnancy. The artificial induction of abortion is indicated, and usually has a prompt beneficial effect; vision, however, is usually permanently impaired, the degree depending upon the duration of the retinitis. The "cotton-wool" patches clear up first, the glistening macular spots more slowly. Partial optic atrophy and slight retinal changes, such as white or pigmented spots at the macula, follow. The disease does not always occur at the first pregnancy; but may after one attack recur at subsequent pregnancies, though by no means always. The patient should, however, be warned of the danger. The retinal changes in puerperal cases are usually severe, and detachment of the retina occurs more frequently in these than in other cases.

Diminution of visual acuity is commonly the only symptom complained of, and, as mentioned, may lead to the discovery of the renal disease. Generally the history of severe headaches can be elicited, and the blood pressure is high, usually about 200 mm. Hg. The condition is almost always bilateral. It very rarely causes complete blindness. It occurs more often in men than in women, owing to the greater incidence of nephritis in men; in children it is commoner in girls than in boys. It is commonest between thirty and sixty years of age, especially in the last decade; but it is probably more frequently associated with the parenchymatous nephritis of children, which is generally syphilitic, than has been hitherto thought (Nettleship).

It is important to note that transient blindness may occur in the course of nephritis, especially associated with uræmia. In these cases the retina shows no abnormality, or at most changes which may occur independently of the disease. This *uræmic amaurosis* is distinguished in being sudden total blindness, whereas the defective vision of albuminuric neuro-retinitis is slower in onset and never complete. Sight usually returns in one or two days (*vide* p. 377).

Diagnosis.—The typical picture of albuminuric retinitis may be nearly simulated in some cases of increased intracranial pressure, particularly when due to cerebral tumours occurring in children (*cide* p. 358). It may occur without any definite signs of nephritis, and somewhat similar appearances are met with in glycosuria and leucæmia. Rarely a star at the macula, with or without slight papillitis, has been met



Fig. 197.—Diabetic retinitis.

with in young persons with anæmia or chlorosis, or without discoverable cause.

The *treatment* of these conditions is purely constitutional.

Diabetic Retinitis. Retinitis is a relatively rare complication of diabetes, occurs in the late stages, and in elderly people; it is not improbable that it is frequently missed owing to the peripheral position of the lesions, opacities of the lens, &c. It is generally, but not always, bilateral. Irregularly scattered small, bright white spots around the macular region are the

commonest manifestation. The snowy patches and stellate arrangement at the macula are usually absent, but it must be remembered that albuminuria is a frequent concomitant of the late stages of diabetes, and all the characteristics of albuminuric retinitis may occur. The white spots may coalesce into larger plaques with crenated edges, which indicate their mode of formation. Punctate hæmorrhages are freely scattered over the fundus: they are more often round and deeply seated than linear and superficial, as in renal retinitis. The vessels and optic disc are generally normal, as well as the remainder of the retina.

The prognosis depends upon the severity of the constitutional condition; 60 per cent. live more than two years.

Diabetic retinitis cannot be due to sugar, di-acetic acid or acetone circulating in the blood, since it does not occur in children in whom these products of perverted metabolism are most markedly present.

A peculiar feature sometimes met with in diabetes is *lipæmia*. It occurs especially in young patients, and the prognosis is grave. The ophthalmoscopic appearances are then striking, the retinal vessels containing fluid which looks like milk. The arteries are pale reddish, the veins having a slight violet tint. The general fundus has much the normal coloration.

Hæmorrhagic Retinitis is a term loosely applied to those forms of retinitis accompanied by hæmorrhages, for hæmorrhages may occur, as has already been seen, in the absence of all signs of inflammation of the retina.

Leucæmic Retinitis.—When retinitis occurs in the course of leucæmia the ophthalmoscopic appearances are characteristic. The fundus is pale and orange-coloured. The veins are dilated and tortuous, often with white lines along them, and are bright red, not dark; the arteries are small and pale yellowish red. Very typical are white spots and patches surrounded by a red rim; they consist of leucocytes surrounded by red corpuscles. These are present only rarely, and are also found in pernicious anæmia. They are most common in the periphery of the retina. In some cases the ordinary picture of hæmorrhagic retinitis is seen. In every doubtful case the blood should be examined.

Septic Retinitis (Roth). Apart from endogenous retinitis (*vide* p. 309), metastatic retinitis may manifest itself in the form of small round or oval white spots near the disc, often accompanied by hæmorrhages. This form occurs in puerperal and other forms of septicæmia. Usually both eyes are affected.

" Retinitis Proliferans."—When hæmorrhage occurs into the vitreous the blood clot is usually almost completely absorbed. This is probably due to the absence of fibroblasts in the vitreous and their scantiness in the retina, for the retinal connective tissue consists of neuroglia, an epiblastic structure which probably takes no part in fibrous tissue formation. The only mesoblastic tissue in the retina is that forming and surrounding the retinal blood vessels. In some cases, however, blood clot may organise, giving rise to masses of fibrous tissue in the vitreous, vascularised by newly formed blood vessels derived from the retinal system (Plate XIII., Fig. 2). This condition is known as " retinitis proliferans." The tissue is most commonly situated near the disc and the vessels spring from this neighbourhood, probably owing to the fact that there is more mesoblastic tissue here than in other parts of the fundus.

There seems to be some special factor necessary to stimulate organisation, and it is found in some general diathesis. In nearly all these cases there is either a history of syphilis or the patient is suffering from nephritis, diabetes, or some other form of toxæmia, the origin of which may be obscure, as in some cases of recurrent hæmorrhage (*vide* p. 308).

The amount of fibrous tissue varies from the most delicate strands and films supporting new-formed blood vessels—specially common in syphilitic cases (*vide* p. 339)—to dense bands and membranes stretching far forwards into the vitreous and hiding the fundus. Most extensive proliferations were met with during the war in cases of rupture of both choroid and retina (*vide* p. 422). The bands are often attached to the retina at their apices, and as the fibrous tissue contracts after formation the retina may be pulled up and detached.

Vision is generally greatly impaired and often lost, usually from detachment of the retina. No treatment is of much avail, but hypodermic injections of fibrolysin may be tried, combined with constitutional régime.

Retinitis Circinata.—Retinal hæmorrhage may give rise to yet another condition in rare cases, generally elderly women. In retinitis circinata there is a girdle of bright white patches with crenated borders around the macula. The diameter of the girdle, which is usually an imperfect circle or ellipse, or horse-shoe-shaped, open towards the temporal side, is generally considerably greater than a papilla diameter, and follows the larger macular branches of the superior and inferior temporal vessels. The vessels pass over the spots. The macula shows yellowish-white areas, slight

pigmentation and often hæmorrhages. The patches develop slowly and are usually well advanced before noticed. The disease is unilateral in about half the cases: exudative retinitis has been observed in the other eye (Coats). Central vision is much reduced, but the field remains full. The patches sometimes disappear slowly and vision improves. Mercury and iodides seem to afford the best chance of amelioration.

Exudative Retinitis. (*Syn.*— *Massive Exudation in the Retina* (Coats)) resembles conglomerate tubercle in its ophthalmoscopic signs. There is usually a large raised yellowish white area or several smaller areas posterior to the vessels. The vessels often show gross degenerative changes, and there is sometimes arterio-venous communication, with enormous dilatation of the veins. The patients are usually boys, otherwise apparently healthy. Detachment of the retina, cataract or glaucoma may occur in the late stages. There is always evidence microscopically of hæmorrhage between the retina and choroid and in the deep layers of the retina; the choroid is at first healthy.

Purulent Retinitis.—This is most commonly due to a septic perforating wound, being then a precursor of panophthalmitis (*vide* p. 423). Rarely it is a metastatic condition, occurring in pyæmia, and probably starting in a septic embolus. In the early stages there is severe retinitis with hæmorrhages. Suppuration rapidly follows, involving the vitreous, so that a yellow reflex is obtained. The condition often passes into panophthalmitis, but less commonly than in cases of exogenous infection. Pyogenic organisms are attenuated in the blood stream and tissues, so that the process may subside with the restoration of useful vision (*vide* p. 319).

Retinitis from Bright Light, which might be termed *photo-retinitis*, occurs after exposure of the unprotected eyes to bright sunlight, as in looking at an eclipse of the sun with unprotected eyes ("eclipse blindness"), or the electric light, as in the intense flash of the short-circuiting of a strong current. The relative parts played by the different rays of the spectrum are not fully understood. Only a few, relatively innocuous, ultra-violet rays reach the retina. On the other hand, practically all the visible rays and many infra-red rays pass unimpeded to the retina (*vide* p. 19). Much of this radiation is absorbed by the pigment epithelium, and it is probable that the pathological changes are produced by the resultant heating effect. It is, in fact, a burn of the retina.

The symptoms are persistence of the after-image, passing on later into a positive scotoma, and metamorphopsia.

Ophthalmoscopically there may be no signs at first, or a pale spot is seen at the fovea with a brownish red ring round it. Later there are usually deposits of pigment and small grey punctate spots around the fovea. Prognosis must be guarded, since, though improvement often occurs, some defect usually remains, and the positive scotoma may persist permanently.

The *treatment* is that of retinitis in general. Smoked glasses should be used, or Crookes's glasses, *i.e.*, such as cut off the ultra-violet rays.

Toxic Amblyopia.—Tobacco, alcohol, quinine, filix mas, carbon disulphide, stramonium, cannabis indica and other poisons sometimes produce defective vision, which is then known as toxic amblyopia. Tobacco and alcohol amblyopias, which are most frequently met with, are usually described as forms of retrobulbar neuritis (*q.v.*), but experimental and pathological evidence tends to show that the condition is primarily retinal in these cases.

Tobacco amblyopia results from the excessive use of tobacco, either by smoking or chewing; and also occasionally from the absorption of dust in tobacco factories. Smokers of shag and tobacco mixtures suffer most. In nearly all cases there is also over-indulgence in alcohol. It is known that alcohol alone may produce toxic amblyopia, and cases in which tobacco caused the disease in total abstainers from alcohol have been published. The patients may have smoked excessively for years with impunity, the attack coinciding with some intercurrent cause of debility, digestive disturbance, &c. They are usually thirty-five to fifty years of age.

The patient complains of increasing fogginess of vision, which is usually least marked in the evening and in a dull light. Central vision is greatly diminished. The field of vision is found to be full, but there is a central colour scotoma for red and green. This is usually small, horizontally oval between the fixation and blind spots, but in rare cases may extend to the limits of the red field, or even be absolute. Both eyes are about equally affected. Ophthalmoscopically there may be congestive haze of the edges of the optic disc followed by undue pallor of the temporal side of the disc, but the changes are usually slight.

The course is chronic, and the prognosis is good if the toxic agents are discontinued absolutely. Recurrence is very rarely seen. Very rarely optic atrophy may result, but it is doubtful if these are uncomplicated cases.

The disease is probably due to poisoning of the ganglion

cells of the retina. In experimental cases, and in one case examined in man, the cells showed vacuolation and breaking up of the Nissl granules. This leads to degeneration of the nerve fibres, demonstrable only after they have obtained their medullary sheaths, *i.e.*, behind the lamina cribrosa. The degeneration is found to be limited to the papillo-macular bundle (*vide* p. 69). The ganglion cells of the fovea and macular region are the most highly differentiated and are liable to suffer first and most severely in any toxic condition. The degeneration is therefore a wedge-shaped area on the temporal side of the nerve immediately behind the globe, but becomes a circular central area more posteriorly (Fig. 58). This degeneration was discovered early in the history of the disease, and combined with the clinical similarity to undoubted cases of retrobulbar neuritis led to the conclusion that it was the primary seat of the disease. Nicotin is generally regarded as the toxic agent, but it is much more probable that it is one of the more volatile decomposition products of nicotin, *e.g.*, collidin or lutidin.

The amblyopia produced by diabetes, carbon disulphide, and iodoform resembles that of tobacco. Diabetics appear to be specially susceptible to tobacco.

Treatment consists in total abstinence from tobacco and alcohol. It should be combined with tonic treatment, especially the administration of strychnine. Iodide of potassium may be given, and copious draughts of water, combined with exercise, have been recommended. All patients with central scotomata, other than those due to total macular degeneration, should abstain from tobacco and alcohol.

Quinine amblyopia differs in some striking characteristics from tobacco amblyopia. Here total blindness (amaurosis) follows the use of the drug, even in such small doses as 12 grains in susceptible persons; 40 grains is the maximum amount of sulphate of quinine which should be given within twenty-four hours (Yarr). The largest doses are usually taken for malaria, but quinine is also used as an abortifacient. The pupils are dilated and immobile. Deafness and tinnitus aurium are present. Ophthalmoscopically the retinal vessels are extremely contracted and the disc is very pale. In less marked cases or at a later stage the fields of vision are much contracted. The fields gradually widen out, but do not regain their normal limits. Central vision may be completely restored. The discs may remain pale for years or become normal. Occasionally blindness is permanent and optic

atrophy ensues. The same condition may follow administration of ethyl-hydrocuprein (optochin) for pneumonia.

Salicylic acid and *salicylates* occasionally produce an amblyopia of the same type and with the same ophthalmoscopic features as that of quinine, but not so severe.

Treatment consists in discontinuing the drug, administering amyl nitrite or nitro-glycerine, supplemented by strychnine and digitalis.

The amblyopias produced by *methyl alcohol*, *arsenic*, *lead*, *nitro-* and *dinitro-benzol*, and *filix mas* differ from those of retrobulbar type in the more serious optic atrophy which generally ensues. There is probably always a stage at which a central scotoma is present, but it is often missed.

Methyl alcohol poisoning from drinking wood-alcohol is common in America, and used to occur in England from drinking methylated spirit before it was intentionally adulterated. Nausea, headache, giddiness, &c., are followed by coma. If the patient survives vision very rapidly fails, passing through the stage of contracted fields and absolute central scotoma to blindness. Vision may improve, but usually again relapses, becoming gradually abolished by progressive optic atrophy. Rarely restoration is complete. Ophthalmoscopically there may be blurring of the edges of the discs and diminished size of the vessels in the early stages. Later there are all the signs of optic atrophy, usually of the primary type (*vide* p. 366).

Arsenic is specially liable to cause optic atrophy, usually total, when administered in the form of trivalent benzol-ring compounds such as atoxyl or soamin, arsacetin, hectine, &c. These were used for attacking the trypanosome of sleeping sickness, but have now been abandoned. The salvarsan group have the arsenic in pentavalent combination and are less toxic: no cases of optic atrophy have been reported from their use.

Lead poisoning is rarely seen since precautions have been taken to eliminate salts of the metal from pottery glazes, &c. The ocular signs are optic neuritis or optic atrophy, which may be primary or post-neuritic. Some cases have retinitis, which may be due directly to lead or of albuminuric type, secondary to lead nephritis.

Filix mas, used as a helminthetic, may cause amblyopia in excessive doses, especially if given with castor oil. The ophthalmoscopic picture is said to resemble that of quinine amblyopia. Later, optic atrophy supervenes. I have seen a case in which a drachm of extract of male fern was ordered three times a day and was taken for ten days. There was total optic atrophy in one eye and partial atrophy with much contraction of the field in the other.

Detachment of the Retina (*Syns.—Ablatio retinæ, Amotio retinæ*).—The retina may become separated from the choroid by being pulled up from within or pushed up from without. The simplest example of the former mode is in the late stages of plastic cyclitis when the strands of connective tissue which become attached to the retina contract during the process of organisation. The simplest example of the second mode is separation by means of a choroidal hæmorrhage, such as may occur from a blow or disease of the vessels.

Clinically detachment of the retina is observed most commonly in three conditions, viz., after a blow, in high myopia, and in sarcoma of the choroid ; to these must be added a not inconsiderable number of cases in which no cause can be assigned. The exact mechanism of detachment in these cases is by no means completely understood ; indeed, detachment of the retina is still one of the difficult problems of pathology.

The patient usually complains that there is a cloud in front of one eye, so that parts of objects, usually the upper or lower parts, are not seen. In other words, there is a positive scotoma, as is confirmed by making a chart of the field of vision. Usually the scotoma corresponding to the detached area is absolute, but in shallow detachments some vision may persist, sufficient nourishment being afforded to the retina from the subretinal fluid. There is then generally a relative scotoma for colours, and acquired colour-blindness of the tritanopic type (*vide* p. 381) is not uncommon. As a rule central vision is intact at first, but all detachments of the retina tend in time to be complete ; when the macular region becomes affected central vision is lost, and when the detachment is total perception of light is lost. The first symptom observed sometimes is transient flashes of light (photopsiæ), due to slight movements which irritate the neuro-epithelium. They should always be regarded with serious attention, but not infrequently occur, especially in myopic eyes, without being followed by detachment.

A small detachment causes much less definite signs. Some obscuration of vision is noticed, but the diagnosis can only be arrived at by careful examination of the fundus and of the field of vision. In sarcoma of the choroid the detachment may be very small and in any position, thus differing from the other forms—so-called *simple detachment*—in which it is generally larger, though often shallow, and confined to the lower parts of the fundus. Simple detachment often commences in the upper part of the fundus, but after a variable

PLATE XV.

PLATE XV. (*To face p.* 351.)

Simple detachment of the retina. The drawing is diagrammatic, since the disc and the highly hypermetropic detached retina cannot be in accurate focus at the same time. Note the change in colour and the convolution of the retinal vessels over the detached part. The red spot near the macula, possibly a hole, is a feature of the individual case.

time the subretinal fluid gravitates to the lower part of the eye and the retina becomes replaced in the upper part. Sarcoma of the choroid may start in the neighbourhood of the macula, and in this case central vision is early affected, as shown by distortion of objects (metamorphopsia, micropsia, &c.) or a relative scotoma for colours. A small detachment due to a sarcoma of the choroid may be accompanied by a large simple detachment in the usual situation below.

Externally the eye looks normal; the anterior chamber may be deep and the tension diminished, though rarely much in the early stages. In cases due to sarcoma of the choroid the tension is always raised in the later stages and the anterior chamber is shallow; occasionally the tension is raised early in these cases.

It is by no means difficult to miss diagnosing a detachment of the retina even when it is large, especially if it is also shallow. The symptoms may be indefinite, for the retina may obtain sufficient nourishment from the fluid which underlies it to retain its functions only partially impaired for a considerable period. Failure in diagnosis is almost always due to the omission of a proper routine examination of the eye. The observer often employs the direct method, possibly after a casual examination by the indirect, without previously examining with the mirror alone. A shallow detachment will then appear little altered from the normal fundus. It is true that it is more hypermetropic, but the observer does not realise that he is exercising rather more accommodation in looking at it than at the rest of the fundus. Now, if the eye is examined with the mirror alone at ordinary reading distance, although perhaps no details of the retina are seen—and they will only be seen if the retina bulges far away from the choroid—yet some difference in the nature of the reflex as the eye is turned in various directions will at once arrest attention.

Further examination by the direct method will then show the following changes (Plate XV.). The detached portion of retina has a different tint from the normal fundus. In the most typical condition it is quite white or grey, with folds which show a bright sheen at the summits and appear greenish grey in the depressions. During slight movements of the eye the folds show oscillations. The retinal vessels are seen coursing over the surface. They naturally follow all the curves of the folds. Very striking is their abnormal colour. Owing to the fact that they are separated from the choroid, which is responsible for the red reflex of the normal fundus,

they are under much the same conditions as an ordinary vitreous opacity, *i.e.*, they cut off the light reflected from the choroid. They therefore look much darker than usual, and may be almost black. They show no central light streak and appear smaller than normal.

If now this portion of the fundus is observed with the highest convex glass with which it can be seen plainly it will be found that the rest of the fundus is out of focus ; this proves that it is displaced forwards and must therefore be detached from the choroid.

In the early stages, and sometimes for a long period in shallow detachments, the colour differs little from the normal red reflex. Such cases are much less easy to diagnose, and may give rise to great difficulty, especially if the apparent detachment is far forwards, for the periphery even of the emmetropic eye is usually seen best with a low convex glass.



FIG. 198. — Diagrammatic sagittal section of eye with partial simple detachment of the retina. (Nettleship.)

When the detachment is very extensive great balloon-like folds may be seen, and these will probably cut off all view of the disc. At the edges of the detachment a considerable degree of pigmentary disturbance may be seen. White spots of exudation, hæmorrhages, &c., may be seen on the surface of the detached retina. Not infrequently a hole is visible, through which the bright red choroid can be seen. It has been said that there is a hole in every detached retina, but it is not always visible ophthalmoscopically.

In total detachment the retina is umbrella-shaped, remaining attached at the disc and at the ora serrata. Still later it becomes bunched up behind the lens, the part attached to the disc being pulled out into a straight cord. In these cases the disturbance to nutrition of the eye leads to the development of a complicated cataract (*q.v.*), so that ophthalmoscopic examination becomes impossible.

The description given applies especially to so-called *simple* detachment, *i.e.*, detachment not due to sarcoma of the choroid. The difference may be slight, but accurate diagnosis is of the utmost importance, since the life of the patient may depend upon it. The chief diagnostic features are given in treating of sarcoma of the choroid, and should be very carefully studied (*vide* p. 385).

The space between the retina and choroid is filled with a highly albuminous fluid secreted by the choroid (Figs. 191, 152, 147).

The prognosis in simple detachment is unfavourable. The only cases in which reposition is likely to occur are those due to albuminuric retinitis (*vide* p. 340). Even if replacement occurs spontaneously, as sometimes happens, or, as the result of treatment, relapses commonly happen and total blindness eventually results. Detachment in a myopic eye should be an indication for extreme care of the other, which may be regarded as predisposed to the same accident. The patient should be warned against stooping, as in gardening, lifting heavy weights, etc.

Treatment.—In spite of the futility of treatment in the majority of cases it should be systematically applied and perseveringly carried out in all cases which come under observation at an early stage, *i.e.*, before the retina has become so degenerated as to have lost all excitability to light. The treatment is very tedious, and is absolutely useless unless minute directions are given and observed.

Of the many modes which have been advocated the following is probably the best. The patient is put to bed and must lie in the supine position without moving more than is absolutely necessary for several weeks. The head should be placed between sandbags. The patient must not be allowed to get up for micturition or defæcation. Atropin is instilled, and the eye is lightly but firmly bandaged, the space between the nose and cheek being filled with cotton wool. A thorough course of pilocarpin injections is given—every day for a week, repeated after a week's interval, and so on. Mercury is administered either by inunction or by the mouth. Iodides are given.

Operative procedures may be adopted. Of these the best are subconjunctival injections of 10 per cent. salt solution once or twice a week. Acoin, 0·1 per cent., may be added to the saline to relieve pain. Not more than 5 minims should be injected at first, as far away from the limbus as possible at the upper part of the globe. The eye should of course be previously cocained.

Probably more efficacious is *scleral puncture.* This is effected as follows. The eye is cocained and washed out (*vide* p. 199). A speculum is inserted, and the patient is told to look as far as possible in the direction away from the detachment, *i.e.*, if the detachment is downwards he looks well up. The eye is fixed

in this position with fixation forceps, held by an assistant. The surgeon takes the conjunctiva far back with another pair of fixation forceps and pulls it well forwards. He then introduces a Graefe cataract knife straight into the globe at a point corresponding with what he considers the site of greatest detachment, usually near the equator. The back of the knife is directed forwards. The knife is pushed in until it perforates the choroid *and* the retina, *i.e.*, for 10 or 12 mm. It is then rotated so that the face of the blade is at right angles to its former position and is withdrawn. In this manner a somewhat valvular wound is made. As the knife is withdrawn the subretinal fluid escapes. The fixation forceps are then removed; the conjunctiva, which has been pulled forwards, retracts and covers the wound. Atropin is instilled and a bandage applied.

Good results have been obtained by using the galvanocautery for puncturing the sclerotic. In this case the conjunctiva must be divided first. It is well to sear a fairly large area of sclerotic before puncturing. The conjunctival wound is then sutured. Posterior sclerectomy with the trephine has also been done.

Scleral puncture is probably useless unless the retina is pierced, thus establishing a communication between the subretinal space and the vitreous, and so aiding the absorption of the subretinal fluid. The retina may become anchored at the site of the wound.

Glioma of the Retina.—See p. 389.

Cysts of the Retina are commonly found in the microscopic examination of degenerated eyes, especially near the ora serrata in old people. Larger cysts occur elsewhere, and are sometimes due to adhesion of folds of detached retina. In rare cases large cysts may simulate detached retina clinically.

Congenital Abnormalities of the Choroid and Retina.

Coloboma of the Choroid and Retina is a congenital malformation in which the choroid and retina are more or less badly developed over a certain area, usually the lower part (typical coloboma). The typical coloboma is due to defective closure of the fœtal, so-called choroidal, cleft. Ophthalmoscopically there is a glistening white area, usually with patches of pigment at the edges (Plate XVI.). There is often also coloboma of the iris (*q.v.*), and the eye may be small

PLATE XVI. (*To face p.* 354)

COLOBOMA OF THE CHOROID.

A. W. Head

(microphthalmia). The condition is often bilateral, and is frequently hereditary. The patch is oval or comet-shaped, with the rounded apex towards the disc, which may be included or not. A few vessels are seen over the surface, some retinal, others derived from the choroid at the edges, but most derived from the posterior ciliaries directly. The surface is often depressed irregularly (ectatic coloboma). The central vision is generally bad, and there is a scotoma in the field corresponding more or less accurately with the coloboma, though this usually contains some retinal elements near the edges.

Similar patches, often symmetrical in the two eyes, occur in other situations (atypical coloboma), notably at the macula



FIG. 199.—Opaque nerve fibres (*O*), stained by the Weigert-Pal method. *R*, retina. *L*, lamina cribrosa. *N*, optic nerve.

(central or macular coloboma). It is probable that some of these are due to intra-uterine inflammation.

Albinism is the defective development of pigment in the eye. There is no pigment in the eye at all, nor indeed in the whole body, in total albinism, so that the iris looks pink, and the patients suffer much from dazzling. Partial albinism is commoner, and the absence of pigment is then limited to the choroid and retina, the irides being blue. With the ophthalmoscope the retinal and choroidal vessels are seen with great clearness, separated by glistening white spaces where the sclerotic shines through. There may be pigment in the macular regions, which may therefore look normal. Nystagmus, photophobia, and defective vision—partly due to myopia, or less frequently, hypermetropia—are generally present. There may be strabismus, usually convergent. The condition is hereditary.

Treatment consists in correction of the refraction by glasses which should be tinted.

Congenital Pigmentation of the Retina.—Small oval grey spots or groups of polygonal greyish-black spots are occasionally met with in the retina in routine examination of the fundus. They are flat and lie below the vessels, and remain unchanged indefinitely. They are probably congenital and due to heaps of retinal pigment epithelium similar to those forming melanomata in the iris (*q.v.*).

Opaque Nerve Fibres.—The medullary sheaths of the fibres of the optic nerve cease normally at the lamina cribrosa. Occasionally patches of fibres regain these sheaths after they have passed through the lamina cribrosa (Fig. 199). They appear ophthalmoscopically as white patches, the peripheral edges of which are radially striated, looking as if frayed out (Plate XVII.). Usually the patches are continuous with the disc: occasionally they are isolated, but rarely far from the disc. Usually the retinal vessels are covered in places by the opaque fibres. When present the blind spot is enlarged, or a scotoma corresponds with the position of the patch. Very rarely the patch is large and involves the macula, so that central vision is abolished. If glaucoma or optic atrophy causes the fibres to degenerate the medullary sheaths disappear and no trace of the abnormality remains. It is important to be able to diagnose them, since they may be easily mistaken for exudates, *e.g.*, albuminuric retinitis. They not infrequently occur in both eyes. They are not strictly speaking congenital, for myelination of the optic nerve progresses from the brain towards the periphery, and is not completed until shortly after birth.

PLATE XVII. (*To face p.* 356.)

OPAQUE NERVE FIBRES.

A.W.Head.

FIG. 1.

FIG. 2.

PLATE XVIII. (*To face p.* 357.)

Fig. 1.—Papillœdema or " choked disc " of intracranial disease. Note the swelling of the disc, as shown by the bending of the vessels, the white exudates, the venous congestion and the hæmorrhages.

Fig. 2.—Myopic changes in the fundus (indirect method). Note the myopic crescent and the visibility of the choroidal vessels, due to the stretching of the posterior pole of the eye and the consequent thinning of the retinal pigment layer. The pigmentation and appearance of splits at the macula and the spots of choroidal atrophy are such as occur in severe cases of high myopia.

CHAPTER XVIII.

DISEASES OF THE OPTIC NERVE.

The optic nerve may be attacked by inflammation at any part of its course. The head of the nerve within the globe is frequently affected alone, and this condition is often called optic neuritis. When the nerve is affected behind the eye the condition is called retro-bulbar or retro-ocular neuritis.

Hyperæmia of the Optic Disc is a condition which can rarely be diagnosed with certainty. Perfectly normal discs of different individuals show variations in colouring, and slight differences of illumination alter the appearances. There is no doubt that hyperæmia occurs as a precursor of optic neuritis, and in some of these cases it is possible to distinguish greater redness of the disc in one eye than in its fellow under the same conditions of examination.

Papillitis (*Intraocular Optic Neuritis*) **and Papillœdema.**—Papillitis occurs in two groups of cases which should be carefully distinguished, viz., as part of a neuro-retinitis (*vide* pp. 329, 340), and as a result of intracranial disease. The pathology of the two conditions is totally different, though the ophthalmoscopic features may be almost or quite identical. These general features will be considered first, the differences being discussed afterwards.

It has already been pointed out that the colour of the disc is a fallacious criterion of abnormality. Attention should be directed especially to the edges, which will always be found blurred in papillitis, usually first on the nasal side, later in the whole circumference. The blood vessels are altered, the arteries being small, the veins distended. In the early stages the disc is usually redder than normal. In the later stages the blurring of the edges is much greater, and the disc looks larger than usual. Exudates cover the vessels in places, and the veins are enormously distended and very tortuous. The papilla is paler than normal and may be white; it shows radial streaks, and small hæmorrhages are generally present upon it and the surrounding retina. The disc is now quite definitely and measurably swollen (*vide* p. 116).

In the optic neuritis of neuro-retinitis the swelling of the disc is usually moderate—2 or 3 D—shelving off gradually into the surrounding retina, which shows the signs of retinitis (*vide* p. 328). The disc is redder than normal, owing to dilatation of the capillaries, but the distension and tortuosity of the veins is moderate. The "optic neuritis" of intracranial disease may in some cases be a true descending neuritis, and will then show the condition just described, except that there is little or no retinitis. The intracranial disease in these cases is usually of an inflammatory nature, *e.g.*, meningitis. More commonly, however, the condition is one of intense œdema with no true inflammation, and this gives rise to papillœdema or "choked disc" (Plate XVIII., Fig. 1). Here there is very great swelling—up to 8 or 10 D—usually delimited much more definitely from the surrounding retina, which shows little change. The veins are enormously distended and very tortuous; the vessels are hidden in places by white exudates. There are frequently hæmorrhages on the swollen papilla and at its edges. In many cases it is impossible to distinguish ophthalmoscopically between the two forms. The swelling may be unusually great in neuro-retinitis, and per contra the appearances due to intracranial disease may simulate the neuro-retinitis of renal disease, including the star figure at the macula, particularly in children. The star figure is rarely complete in these cases; usually it is a fan-shaped figure on the disc side of the macula.

The symptoms of papillitis may be extremely vague. Central vision may be quite normal, in which case the pupils will also be normal in size and reactions. It is very important to remember this fact, which emphasises the necessity of examining the fundus in all cases of headache, &c. Even in this stage there may be some concentric contraction of the field of vision. Transient attacks of blurred vision, lasting for a few minutes up to an hour or so, are not uncommon in the early stages of papillœdema. Later central vision is reduced, even to complete blindness. The pupils will then be large and immobile. In less severe cases or in the intermediate stage central vision is defective, the field is markedly reduced concentrically, and relative scotomata—first to green and red—or absolute scotomata may be present. There may be hemianopia or other defects in the field dependent upon the nature of intracranial mischief.

Both in neuro-retinitis and in intracranial disease the condition is generally bilateral, though not necessarily equal on

the two sides. The relative amount of swelling may be of localising value in the case of intracranial disease, but its value has certainly been over-estimated ; in frontal tumours and middle ear disease the swelling is usually greater on the side of the lesion. The time of onset of the papillœdema is really more to be taken into account than the amount of swelling, the localising value being attached to the side first affected. Thus the swelling may be actually less on the side first affected owing to subsidence associated with commencing atrophy. Unilateral papillœdema, with or without " secondary " optic atrophy on the other side, suggests a tumour of the opposite olfactory lobe or orbital surface of the frontal lobe or of the pituitary body (*vide* p. 362). Unilateral papillitis occurs in the early stages of papillœdema from intracranial causes and in orbital diseases such as tumours of the optic nerve or orbit, cellulitis of the orbit, hæmorrhage into the sheath of the optic nerve, &c.

The course is chronic, the prognosis bad. Occasionally, especially in syphilis, the disease subsides under treatment and good vision is preserved. The same applies to cases due to intracranial pressure if the pressure is relieved early. Palliative decompression by trephining the skull has a remarkable effect. Headache, vomiting and stupor are relieved, vision improves rapidly unless the nerves have been irretrievably damaged, and the papillœdema quickly subsides. Rarely the discs regain a normal appearance, but generally the nerve fibres are destroyed, post-neuritic atrophy (*q.v.*) follows, and the patient becomes blind. Recurrence of papillœdema is very rare, but has been recorded.

Diagnosis is easy in severe cases ; it may be very difficult in slight cases. Here the colour of the disc is no sure guide unless there is undoubted difference between the two eyes. Attention should be directed to the edges of the disc ; if these can be seen clearly defined with any lens there is no neuritis, but it does not follow that there is neuritis if they appear blurred. Astigmatism causes apparent blurring of the disc margin and there is a condition nearly simulating slight neuritis known as pseudo-neuritis, seen particularly in hypermetropic eyes and due largely to a peculiar reflex (Fig. 200). In such cases attention must next be directed to the amount of swelling. In the absence of other indubitable signs such as exudates or hæmorrhages papillitis should not be diagnosed unless at least 2 D of swelling can be demonstrated. In some cases it is necessary to keep the

patient under careful observation for a considerable period before certainty can be arrived at.

The following stages may be distinguished in the development of papillœdema (Marcus Gunn, de Schweinitz and Holloway) :—

(1) Increased redness of the disc, with blurring of its upper and lower margins, with a gradual progression of the blurring

Fig. 200.—Pseudo-neuritis.

to the nasal edges, while the temporal margin is still visible, represents the first stage.

(2) Increased œdema of the nerve-head, beginning filling in of the physiological pit, involvement of the temporal margin of the disc, with a tendency of the œdema to spread into the surrounding retinal area, and uneven distension and darkening of the retinal veins, represent the second stage.

(3) Decided increase of œdema, elevation and size of the nerve-head, with vascular striation of the swollen tissue and striæ of œdema in the form of lines in the swollen retina between the disc

and macula, marked distension of the retinal veins and retinal hæmorrhages, represent the third stage.

(4) Increase in the prominence of the disc, which assumes a mound-shape and begins to lose its reddish and juicy colour and to become opaque, exudation in and on the swollen disc and surrounding retina, elaboration of the retinal hæmorrhages in size and number, represent the fourth stage.

(5) Decided subsidence of the vascularity of the papillœdema and increasing pallor, with or without sinking of its prominence, apparent contraction of the retinal arteries and thickening of their perivascular lymph-sheaths, spots of degeneration of the retina, especially in the macula, represent the fifth stage, which passes into the final stage of so-called papillitic atrophy.

Ætiology.—The chief causes of papillitis are those of neuro-retinitis (*q.v.*) and intracranial disease. The latter is the most frequent cause, especially intracranial tumour, at least 80 per cent. of these cases giving rise to the condition. Any intracranial tumour in any position, with the exception of the medulla oblongata, may cause papillœdema, the highest percentage being found with tumours of the mid-brain, parieto-occipital region, and cerebellum (*vide* p. 552). Meningitis is the next commonest cause, especially tuberculous meningitis—basal meningitis relatively rarely. Other intracranial causes are abscess, thrombosis of the cavernous sinus, aneurysm, hydrocephalus (rarely), &c.

Toxæmia accounts for most of the other cases. Syphilis may act in this manner as a basal meningitis, or more frequently as an intracranial gumma ; it is a frequent cause. Papillitis may be due to any of the acute febrile diseases, but only in exceptional cases, and to acute anæmia from sudden loss of blood. It occurs sometimes with poisoning with lead and other substances which usually give rise to toxic amblyopia of retrobulbar type (*vide* p. 348). It has often been described in chlorosis, and suppression of menses has been given as a cause. This ætiology is in my opinion doubtful ; some cases have subsequently proved to be due to other more serious causes, and many are probably errors of diagnosis (*see* Pseudo-neuritis, p. 359). Slight papillitis has been observed in some of those rare cases in which there is persistent escape of cerebrospinal fluid from the nose. Malformation of the cranium, *e.g.*, oxycephaly, intra-orbital tumours and inflammations—caries, periostitis, &c.—and tumours of the optic nerve act directly upon the nerve.

Pathology.—In neuro-retinitis there is true inflammation of

the nerve, but even then œdema plays a prominent part owing to the obstruction to the outflow of venous blood at the site of the lamina cribrosa (*vide* p. 120). Most cases of "choked disc" are at first caused by simple œdema, without inflammation; in the later stages there may be some inflammatory reaction—infiltration with leucocytes, &c.—due to the irritation of the necrosed tissues. In a minority of cases due to intracranial disease, especially meningitis, there may be a descending neuritis, with true inflammation of the nerve.

The mechanism whereby œdema is induced by intracranial disease has been the subject of much dispute and is still an unsolved problem. There is no doubt that the predominant factor is increased intracranial pressure. This is proved by the fact that the œdema almost invariably subsides, even though the intracranial disease continues, if the pressure is relieved by freely opening the skull. If communication is cut off from one intravaginal space by pressure of a tumour of the pituitary body or olfactory region papillœdema does not occur on this side, but the optic nerve passes into a condition of partial or complete "secondary" atrophy (*vide* p. 367). The mode of action of increased intracranial pressure is unknown. The following theories have been advanced: (1) impediment to the return of lymph along the intravaginal space of the nerve; (2) impediment to the return of venous blood—this is improbable owing to the free anastomosis in the orbit (*vide* p. 10); there is, however, excessive pressure on the central vein of the retina where it traverses the intravaginal space behind the globe; (3) arterial anæmia, leading to altered conditions of osmosis at the nerve-head.

The *treatment* of optic neuritis is essentially that of the underlying cause. Local treatment is of no avail, but all sources of irritation, such as bright light, &c., should be guarded against. Intracranial pressure should be relieved before vision is lost, even if the cause is a tumour which cannot be completely extirpated.

If the localising symptoms are positive trephining should be performed over the site of the tumour if possible. If this is impossible or if localising signs are masked by pressure symptoms a free opening should be made in the skull supra- or subtentorially according to the indications of the case. The relief of pressure will alleviate the cerebral œdema and unmask the localising signs, so that it may be possible at a later stage to undertake a radical operation. Paralysis of the external recti is often due to the pressure on the sixth nerves in their long

intracranial course, and in these cases is of no localising value.

If decompression is done early the prognosis as to sight is very favourable. If delayed until there is great swelling and exudation, with marked depreciation of vision, and especially if signs of subsidence and commencing atrophy are present, further diminution of vision is to be anticipated. Subsidence of the papillœdema is usually rapid after operation, a decided change being seen in a week to a fortnight, though there is considerable variation in different cases. In cerebral abscess there may be temporary increase of the papillitis or swelling after operation, without however seriously compromising the prognosis.

Retrobulbar Neuritis.—The intra-orbital portion of the optic nerve may become inflamed as the result of extension from the surrounding tissues. Retrobulbar neuritis is usually divided into an acute, so-called symptomatic form, and a chronic, idiopathic form. The latter is the condition which we have described as toxic amblyopia and have attributed to a primary retinal lesion.



FIG. 201.—Diagrammatic meridional section of optic disc in papillitis.

Acute retrobulbar neuritis is usually unilateral. The patient complains of rapidly increasing defect of vision, and sometimes of pain on moving the eye. The pain is increased by pressure upon the globe, and neuralgia and headache may be present. Ophthalmoscopic examination will probably reveal a quite normal fundus. It is therefore very easy to overlook the true condition and to attribute the symptoms to hysteria. Careful methodical examination will minimise this danger. Diagnosis at this stage will depend upon thorough investigation of the pupil reactions and of the field of vision. The patient should be asked if he ever squinted (*vide* p. 376).

At first glance the pupil reactions will be apparently normal, both directly and consensually to light, as well as to accommodation. More minute inspection will show, however, that though the pupil of the affected eye reacts to light the contraction is not maintained under the bright illumination, *i.e.*, instead of remaining contracted the pupil slowly dilates while the light is still kept upon the eye. This phenomenon, if it can

be placed beyond dispute, is of the greatest diagnostic significance.

The field of vision shows a central scotoma, which may be relative for colours or absolute. It is not always quite central, but may be paracentral or sectorial or in the form of a ring around the fixation point. There is usually some peripheral loss of the field and there may be complete blindness.

In the later stages or more severe cases there are usually ophthalmoscopic changes. These are distension of the veins, with diminished calibre of the arteries, or actual papillitis, moderate in degree. They are most likely to occur if the focus is close behind the eyeball. With or without these preliminary changes atrophy of the optic disc may ensue. In every case in which the inflammation in the nerve behind the globe is so great as to lead to destruction of the nerve fibres in this situation the degeneration extends not only towards the brain but also towards the eye. In the milder cases pallor of the disc is limited to the temporal side, corresponding with degeneration of the papillo-macular fibres. It has already been noted that the ganglion cells at the macula are more liable to be affected by deleterious agents than those in other parts of the retina. The same fact applies to their axis cylinder processes, contained in the papillo-macular bundle, wherever in their course the noxious agent acts. This accounts for the clinical similarity between true retrobulbar neuritis and toxic amblyopia.

The causes of acute retrobulbar neuritis are local and general. Among the former periostitis and transmission of inflammation from the ethmoidal and sphenoidal sinuses may be mentioned ; hence attention should be directed to any nasal complication. Hæmorrhage into the optic nerve sheath or orbit and fracture of the base of the skull, involving the optic foramen, may produce a similar clinical picture. Pressure on the chiasma by hypertrophy or tumour of the pituitary body may in the early stages cause the symptoms of a unilateral retrobulbar neuritis (*vide* p. 374). Among general diseases which cause retrobulbar neuritis insular sclerosis is one of the most important ; it is a very frequent early symptom in this disease. Other alleged causes are rheumatism, chills, diabetes, infectious diseases, septic foci somewhere in the body (mouth, intestinal tract, &c.), and so on. In cases due to such causes relapses are common, and both eyes may be affected, together or alternately. Acute retrobulbar neuritis may precede peripheral facial palsy of the same or opposite side, and shows some analogies to this condition (Marcus Gunn). The prognosis depends upon the cause

and the possibility of combating it, but is generally good. In multiple sclerosis the affection of the nerve rarely leads to complete blindness.

Treatment consists in attacking the cause, the mouth and nasal sinuses being specially carefully investigated. The eyes must be protected from bright light, and kept at rest by atropin and abandonment of near work. When the cause is obscure, mercury, iodides, salicylates, diaphoresis, and tonics may be used. Smoking should be prohibited. In diabetic cases appropriate régime should be instituted.

Hereditary Optic Neuritis (*Syns.—Hereditary Optic Atrophy, Leber's Disease*) is a form of retrobulbar neuritis, usually commencing at about the twentieth year of life. Descent is generally through an unaffected female to the males, though females are also sometimes affected. Vision generally fails rapidly at first, then gradually, then remains stationary or gradually improves. Both eyes are always affected, though one may precede the other by a few days up to eighteen months. In two-thirds of the cases there is a central scotoma, either partial for colours or also for white. The peripheral field is usually normal, but concentric contraction or sector-shaped defects may occur. Total and permanent colour-blindness has been known to follow. The central scotoma generally persists, but progressive constriction of the field to complete blindness is rare. Members of the same family often show identical peculiarities in the progress of the cases. The fundus is at first normal or there is slight blurring of the edges of the disc. In later stages, after months, optic atrophy ensues, with pallor confined to the temporal side or involving the whole disc. Apart from headache, migraine, &c., the general health is good. Herbert Fisher has suggested that Leber's disease is due to transitory changes in the pituitary body (*vide* p. 379), resulting in pressure upon the chiasma and associated with the periods of physiological change in the sexual life. If this be true treatment with thyroid and pituitary extracts may do good.

Optic Atrophy is the term usually applied to the condition of the disc when the optic nerve is degenerated. It has been pointed out that injury to the nerve fibres in any part of their course from the retina to the external geniculate body leads to degeneration not only on the proximal (cerebral) side—as might be anticipated for afferent fibres—but also on the distal (ocular) side (*vide* p. 364). Optic atrophy follows extensive disease of the retina from destruction of the ganglion cells, *e.g.*, in "retinitis" pigmentosa; it also follows destruction of the nerve in the orbit, as in rupture of the nerve at the optic

foramen in fracture of the base of the skull, severe retrobulbar neuritis, &c. The break in continuity of the fibres may be at the disc itself, such as results from the strangulation of the papillitis of neuro-retinitis or papillœdema. These cases are distinguished as *post-neuritic atrophy*, owing to special features which they often exhibit. Besides these causes there is a well-defined group of cases in which optic atrophy occurs without previous evidence of severe local inflammation, but associated with general disease, usually of the central nervous system, or without discoverable disease. Such cases are described as *primary atrophy*.

The essential ophthalmoscopic features of optic atrophy in general are alteration in the colour of the disc and changes in the blood vessels. The disc is always pale, but may show varieties of tint specially associated with various types of atrophy. The pallor affects the whole disc and must be carefully distinguished from the white centre, often encroaching upon the temporal side, due to physiological cupping. The change in the vessels is always on the side of contraction, but may be absent.

Fig. 202.—Diagrammatic meridional section of the optic disc in atrophic cupping. Note that the lamina cribrosa is not displaced (*cf.* Fig. 157).

In primary (grey, tabetic, spinal) atrophy the disc is grey or white, sometimes with a greenish or bluish tint (Plate XIX., Fig. 1). The stippling of the lamina cribrosa is seen; the edges are sharply defined, and the surrounding retina looks normal. Owing to the degeneration of the nerve fibres there is slight cupping (atrophic cupping) (Fig. 202), which must be carefully distinguished from glaucomatous cupping. It is shallow and saucer-shaped, as shown by the slight bending of the vessels, but is scarcely measurable with the ophthalmoscope. There is no retraction of the lamina cribrosa. The vessels are normal or only slightly contracted. Both eyes are generally affected.

In the "secondary" atrophy of retrobulbar mischief the condition nearly resembles primary atrophy, but there is greater pallor, and the vessels are more likely to be contracted. In the "consecutive" atrophy of retinal and choroidal disease, as typically represented by the late stages of "retinitis" pigmentosa, the disc has a yellowish waxy appearance, the edges are less sharply defined, and the vessels are very markedly contracted, sometimes to almost complete disappearance.

PLATE XIX. (*To face p.* 366)

OPTIC ATROPHY.

Fig. 1.—Primary optic atrophy. Note the pallor and sharply cut margin of the disc, the exposure of the lamina cribrosa, and the absence of circumpapillary retinal changes.

Fig. 2.—Postpapillitic atrophy. Note the pallor and blurring of the edges of the disc, the filling in of the atrophic cup with fibrous tissue so that the lamina cribrosa is not seen, the disturbance of the retina in the immediate neighbourhood of the disc, and the thread-like arteries.



FIG. 1.

FIG. 2.

Post-neuritic atrophy may be indistinguishable from the other forms ophthalmoscopically. More commonly it can be diagnosed with considerable probability by characteristic signs (Plate XIX., Fig. 2). These depend upon the fact that the absorption of the exudates leads to a certain amount of reactionary organisation, with the formation of a variable quantity of fibrous tissue upon the disc. This tissue obscures the lamina cribrosa and fills in the atrophic cup. It extends over the edges, which are thus indefinite, and along the vessels as a thickening of the perivascular sheaths. Further, it throttles the vessels, so that they become markedly contracted, especially the arteries. Owing to previous neuro-retinitis the surrounding retina often shows permanent changes, chiefly manifested by pigmentary disturbance. When such changes are well marked previous papillitis may be hypothecated with great probability, but in their absence the conclusion that there has not been papillitis is not justifiable. The amount of reactionary organisation varies greatly in different cases, and the tissue laid down is in the course of time gradually absorbed to some extent. It must also be remembered that the amount of fibrous tissue on the normal disc varies considerably and that slight filling in of the physiological cup with shreds of fibrous tissue extending outwards along the vessels occurs as a congenital peculiarity.

In total optic atrophy the pupils are dilated and immobile to light and the patient is blind. When unilateral the consensual reaction to light is exaggerated. In partial optic atrophy central vision is depressed and there is concentric contraction of the field, with or without scotomata, relative or absolute, according to the cause. In primary atrophy the disease is usually slowly progressive, ending after months or years in complete blindness. It is important to note that no deduction as to the amount of vision can be made from the ophthalmoscopic appearances. The presence of all the signs of atrophy is not inconsistent with a certain, sometimes considerable, amount of vision.

The chief cause of primary optic atrophy is tabes. It may be the first sign, and the other symptoms and signs may be long delayed. The patient should be carefully investigated for a history of syphilis, the presence of Argyll Robertson pupils, the absence of knee-jerks, the presence of anæsthesia (especially of the fifth nerve), inco-ordination, a positive Wassermann reaction, and so on. If optic atrophy occurs early in tabes, ataxy may be long delayed or never supervene. On

the other hand, if it occurs late there is no appreciable difference in the course of the ataxy. Both eyes are affected, but often in unequal degree. The disease advances slowly but surely to complete blindness—progressive optic atrophy par excellence. The lesion was formerly supposed to be in the ganglion cells of the retina, but there is now considerable evidence that the earliest change is an inflammatory exudation into the intracranial part of the nerve and the chiasma. A similar condition may be due to general paralysis and insular sclerosis, less commonly to other forms of central nervous disease: in insular sclerosis the course of the disease is usually different —variations or repeated acute attacks, seldom causing blindness. The more favourable prognosis in disseminated sclerosis may be due to the fact that in this disease the axis cylinders often escape in spite of much destruction of their medullary sheaths. Primary atrophy may be due to poisons, such as methyl alcohol, lead, atoxyl, &c. (*vide* p. 349). Excessive hæmorrhage, menstrual defects, cold, arterio-sclerosis, &c., have been ascribed as causes in doubtful cases.

In primary atrophy, though central vision is early affected, there is usually no central scotoma. The field shows progressive concentric contraction, often with marked indentations which are rather more common on the nasal side. Finally, usually in a year or two in tabes, the fixation point is engulfed, though eccentric perception of light may still persist for a while on the temporal side. Contraction of the colour fields precedes that of the field for white, so that there is a stage of acquired colour-blindness. The prognosis is very bad.

"Secondary" atrophy is caused by compression of the optic nerves, chiasma, or tracts by tumours, aneurysm, distension of the lateral ventricles (hydrocephalus), &c. These tumours, &c., are so situated as to press directly upon the nerve fibres without causing much rise of intracranial pressure, as in the case of tumours of the pituitary body, aneurysm of the internal carotid in the cavernous sinus, &c., or to press directly on the chiasma or optic nerves behind the optic foramen, thus blocking communication with the intravaginal lymph-space, as in the case of tumours of the olfactory lobe and inferior aspect of the frontal lobe. The same type of atrophy may follow rupture of the optic nerve at the foramen, hæmorrhage into the dural sheath, section of the nerve, compression by blood clot, oxycephaly, &c., without previous neuritis.

The prognosis and treatment of post-neuritic atrophy have been dealt with in discussing papillœdema.

Treatment of primary atrophy is that of the cause. For the lesion of the nerve itself mercury, iodides, strychnin, nitroglycerin, the constant current, &c., may be used, but seldom have any effect.

Tumours of the Optic Nerve.—See Chap. XXXIII.

Congenital Abnormalities of the Optic Disc.

Coloboma of the Optic Disc.—This occurs in two forms, one of which is common, the other rare. The common form is due to incomplete closure of the choroidal fissure, and manifests itself as an *inferior crescent*, much resembling the myopic crescent (*q.v.*), but situated at the lower edge of the disc (Plate VIII., Fig. 1). It is a crescent, whiter than the disc itself, situated at the lower border. It occurs most commonly in hypermetropic and astigmatic eyes, which are often found to have slightly defective vision though the error of refraction is corrected.

In what is commonly known as *coloboma of the disc* (or nerve sheath) there is greater failure of the fœtal fissure to close. The disc then looks very large and the vessels have a very abnormal distribution, appearing only above or irregularly round the edges. The apparent disc is really the sclerotic and inner surface of the sheath of the nerve, the nerve itself being usually spread out as a pink horizontal linear band at the upper part. The floor of the coloboma is white and measurably depressed, often quite ectatic. The eye usually has defective vision.

Rarer anomalies allied to coloboma are round "*holes*" in the disc, usually looking grey or black owing to the shadow, and situated in the temporal portion of the disc; and patches of *pigment* due to inclusion of retinal pigment epithelium.

Mention has already been made (p. 367) of excess of *fibrous tissue on the disc* and extending a short distance along the vessels. Sometimes the fibrous tissue takes the form of a delicate semi-transparent membrane covering the disc and appearing to be slung from the vessels.

CHAPTER XIX.

SYMPTOMATIC DISTURBANCES OF VISION.

Apart from the disturbances of vision which have been already considered and have their origin in the eye itself, there are others dependent upon lesions in the visual nervous tracts. They not infrequently closely simulate the disorders due to peripheral causes, or, being early evidence of disease, lead the patient to consult an ophthalmic surgeon. There are also visual defects the cause and seat of which are imperfectly elucidated; though some are probably peripheral in origin it will be convenient to consider them here.

Hemianopia (*Syns.—Hemianopsia, Hemiopia*).—Hemianopia denotes loss of half of the field of vision. The commonest clinical form is so-called *homonymous hemianopia*, in which the right or left half of the binocular field of vision is lost, owing to loss of the temporal half of one field and the nasal half of the other. The condition may be due to a lesion situated in any part of the visual tracts from the occipital lobe to the chiasma. A focus of disease in this area causes loss of vision of the corresponding halves of each retina (hence the designation homonymous), *i.e.*, loss of the opposite halves of the visual fields.

In most cases of hemianopia the fixation point of each eye escapes (Fig. 203). This is probably due to two causes—widespread representation of the foveal region in the occipital lobe of the same side, owing to spread of the fibres of the papillo-macular bundle and of the corresponding neurons of the third order; and bilateral representation of each foveal region in each occipital lobe. On Gordon Holmes and Lister's view of unilateral representation of the macula (*vide* p. 69) the escape of the fixation point in vascular lesions is attributed to the fact that the occipital pole is supplied by branches of the posterior and middle cerebral arteries, and that both of these arteries are seldom blocked at the same time.

Lesions of the external geniculate body cause homonymous hemianopia; those limited to the pulvinar and superior colliculus do not. Right hemianopia is much more quickly dis-

covered than left, owing to the fact that reading is impossible. Left hemianopia is often discovered by the fact that the patient does not see food on the left side of the plate.



Fig. 203.—Charts of fields of vision in homonymous hemianopia.

Cortical and Sub-cortical Lesions.—The majority of cases of hemianopia are due to lesions above the primary visual centres, usually in the occipital lobe or optic radiations. The injury or disease rarely affects the grey matter of the occipital cortex

only; the sub-cortical white matter is almost invariably involved. The chief causes are injury by falls on the back of the head or gun-shot wounds, cerebral tumour, cerebral softening due to syphilitic or other disease of the blood vessels, &c. In gun-shot wounds both occipital lobes are not infrequently injured. There is usually unconsciousness from concussion at first; with the gradual recovery the hemianopia becomes manifest. If both lobes are extensively injured there is bilateral hemianopia with complete blindness. Often, however, some portion of the cortex of one or other calcarine fissure escapes, and in these cases some measure of central vision is regained. In less extensive injury the hemianopic symptoms may gradually improve. The first sign of improvement is the perception of the movement of objects in the affected field, the nature and details of the objects being as yet quite unrecognised. The onset of hemianopia due to disease of the cortex is more gradual, and careful investigation with the perimeter shows that the colour fields are often lost before the field for white light, but the field for white is always contracted in these cases (Gordon Holmes). This *hemiachromatopsia* is itself of gradual onset, the colour fields becoming contracted. In cortical and sub-cortical lesions the pupil reactions are normal (*vide* p. 56), and in many cases the fundi reveal no ophthalmoscopic changes. The chief exception to the latter statement is in the case of tumours of the occipital lobe, in which case the rise of intracranial pressure leads to bilateral papillœdema. Cortical lesions are liable to be accompanied by word-blindness, probably due to involvement of the angular gyrus. When the lesion is in the posterior part of the internal capsule hemianæsthesia, with or without hemiplegia, is likely to be present.

Rare cases of homonymous *quadrant hemianopia* have been reported, in which corresponding quadrants of each field the upper or lower half of one temporal, and the upper or lower half of the other nasal have been lost. These are generally caused by cortical or sub-cortical partial lesions of one occipital lobe, destruction of the part above the calcarine fissure leading to loss of the lower quadrants and *vice versâ*. Figs. 61, 62 show the probable representation of different portions of the field in the visual cortex according to Gordon Holmes and Lister.

Lesions of the Optic Tract.—In this case, since the afferent pupillary fibres part company with the visual fibres before the latter enter the so-called primary optic centres—external geniculate body, pulvinar, and superior colliculus (*vide* p. 56)—Wernicke's hemianopic pupil reaction should be present



Showing the eight stages of a progressing right temporal field defect in pituitary disease. (Harvey Cushing and Clifford B. Walker.)

To face p. 373

(*vide* pp. 56, 88). It must be remembered, however, that this reaction is always difficult to elicit, and with the methods usually employed is seldom conclusive unless definitely present. More assistance in diagnosis is afforded by collateral symptoms. The proximity of the crus cerebri, third and other cranial nerves, leads to not infrequent involvement in the pathological process. The association of hemianopia with contra-lateral third nerve paralysis and ipso-lateral hemiplegia suggests a tract lesion. It is said that the fixation point does not escape in tract hemianopia. Partial atrophy of both optic nerves manifests itself by pallor of the discs in these cases, preceded in cases of raised intracranial pressure by papillœdema. The lesion is usually syphilitic meningitis or gumma, tubercle, or tumour of the optic thalamus or temporo-sphenoidal lobe; softening and hæmorrhage are rare.

Bitemporal hemianopia is usually caused by disease of the pituitary body, which then presses upon the chiasma, so that the fibres going to the nasal halves of each retina are destroyed (Plate XX).

Disease of the pituitary body may manifest itself in forms which are attributed to (1) hyperpituitarism, (2) hypopituitarism, and (3) dyspituitarism. The organ consists of a glandular anterior lobe, and a posterior lobe, composed of nervous tissue, covered anteriorly by a glandular veneer, the pars intermedia. The anterior lobe appears to be specially concerned with skeletal growth, the posterior, including the pars intermedia, with tissue metabolism: *i.e.*, over-activity of the anterior lobe causes excessive growth in the bones; over-activity of the posterior lobe leads to emaciation and glycosuria with polyuria; diminished activity of the posterior lobe leads to adiposity, sugar tolerance, sexual infantilism, low pulse, low temperature, &c. (Cushing). Hyperpituitarism in infancy causes gigantism, in later life acromegaly. Hypopituitarism causes adiposity and persistence of skeletal and sexual infantilism when originating in childhood, adiposity and reversion to sexual infantilism, with development of feminine characteristics in the male, when originating in the adult (Fröhlich's syndrome). Hyperpituitarism often gradually gives place to hypopituitarism; mixed or transition cases exhibit features of both states (dyspituitarism). The adiposity associated with defective action of the posterior lobe is accompanied by excessive sugar tolerance, *i.e.*, 300 or 400 gms. of glucose or lævulose can be assimilated without the development of glycosuria, whereas about 100 gms. is the normal amount. Changes in the pituitary body are accompanied by changes in other ductless glands, such as the thyroid, pancreas, testicles and ovaries.

Enlargement of the pituitary body, whether from functional hyperplasia, adenoma, or malignant growth, generally leads to pressure upon the chiasma, which lies immediately above it (Figs. 204, 205), and upon the inner sides of the optic tracts. The earliest visual symptom is unilateral central scotoma, simulating retrobulbar neuritis (Nettleship), for one side is usually compressed before the other. This may be followed by



FIG. 204.—Antero-posterior section of sella Turcica. *Pit. B.*, pituitary body. *Sph.*, sphenoid. *Chi.*, chiasma.



FIG. 205.—Transverse section of sella Turcica. *Pit. B.*, pituitary body. *Sph.*, sphenoid. *Cav. S.*, cavernous sinus. *Car. A.*, internal carotid artery. *III, IV, VI*, third, fourth and sixth nerves. V_1, V_2, V_3, first, second and third divisions of the fifth nerve.

homonymous hemianopia from pressure on one tract, or rarely by *altitudinal hemianopia*, *i.e.*, loss of the upper or more rarely lower halves of the fields, from pressure upon the chiasma. More commonly temporal hemiachromatopsia, passing into a complete hemianopia, supervenes. The field does not show the accurate delimitation characteristic of homonymous hemianopia, but gradually contracts from the temporal side inwards and from above downwards, finally involving the

nasal field from below upwards and leading to complete blindness in the eye affected. Then, or at a much earlier stage, the vision of the other eye becomes affected in a similar manner. If the second eye becomes affected before vision is lost in the first the fields show bitemporal hemianopia, but one eye is almost invariably more affected than the other, owing to the asymmetry of the growth. Complete temporal hemianopia in one eye, for example, may be associated with temporal achromatopsia in the other. Such cases emphasise the importance of charting the colour fields in all cases. A considerable proportion of cases show homonymous hemianopia, due to pressure and traction on one optic tract. Variations in the progress of the visual defect are not uncommon. In some cases headache disappears, other symptoms are ameliorated, and the vision ceases to deteriorate. Some of these cases are probably to be attributed to cysts which have ruptured spontaneously.

If one eye is blind the history may reveal the fact that the field was lost earliest on the outer side. In cases of acromegaly, the enlargement of the jaw and characteristic facies, the large hands, the loss of sexual desire, and the presence of impotence or amenorrhœa, render the diagnosis easy. Tumours of the hypophysis are less readily diagnosed, but here also loss of sexual power is usually an early symptom, often accompanied by excessive subcutaneous fat. In all cases a skiagram of the skull should be taken; the sella Turcica will be found often, though by no means invariably, enlarged. Associated signs are slight proptosis and paralysis of ocular nerves, generally the third nerve. This is of diagnostic importance, since the sixth nerve is more commonly affected in intracranial lesions. As already mentioned Fisher regards Leber's disease (*vide* p. 365) as probably due to pituitarism.

Some cases improve when treated with thyroid extract, others with pituitary extract, but spontaneous variations in the amount of visual defect are common. Usually the cases come under observation at a late stage when over-activity has given place to insufficiency. Consequently, treatment with active extracts of the gland or of the anterior lobe is indicated. If, as is usual, the progress of the visual defect remains unmitigated the question of operation arises, since otherwise total blindness from optic atrophy is inevitable. The danger to life, whether by the sub-temporal or the intra-nasal route, is considerable.

Bitemporal hemianopia is occasionally due to basal syphilitic inflammation, or disease of the sphenoid. It has also resulted

from antero-posterior rupture of the chiasma in fracture of the base of the skull.

Binasal hemianopia is very rare if, indeed, it can be said to occur at all in typical form. It necessitates two lesions, one on each side of the chiasma, destroying the fibres to the temporal halves of each retina while leaving the nasal fibres intact. In cases which have been reported there has usually been increased intracranial pressure with choked discs, and the condition has been attributed to distention of the third ventricle, causing the optic nerves to be pressed downwards and outwards against the internal carotids. Other cases have been referred to atheroma of the carotids or posterior communicating arteries.

Cases have been described in which there has been loss of half of one field and general blurring or amblyopia of the whole of the other. In some of these cases a lesion has been found post-mortem involving one occipital lobe and extending into the angular gyrus. *Heteronymous amblyopia* has therefore been ascribed to lesion of the angular gyrus. It is probable that this is merely a survival of the erroneous idea that the cortical representation of vision is in the angular gyrus. It is probable that this gyrus has to do with visual conceptions, *i.e.*, mental visual impressions of a higher order than the perceptions represented in the occipital cortex.

Amblyopia (ἀμβλύς, blunt) and **Amaurosis** (ἀμαυρός, dark) are the terms used for partial and complete loss of sight respectively in one or both eyes. They are not used of all cases of partial or complete blindness, but have become restricted to certain forms of a more or less indefinite character devoid of ophthalmoscopic or other marked objective signs.

Unilateral amblyopia is usually either *congenital* (*vide* p. 521) or from psychical suppression of the retinal image—*amblyopia ex anopsia* (*vide* p. 521) : these varieties are discussed elsewhere. Unilateral amblyopia may be due to high refractive errors in the eye. It is then not a true amblyopia, since correction with suitable glasses in early life cures the condition if sufficient perseverance is exercised. In older people glasses often fail : this may be attributed either to the development of a true amblyopia from disuse or, more probably, to defective receptivity of the higher centres. Unilateral amblyopia is also a symptom of retrobulbar neuritis (*q.v.*).

Bilateral amblyopia is found in the various forms of toxic amblyopia (*q.v.*). Bilateral amaurosis occurs in uræmia and in meningitis. Both amblyopia and amaurosis occur in hysteria.

Uræmic Amaurosis occurs particularly in acute nephritis, *e.g.*, in pregnancy, after scarlet fever, &c., but is also found with chronic nephritis. The onset of blindness is sudden or rapid (8–24 hours); it is bilateral and complete. The fundi show no changes, unless, as in some cases, there is a coincident albuminuric retinitis. Vision usually improves in 10–18 hours, and is fully restored in about 48 hours. In cases during pregnancy there is usually eclampsia. In uræmic amaurosis the pupils are dilated, but usually react to light. It is probably due to circulation of toxic material which acts upon the cells of the visual centres. The retained reaction of the pupils to light shows that the lower centres are relatively immune. Exophthalmos sometimes occurs in cases of nephritis in which uræmia is present or imminent, and may therefore be of some prognostic significance. It is accompanied by pain and limitation of movement of the eyes, and is probably due to œdema of the orbital tissues.

Hysterical amblyopia, as might be expected, exhibits protean manifestations. It may be unilateral, but is more commonly bilateral. There is usually concentric contraction of the fields, with or without colour defects. A spiral field is very characteristic, *i.e.*, the field continually diminishes while it is being taken, so that it may be finally limited to the fixation point. The patients, however, get about perfectly well unaided, an impossibility in cases of genuine highly contracted fields (*vide* p. 325). The condition is sometimes called anæsthesia of the retina, an undesirable designation. Sometimes there are irritative symptoms—blepharospasm, blinking, lacrymation, &c.—hyperæsthesia of the retina. The pupillary reactions are perfect, affording an invaluable objective diagnostic sign. The prognosis in hysterical amblyopia is good, though treatment is usually tedious. The chief difficulty consists in eliminating organic disease, such as retrobulbar neuritis, injury, embolism of the central artery of the retina, sympathetic ophthalmia, and so on.

Amaurosis fugax is a term given to sudden temporary failure of sight, and is a symptomatic condition due to various causes. In its simplest form it occurs in normal people on rising suddenly from the sitting or recumbent to the upright posture. It is then due to the effects of gravitation upon the blood and is merely momentary, accompanied by slight giddiness and even faintness. Transient blindness, seldom complete, occurs as a prodromal symptom of obstruction of the central artery of the retina, and is probably due to spasm of the arteries or to

the effects of changes in blood pressure associated with arterio-sclerosis. It has been met with in people with mild signs of Raynaud's disease. Temporary amblyopia also occurs in migraine (*q.v.*) and in early stages of papillœdema (*q.v.*) from increased intracranial pressure.

Scintillating Scotomata of various kinds occur in migraine. In typical migraine the patient feels unusually well before the attack. A positive scotoma appears in the field of vision; while obscuring sight it has a peculiar shimmering character. It gradually increases in size until half the field is clouded, the fixation point remaining relatively clear. In the dark field there are often seen bright spots and rays of various colours, and these are often arranged in zig-zags and are then called fortification spectra (*teichopsia*). Both half fields are affected, so that there is homonymous hemianopia. In other cases the whole field becomes clouded, but usually even then the fixation point is seen momentarily, and then becomes obscured until the eyes are moved to a fresh spot. Vision usually clears in about a quarter of an hour. The attack is soon followed by violent headache, generally limited to the opposite side of the head to the hemianopic field (*hemicrania*), and accompanied by nausea and even sickness ("bilious attack"). During the attack there is frequently numbness in the mouth and tongue and slight aphasia. Attacks occur periodically, but vary greatly in number and severity. In mild attacks the scotoma or slight aphasia may occur without the headache and *vice versâ*.

Migraine is to be attributed to vaso-motor changes in the brain. Vaso-dilatation, associated with a feeling of well-being, is followed by vaso-constriction, especially in the occipital lobes. There is often a copious secretion of urine of low specific gravity during the attack. Constriction of the retinal arteries has been described during the scotomatous stage, but it is very doubtful if it ever occurs.

Migraine occurs chiefly in highly-strung people and is undoubtedly accentuated, if not caused by chronic forms of peripheral nerve irritation, especially such as are due to astigmatism, anisometropia, &c. Many cases, but by no means all, have been cured by accurate correction of the errors of refraction and the wearing of suitable glasses. Rest, warmth, and sleep are the best cures for the actual attacks. They can sometimes be warded off or alleviated by aspirin. Nitroglycerin or amyl nitrite have been used, but are not reliable cures.

Occasionally people who suffer from ordinary migraine have attacks in which, without any scotoma, the headache is followed by partial paralysis of the third nerve (*ophthalmoplegic migraine*) on the same side as the hemicrania. Slight ptosis, diplopia, and sluggishness of the pupil reactions continue for some hours and gradually disappear. The paresis is worse and persists longer with succeeding attacks, and has sometimes eventually become permanent. Care should be taken to exclude an organic lesion in cases of this type.

Night-blindness occurs par excellence in retinitis pigmentosa (*q.v.*) and in xerophthalmia (*q.v.*). In rare cases it is a congenital affection (*vide* p. 327). It is also found in endemic form, especially after exposure to bright sunlight in hot countries, *e.g.*, amongst soldiers and sailors. The patients are usually overwrought, as by long marching, or debilitated, as by scurvy, fasting in Lent, &c. The condition generally improves rapidly if the eyes are protected and the nutrition attended to; cod-liver oil is specially indicated. The affection is purely local, due to abnormal retinal fatigue, as is shown by the fact that covering one eye with a bandage during the day has been found to restore sight enough for the ensuing night's watch on board ship, the unprotected eye remaining as bad as ever. Night-blindness is to be attributed to interference with the functions of the retinal rods: these structures appear to be more vulnerable than the cones. Soldiers often complain of it, but not infrequently it is a functional nervous disorder in these cases, associated with other symptoms of neurosis or malingering.

Day-blindness occurs in some cases of congenital amblyopia. It also occurs in all cases of reflex blepharospasm. In less noticeable form it occurs in lesions affecting the conducting paths of visual impulses, such as tobacco, amblyopia, retrobulbar neuritis, and the early stages of optic atrophy. Patients suffering from these disorders often see relatively, and sometimes absolutely, better in a dull than in a bright light.

Coloured Vision is sometimes complained of, and red is the colour usually noticed. *Erythropsia* occurs particularly after cataract extraction if the eyes are exposed to bright light and are over-strained. In these cases it may persist for several hours or days. Objects look red, but the visual acuity is not affected, and no permanent damage results. Patients should be warned of the possibility of erythropsia, as it is somewhat alarming and suggestive of hæmorrhage. It is met with also in snow blindness. Red vision is sometimes complained of by

neurotic hypermetropic children. Chromatopsia also occurs in some cases during the resolution of optic neuritis when the ensuing atrophy is not complete.

Metamorphopsia ; Micropsia ; Macropsia or **Megalopsia ; Photopsiæ** (*vide* pp. 315, 328).

Muscæ volitantes (*vide* p. 306).

Colour Blindness or **Achromatopsia** may be congenital or acquired. *Acquired colour blindness*, partial as in cases with relative scotomata, or complete as in disease of the optic nerve, has been referred to incidentally in treating of the various disorders of the eye in which it occurs. It may also be a symptom of disease of the central nervous system, especially perhaps the lowest parts of the occipital cortex. In most diseases of the retina and choroid, *e.g.*, detached retina, changes in colour perception affect mostly the blue end of the spectrum. Slight diminution in acuity of perception of these rays is caused normally, owing to their physical absorption, by the increase of amber pigment in the nucleus of the lens (*blue blindness*), and this may be abnormally great in sclerosing lenses (black cataract). The effect has been noticed in the pictures of artists in their old age (Liebreich). Slight absorption of rays of short wave-length is normal at the macula, owing to the yellow pigment present here ; it varies in different individuals.

Congenital Colour Blindness occurs in two chief forms, total and partial. The former is very rare and is always associated with nystagmus and a central scotoma. All colours appear grey, of different brightness. The spectrum appears as a grey band exactly like the normal scotopic spectrum (*vide* p. 63), and like it with the maximum brightness at 530 $\mu\mu$. It does not change, except in increase of brightness, when the intensity is increased, but at high intensity photophobia occurs. It is probable that total colour blindness is caused by defective development of cones or their complete absence.

The partial form is seldom discovered unless specially tested for, since the subjects compensate for their defect by attention to shade and texture, combined with experience. Gross cases occur in 3 to 4 per cent. of males, but are rare in females (0·4 per cent.). It is an inherited condition, being transmitted through the female, who is usually unaffected. In most cases reds and greens are confused, so that the defect causes grave danger in certain occupations, *e.g.*, railway signalmen, engine-drivers, and sailors. The red-green cases fall into two chief groups, protanopes and deuteranopes. For the former the red

end of the spectrum is much less bright than for normal people and is often actually shortened. These groups are explained on the Young-Helmholtz theory by the hypothesis that one of the primary sensations (*vide* p. 63) is lacking. They are therefore often said to have dichromatic vision. In the protanopes the so-called red sensation is said to be absent, and they are called red-blind; in the deuteranopes the green sensation is absent and they are called green-blind. On Hering's theory both groups are varieties of red-green blindness, which is explained as due to absence of the red-green substance (*vide* p. 64). In both groups the defects may not be complete and these cases are called protanomalous and deuteranomalous respectively. Since, on the Young-Helmholtz theory, all three sensations are represented, though one is defective, these are said to have anomalous trichromatic vision. It is clear that theoretically there might be other cases of colour blindness due to absence of the blue sensation or the blue-yellow substance, and such have been described, but are very rare (tritanopes).

There are two objects to be aimed at in testing for colour blindness: (1) the exact scientific nature of the defect; (2) whether the subject is likely to be a source of danger to the community. The first is exhaustive and may be the only means of arriving at the second, especially in the anomalous trichromats. In the more difficult cases, besides the simpler tests, more stringent tests with a pure spectrum must be employed. In them only those of large experience will gather any useful information from the names given by the subject to various colours, for these are named chiefly by reference to their various brightnesses and the answers appear to be quite inconsistent. In testing for danger only, it is obvious that the names given to the colours are of value, for if a man repeatedly calls red green or *vice versâ* he is clearly unsuited to be an engine-driver or look-out man on a ship.

Whatever the object in view several tests should be employed. For the spectrum tests the student must be referred to special monographs on colour vision. The following are the chief other tests.

(1) *The Lantern Test.* The subject names various colours shown by a lantern, and is judged by the mistakes he makes. Much here depends upon the size of the apertures of the lantern (*i.e.*, the size of the retinal areas stimulated) and the nature and intensity of the light source. Many lanterns are worse than useless. The best is that used by the Board of

Trade, but this is not available for general use. Edridge-Green's lantern is efficient if used by an expert.

(2) *Holmgren's Wools.* These consist of a large selection of skeins of coloured wools, and the candidate is required to make a series of colour matches.

Test I. consists in presenting to the candidate a pale green sample and telling him to select from the heap of wools all those which seem to correspond in colour. If he is colour blind he will probably select several of the "confusion colours" greys, buffs, straw colour, &c.—as well as greens. He is next given a rose colour (II.): if he matches this with blues or violets he is red-blind; if with greys or greens he is green-blind. He is then given a bright red skein (III.): if he is red-blind he will choose dark greens and browns, if green-blind pale greens and browns. Sir William Abney has recommended the addition of two other tests. IV. is a purple skein: if the candidate is colour blind he will probably select any shade of blue or green, also pinks and greys. V. is a yellow skein: the colour-blind candidate will probably select greenish yellows, light yellow-greens, fawns and pinks. In blue-blindness purples, red and orange are confused in test II.

Holmgren's wools have been much criticised, but if the tests are properly carried out gross defects of colour vision are easily recognised and an expert will be put on his guard in almost every case of even minor defect.

(3) *Stilling's Tests.* These consist of coloured lithographic plates, in which bold numerals are represented in dots of various tints set amid dots of the same size but of tints which are most readily confused with those of the figures by colour-blind people. Normal trichromats can easily read the numbers, some of which are indistinguishable to the various types of colour blind.

(4) *Nagel's Anomaloscope.* This is an instrument in which on looking down a telescope a bright disc is seen, divided into two halves by a horizontal line. One half is illuminated by light of the sodium line of the spectrum (yellow), and this has to be matched by a mixture of red (lithium line) and green (thallium line) in the other half. By turning a screw the relative amounts of red and green in the mixture can be varied. By turning another screw the brightness can be varied.

Defects of colour vision have led to much acrimonious discussion. Their detection may be easy, but is often difficult. No theory which has yet been brought forward is wholly satisfactory, and no single test is infallible.

Malingering.—Cases occasionally occur of men who hope to gain some advantage by pretending to be blind. It is rare for complete blindness to be assumed, and such cases can only be detected by constant watching of the person's behaviour. When one eye is said to be blind, in spite of absence of sufficient objective evidence to account for the condition, the demonstration of malingering resolves itself into a contest of wits between the surgeon and the individual. Many tests have been devised, and several should be employed in each case.

(1) A low concave or convex glass (O. 25 D) is placed before the "blind" eye, and a high convex (+ 10 D) before the "good" eye, and the examinee is told to read the distant types. If he succeeds malingering is proved.

(2) A prism is placed base downwards before the "good" eye and the examinee is told to look at a candle. If he admits to seeing two candles malingering is proved.

(3) The surgeon stands behind the patient and covers the "blind" eye with his hand, at the same time holding a prism of 10 degrees base down before the "good" eye in such a manner that the edge of the prism passes horizontally across the centre of the pupil. Monocular diplopia results. The surgeon then simultaneously removes his hand from the "blind" eye and shifts the prism upwards so that the whole pupil is covered by it. If the examinee still admits to seeing two candles malingering is proved.

(4) While the examinee looks at a candle a prism of 10 degrees is placed base outwards before the "blind" eye. If the eye moves inwards in order to eliminate diplopia it is not blind.

(5) Snellen's coloured types may be employed. The letters are printed in green and red. If a red glass is placed before the "good" eye, and the patient reads all the letters the other eye is not blind, for the eye looking through the red glass can only see the red letters. Care must be taken in this test that the red glass cuts off all the rays from the green letters as tested by the surgeon's own vision.

Word-blindness occurs as a not very uncommon congenital defect—0·1 per cent. of elementary school children (Thomas). It is much commoner in boys than girls. Owing to backwardness in learning to read the children are often brought to the ophthalmic surgeon owing to suspected visual defect. In spite of normal fundi and often quite normal acuity of vision, the patients fail to recognise printed or written words. The auditory memory of words is unimpaired, and generally

numerals and music can be read. Hence the patients learn well orally, and are good at arithmetic. They are often quite intelligent and may be wrongly punished for inattention and stupidity. The condition sometimes runs in families. The defect is not necessarily complete, and much improvement can be obtained by careful individual tuition and perseverance.

CHAPTER XX.

INTRA-OCULAR TUMOURS.

Intra-ocular tumours are rare, but of great importance, since they are usually malignant and endanger the life of the patient.

Tumours of the Iris.—Sarcoma of the iris is extremely rare. It is not uncommon to see irides with dark brown spots (melanomata), due to congenital aggregations of retinal pigment epithelium. As a rule these are benign, but occasionally they take on malignant proliferation. Any increase of size must be watched with suspicion. True sarcoma, composed of pigmented or unpigmented spindle-shaped or round cells, occurs as an isolated nodule. It grows rapidly, and if left attacks the corneo-sclera and perforates the globe. It gives rise to metastases from which the patient dies.

Although it is the only new growth of importance met with in the iris, it may be diagnosed from gumma or tubercle only with difficulty. The chief points of difference are the absence of inflammation as shown by synechiæ, &c., the absence of satellites, the frequency of pigmentation, and the rapid increase in size.

Treatment.—The growth should be watched for a short time and if found to increase in size should be removed by iridectomy if this is feasible. The specimen is subjected to expert microscopic investigation. If the growth is malignant and has not been completely removed or shows signs of recurrence the eye must be excised at once.

Sarcoma of the Choroid is not so rare as sarcoma of the iris. The growth arises from malignant proliferation of the stroma cells of the outer layers. It forms at first a lens-shaped mass, raising the retina over it. By the process of growth increased tension is thrown upon the elastic membrane of Bruch, which finally ruptures. The cells then proliferate through the opening and form a globular " head " in the subretinal space, separated from the mass in the choroid by a narrow " neck " (Fig. 206). The retina remains in contact with the tumour at the summit

of the head, but is detached from the choroid at the sides, the intervening space being filled with albuminous fluid. The growth may be in any situation, and the fluid may sink down to the lowest part of the eye, forming there a " simple " detachment isolated from that over the tumour. As the tumour progresses the retina is more and more detached, until no part remains *in situ*. The nutrition of the lens then suffers, so that it becomes opaque. The tumour may fill the globe before perforating the sclerotic, or this may occur relatively early along the perivascular spaces of the vortex veins or ciliary vessels. The orbital tissues then become infiltrated. The



FIG. 206.—Section of sarcoma of the choroid (× 9), showing the typical mushroom shape. *Ch.*, pars plana of ciliary body, continuous posteriorly with the choroid, behind *R*, retina.

lymphatic glands are not affected, but metastases occur in the liver and elsewhere.

The growth is usually pigmented—melanotic sarcoma, rarely unpigmented—leucosarcoma. The cells are spindle-shaped or round. They may be arranged in columns (cylindroma) or around blood vessels (angio-sarcoma), and are then probably endotheliomata.

The course of sarcoma of the choroid is commonly divided into four stages: (1) the quiescent stage; (2) the glaucomatous stage; (3) the stage of extra-ocular extension; (4) the stage of metastasis. This is probably the typical chronological order of events, but secondary glaucoma may arise at a very early stage or be delayed until after extra-ocular extension has

taken place, and there is evidence to show that metastases may occur at an early stage.

The cause of the glaucoma is obscure: in most cases it is due to the lens and iris being forced forwards, so that the angle of the anterior chamber becomes blocked. In other cases, particularly those of early onset, obstruction to the venous outflow from the eye is the probable explanation, the tumour being in some instances so situated as to press upon a vortex vein.

Sarcoma of the choroid usually occurs in adults between forty and sixty. It is always primary, single, and unilateral. The patient may come under observation in the early stage when there is only detachment of the retina. The earliest cases to seek advice are those in which the tumour is near the macula, since vision is then most strikingly affected. In other cases the tumour has usually attained a considerable size, and the patient may apply for treatment for relief of the pain of glaucoma.

It is of the utmost importance that the cause of the detachment of the retina should be diagnosed in these cases. We have already seen that "simple" detachment of the retina is most frequently found in myopic eyes or after a blow, though many cases are apparently "idiopathic." In simple detachment the lower part of the retina is usually affected, though not always. There is therefore presumptive evidence—of only slight weight—that a detachment in the upper part is due to a tumour of the choroid. In detachments of considerable size the tension of the eye is normal or subnormal in the absence of a growth; if the tension is raised quite definitely a growth may be diagnosed almost with certainty. A simple detachment shows numerous, more or less parallel, folds, and undulations can be seen to travel over the surface when the eye moves. The detachment at the summit of a tumour is usually rounded and fixed, though in the surrounding parts it may show all the signs of a "simple" detachment. Patches of pigment upon the rounded part support the diagnosis of a tumour, but pigment disturbance, more particularly at the periphery, is not uncommon in simple detachment. Rarely a system of blood vessels having an entirely different mode of distribution from the retinal vessels can be made out between the latter vessels. This is the most positive evidence of growth, but it is only occasionally seen. A very small, round detachment in the macular region or upper part of the globe is almost certain to be due to a tumour of the choroid. If the detachment is sufficiently anterior

transillumination with a specially devised lamp will afford assistance in diagnosis, a simple detachment being transparent, a choroidal growth opaque.

Diagnosis may be extremely difficult if the patient is first seen when glaucoma has already supervened. Dependence must then be placed largely upon the history. Defective vision may have been noticed, but the premonitory haloes of glaucoma have been absent, and vision has gone from bad to worse without remissions. One eye only is involved. The other may be perfectly normal, or at least not of the glaucomatous type with small cornea and so on, and the field of vision in this eye will show no contraction on the nasal side. The affected eye will probably have no perception of light, so that if any doubt remains it should be excised.

Treatment.—The eye should be excised as soon as possible after arriving at the diagnosis. Although sarcoma of the choroid rarely travels down the nerve, it is wise to cut it as long as may be. If the growth has already burst through the globe the orbit should be exenterated.

Prognosis is fair if the tumour is small and entirely intra-ocular, but even in these cases metastasis may have occurred. The disease is invariably fatal, usually within five years, if not eradicated by operation.

Flat Sarcoma of the Choroid.—In rare cases the choroid becomes infiltrated with sarcoma cells which cause a uniform thickening and shallow "detachment" of the retina. These are probably endotheliomatous, spreading along the lymphatic spaces of the choroid in the same manner as secondary carcinoma (*q.v.*).

Sarcoma of the Ciliary Body is fundamentally of the same nature and gives rise to the same symptoms as sarcoma of the choroid, the differences being only those dependent upon the anatomical disposition of the parts. Thus the retina, being here more adherent to the underlying uvea, and being reduced to a double layer of epithelial cells, is not detached. When the growth has spread to the choroid the retina proper becomes detached. The tumour may attain considerable size before it causes symptoms, which are then referable to displacement or distortion of the lens and interference with the ciliary muscle. The ciliary circulation is impeded, and conspicuous dilatation of one or two anterior perforating ciliary vessels should always arouse suspicion. The growth may invade the angle of the anterior chamber. It then has the appearance of an iridodialysis, a dark crescent showing at the root of the iris. That it

is not an iridodialysis is shown by the fact that no reflex can be obtained through it on illuminating with the ophthalmoscopic mirror and from the absence of history of a blow. In the case of a leucosarcoma the crescent may be yellowish, but vessels will usually be visible upon the surface, and these render the diagnosis easy. The growth may be visible by oblique illumination with a widely dilated pupil.

Sarcoma of the ciliary body is less common than that of the choroid. The treatment and prognosis are the same.

Ring or Annular Sarcoma of the Ciliary Body resembles flat sarcoma of the choroid in its infiltrating character. It is very rare.

Secondary Carcinoma of the Choroid occurs sometimes in late stages of scirrhus of the breast, rarely in cancer of other organs. There is obscuration of vision, and ophthalmoscopic examination reveals a widespread shallow detachment of the retina. The disease is nearly always bilateral, and as it is usually only one of many metastatic deposits and the patient is generally in the last stages of general carcinomatosis, no special treatment is indicated.

Glioma of the Retina (*Syn.—Neuroepithelioma retinæ*) is a malignant disease confined to infants. It is probably always congenital, though it may remain quiescent or pass unnoticed until the fifth or sixth year or even later. The disease is rare; the second eye is affected, independently and not by metastasis, in about one-fourth of the cases, but frequently the growth cannot be recognised even on careful examination until after months or even years. Several children of the same family are sometimes affected.

The child is brought to the surgeon on account of a peculiar yellow reflex from the pupil, sometimes called "amaurotic cat's eye." If left untreated glioma runs through the same stages as sarcoma of the choroid, viz., (1) the quiescent stage; (2) the glaucomatous stage; (3) the stage of extra-ocular extension; (4) the stage of metastasis. The second stage results in enlargement of the globe, with apparent or real exophthalmos. Pain is severe during this stage, but is relieved when the tumour bursts through the sclerotic. Perforation often occurs at the limbus, and is followed by rapidly fungating growth. Metastasis first occurs in the preauricular and neighbouring glands, later in the cranial and other bones. Direct extension by continuity to the optic nerve (which is early affected) and brain is commoner, and metastases in other

organs, usually the liver, are relatively rare (cf. sarcoma of the choroid). In most cases the first stage lasts from six months to a year.

Glioma is a malignant growth of the retina composed chiefly of small round cells with large nuclei, characteristically arranged around blood vessels, and showing great tendency to degenerative changes (Fig. 207). It is a disease sui generis, and bears no resemblance to glioma of the brain, from which

Fig. 207.—Section of glioma of the retina (× 3). Note the rings of deeply stained cells surrounding blood vessels; also the infiltration of the anterior part of the optic nerve.

it must be carefully distinguished. It probably originates in fœtal, undifferentiated retinal cells: when the other eye is affected it is a separate focus, and not an extension *viâ* the chiasma. It is invariably multiple (cf. sarcoma of the choroid). When seen very early, as for example in the second eye, a larger mass is seen surrounded by numerous punctate satellites. Microscopically, minute deposits are seen scattered in various situations throughout the globe. It may grow principally outwards, separating the retina from the choroid—glioma exophytum, or inwards towards the vitreous—glioma endophytum. There is no fundamental distinction, but the

ophthalmoscopic appearances differ in the two types. In the former the condition resembles a mere detachment of the retina; in the latter polypoid masses may be seen stretching into the vitreous. Hæmorrhages upon the surface are not uncommon, especially in glioma endophytum.

The growth consists chiefly of small round cells with large nuclei resembling the cells of the nuclear layers of the retina; masses of these stain badly, showing that they are undergoing necrosis. Among them may be found rosette-like formations of cells resembling the rods and cones. The growth probably originates in an island of undifferentiated embryonic retina which has failed to develop normally.

Several conditions occurring in children may give rise to similar signs, and cause great difficulty in diagnosis. These have been grouped together under the term *pseudoglioma*. The chief are (1) inflammatory deposits in the vitreous, with or without detachment of the retina; (2) tubercle of the choroid, especially the confluent type; (3) congenital defects, due to persistence of part of the fibrovascular sheath at the back of the lens. The first group are due to a quiet form of cyclitis, which may have passed unnoticed or may have caused slight redness of the eyeball with inflammatory deposits in the anterior chamber; these, however, are usually soon absorbed. A history of fits, unconsciousness, attacks of screaming, ear disease, meningitis (especially post-basic), one of the acute specific fevers, syphilis, &c., may be obtained. Iritis, or the results of iritis or iridocyclitis, *e.g.*, posterior synechiæ, retraction of the base of the iris, and so on, are often present.

In all cases atropin should be instilled and both eyes should be thoroughly examined ophthalmoscopically, under chloroform if necessary. The tension may then be satisfactorily tested, and may afford useful information which cannot be obtained without an anæsthetic. Raised tension is in favour of glioma, lowered of pseudoglioma. Even when every precaution is taken there is a considerable group of cases in which it is quite impossible to be certain of the diagnosis. Considering that the life of the patient is at stake and that the eye is in any case useless as an organ of sight, these cases should be treated as glioma.

Treatment.—The treatment of glioma is excision of the eye at the earliest possible moment. The optic nerve should be cut long, and the cut end invariably submitted to microscopical examination. If there is any doubt of extension of the disease

to the conjunctiva or orbital tissues exenteration of the orbit is imperative. In cases where the diagnosis is doubtful the eye should be removed, for in inflammatory pseudoglioma the eye is destined to shrink and become unsightly. In no case should both eyes be removed at the same operation, but if one is proved by microscopical examination to be gliomatous and the other contains a similar growth, the second eye should also be excised if consent of the parents can be obtained.

The prognosis of glioma, if untreated, is absolutely bad, the patient invariably dying. The prognosis is fair if the eye is removed before extra-ocular extension has occurred. In the absence of disease of the second eye the patient may be regarded as out of danger if there is no recurrence in the orbit within three years, but the remaining eye should be carefully examined under atropin at frequent intervals for a much longer period. There are several cases on record of cure after removal of both eyes for glioma retinæ.

CHAPTER XXI.

INJURIES TO THE EYE, PANOPHTHALMITIS, AND SYMPATHETIC OPHTHALMIA.

The eye is protected from direct injury by the lids and the projecting margins of the orbit. Nevertheless, it is not exempt from foreign bodies, the action of caustics, contusions by blunt and wounds by sharp instruments.

Foreign Bodies, Burns, &c.

Foreign Bodies, which are usually small—particles of coal dust, emery, steel, &c.—may pitch upon the conjunctiva or upon the cornea. In the former case they cause sudden discomfort and reflex blinking. The foreign body sticks to the palpebral conjunctiva and is liable to be dragged across the cornea, which it excoriates. It may get floated by tears towards the inner canthus, and so into the nasal duct. Very frequently it becomes lodged at about the middle of the upper sulcus subtarsalis (*vide* p. 561), where it is most likely to irritate the cornea, or in the upper fornix. It may occasionally become imbedded in the bulbar conjunctiva. Quite large foreign bodies, such as a grain of corn, may be retained for a long time in the upper fornix and give rise to much irritation and some discharge. They are liable to be overlooked unless the upper lid is everted. They are generally imbedded in a mass of granulation tissue, which may simulate the cockscomb type of tubercle (*vide* p. 175).

Fragments of aniline pencil in the eye cause much irritation and a very unsightly staining. The eye should be irrigated with a weak solution of alcohol, and glycerine drops used (Swanzy and Werner), since these substances are solvents of aniline violet.

Particles of steel and emery are very liable to fly straight on to the cornea and penetrate into the epithelium or substantia propria. Larger particles of steel, or less commonly stone, glass, &c., may perforate the globe (*vide* p. 405). When situated in the cornea they cause great pain and irritation. The pupil is often constricted. If allowed to remain they

expose the cornea to the dangers of infection by organisms in the conjunctival sac and ulceration. This may lead to a small superficial slough being cast off, carrying the foreign body with it. The small ulcer thus formed may heal, but if virulent organisms are present, a spreading ulcer, with or without hypopyon, may develop.

It is not always easy to discover a foreign body upon the cornea. A leash of dilated blood vessels at some spot near the limbus may direct attention to its position. In case of doubt the eye should be cocained and the cornea thoroughly examined under oblique illumination with a loupe. The use of fluorescein will sometimes, but not always, reveal the position. In some cases the foreign body can be detected by reflecting light into the eye with a plane mirror, especially if a convex lens is used to magnify the object (*vide* p. 104).

The wing cases of insects and the husks of seeds may adhere by their concave surfaces to the cornea, usually at the limbus, for several days or even weeks.

Treatment.—Foreign bodies must be removed as soon as possible, and as far as possible with antiseptic precautions. If situated in the lower fornix they are easily removed with a clean handkerchief after everting the lower lid. If not found in this position the upper lid should be everted (*vide* p. 75); the particle will generally be found in its favourite situation, and can be removed in the same manner or by passing the finger over the surface. If it is still not seen the upper fornix should be brought into view (*vide* p. 76) and the particle removed. In case of difficulty, previous application of cocain will materially assist.

If the foreign body is imbedded in the bulbar conjunctiva it should be picked out by a needle after application of cocain. If a discission needle is not at hand a darning needle may be used; it should be passed through a flame first, so as to sterilise it. This little operation is performed in the same manner as removal of foreign bodies from the cornea (*vide infra*). It may be necessary to snip off the small piece of conjunctiva containing the foreign body with scissors.

Removal of foreign bodies from the cornea is effected as follows: The eye is cocained and the patient seated in a chair. The surgeon stands behind the patient and holds the lids apart with the first and second fingers of his left hand, pressing slightly backwards so as to steady the globe. An assistant focusses the light upon the cornea, the patient being told to look in the direction which affords the best view of the particle.

A sterilised spud (Fig. 208) is used to remove it. Only if this fails after repeated efforts should a needle be resorted to. It may be a discission needle, or in default of that a darning needle (*vide supra*). The greatest care should be exercised not to scrape up the epithelium more than is absolutely necessary. Emery and steel particles cause a little ring of brown stain around them, which should be scraped off; if it is not done the patient is likely to return, under the impression that the foreign body has not been removed. If there is any sign of ulceration and *the patient is young, with normal tension*, a drop of 1 per cent. atropin should be instilled, warning being given that the sight will be misty for a few days. If the patient is over forty or has any signs indicative of the possibility of glaucoma arising (*vide* p. 268), atropin should be avoided and the eye should be examined daily for a time. In every case the eye is kept bandaged for a day, and boric lotion is ordered. If ulceration occurs it is treated in the appropriate manner (*vide* p. 194). Special attention should be given to particles of stone, which show a greater tendency than steel, &c., to cause ulceration, probably because steel particles are often hot, and therefore sterile, when they enter the eye.

Fig. 208.—Spud for removing foreign bodies from the cornea.

Occasionally sharp steel and other particles penetrate deeply into the cornea, without, however, perforating. The efforts made to remove them may push them still deeper or even into the anterior chamber. When such an accident is feared, special precautions must be adopted. If the particle is steel and a large magnet, *e.g.*, Haab's or Mellinger's, is available, this method should be tried (*vide* p. 418). It will generally fail, the particle being so small that an insufficient number of lines of force pass through it. In these cases or when the particle is non-magnetisable a broad needle should be passed into the anterior chamber and pressed against the back of the cornea while the foreign body is being removed with a needle. Usher has invented a spatula, curved on its anterior surface to fit the back of the cornea, for this purpose: it is introduced into the anterior chamber through a keratome incision. If the foreign body escapes into the anterior chamber it must be removed by other methods (*vide* p. 416).

Prophylactic Measures.—Foreign bodies in the eye are extremely common in industrial workers, especially in grinding tools, lathe work, &c. Apart from the danger to the sight of the worker, they are a source of great economic loss from loss of time, compensation, &c. They could in most cases be entirely prevented by the use of protective goggles, but it has hitherto been found impracticable to enforce this measure among British workmen. Every attempt should be made by the provision of comfortable goggles and by educative means, such as "Safety First" notices and lectures by welfare officials, to point out the dangers and to encourage the workmen to use goggles.

Burns and Injuries by Caustics.—Burns by hot water or steam, hot ashes, exploding powder, molten metal, &c., and injuries by caustics, such as lime, usually from fresh mortar or whitewash, strong acids and alkalies, &c., endanger the eye chiefly in two ways, viz., by injuring the cornea and by producing symblepharon. Strong ammonia is particularly harmful, causing necrosis of the cornea, hydrochloric acid (spirits of salt) much less so. Many eyes have been lost through cutting open golf balls: the central core often contains caustics (barium sulphate, caustic soda, etc.) which spurt into the eye. Immediately after the accident there is intense conjunctivitis and chemosis, but the cornea looks clear; in this stage it is difficult to be certain of the extent of the injury. A drop of fluorescein solution will reveal the extent of the area denuded of epithelium. Prognosis should therefore be guarded, care being taken to impress upon the patient the gravity of the injury and the necessity for constant supervision. In the worst cases the cornea is dull or opaque. In the succeeding days an eschar forms and is thrown off. This is followed by granulation of the injured conjunctiva and frequently by ulceration of the cornea. The corneal condition must be treated like a corneal ulcer (*vide* p. 194). In bad lime burns, &c., the whole cornea may be destroyed, perforation takes place, and the eye shrinks. In less severe cases a dense leucoma forms, porcelain-like in lime burns, and sight is lost. The chief danger derived from the condition of the conjunctiva is that of adhesion of the lid to the globe. It is most likely to occur with the lower lid, the caustic acting principally upon the lower fornix, which is obliterated by organisation of the granulation tissue. The symblepharon thus produced impedes the movements of the globe and may even interfere with its nutri-

tion. Every precaution must be adopted to prevent its occurrence.

Treatment.—In the earliest stages of injury by caustics the excess of deleterious material must be removed. Acids may be neutralised by dilute alkalies (lotio sodii bicarbonatis, 3 per cent.) and alkalies by weak acids (lotio acidi borici) or milk. Particles of lime must be perseveringly picked out with forceps, after previous application of cocain : irrigation with 10 per cent. solution of neutral ammonium tartrate has been advocated. A few drops of ol. ricini or olive oil may be instilled.

To prevent symblepharon a glass rod is well coated with vaseline or boric acid ointment, or if the cornea is involved atropin ointment, and the point is swept round the upper and lower fornices, so that they are well packed with ointment. In severe cases cold compresses should be applied, and the patient put to bed.

In the succeeding days, if symblepharon still threatens, the treatment with ointment and the glass rod is repeated once or more times daily according to the severity of the case. A strip of tin-foil, gold-beaters' skin, guttapercha tissue, or linen, smeared with ointment, may be inserted into the fornix and kept in place by a bandage. If symblepharon occurs it must be suitably treated (*vide* p. 580), but prevention in this case is very much easier than cure.

Contusions by Blunt Instruments.

Injuries by blunt instruments vary in severity from a simple corneal abrasion to rupture of the globe. There is no part of the eye which may not be so injured by contusion as seriously to diminish vision. Moreover, in some cases the changes are progressive, so that *in all cases a very guarded prognosis should be given.* The various conditions which may follow contusion will be briefly enumerated.

Cornea.—A simple *abrasion* may be caused. It is recognised by distortion of the corneal reflex and by the use of fluorescein (*vide* p. 81). There is much pain, like that due to the presence of a foreign body, increased on moving the lids, much lacrymation and reflex blepharospasm. It may become infected and give rise to a corneal ulcer, especially if a mucocele is present (*vide* p. 594). In the simple cases the use of a lotion, *e.g.*, boric acid, boric ointment to prevent the lids from sticking together, and a pad and bandage for a few days suffices (cf. p. 395). Ulceration must be treated suitably (*vide* p. 194).

Recurrent Erosion (*Syn.—Recurrent Traumatic Keratalgia*) is particularly liable to occur after scratches with babies' fingernails. The abrasion, however produced, usually heals quickly, but is followed some days, weeks, or even months later, by acute pain and lacrymation, generally on first opening the eyes in the morning. If the cornea is then stained with fluorescein an abrasion will be found, usually at the original site, but sometimes elsewhere, or there may be a vesicle or group of vesicles. The attack rapidly passes off with appropriate treatment, but often recurs again and again. There is no doubt that in these cases the epithelium is abnormally loosely attached to Bowman's membrane, and is liable to be torn off by the lid on waking. Such looseness of epithelium and formation of vesicles is characteristic of lesions of the fifth nerve (*vide* pp. 215, 219), and it is probable that recurrent erosion is due to this cause, though the actual rationale is unknown. Early attacks should be treated in the same manner as a simple abrasion, boric ointment being plentifully applied at night. Instillation of 2 per cent. holocain has been advocated (de Schweinitz). If the attacks are repeated the spots should be curetted and touched with pure carbolic acid (*vide* p. 207).

Deep opacity may be found, usually in the form of delicate grey striæ interlacing in different directions. They are due to accumulation of lymph in the interlamellar spaces, occasionally to wrinkling of Descemet's membrane (*vide* p. 234). They generally clear up without leaving a permanent opacity.

Blood-staining of the cornea occasionally results from contusion which has caused hæmorrhage into the anterior chamber. Probably in all the cases the tension of the eye is raised in the early stages. The whole cornea is at first stained, the colour varying according to the duration of the condition. It may be reddish-brown, or greenish. In the latter case the condition nearly simulates dislocation of the clear lens into the anterior chamber (*vide* p. 400). The cornea gradually and very slowly clears from the periphery towards the centre, the whole process taking more than two years. Microscopically there are myriads of minute, highly refracting rods packed in the lamellæ of the substantia propria, and sometimes round granules of pigment in the corneal corpuscles. These are derivatives of hæmoglobia, which may or may not contain iron. They are probably removed by the phagocytic action of leucocytes —a slow process. In the absence of other cause of defective vision sight may eventually be completely restored.

Rupture of the cornea is very rare. Descemet's membrane may be ruptured alone. In complete rupture an attempt may be made to save the eye by suture.

Sclerotic.—*Rupture of the globe* is generally due to it being suddenly and violently forced against the orbital walls. It is often due to falls upon some projecting object, such as a knob or a key in a door, and is frequently caused in country districts by a blow from a cow's horn. The force is usually applied from the direction down and out, where the eyeball is least protected by the orbital margin ; the eye is forced into contact with the pulley of the superior oblique muscle. The sclerotic gives way up and in at its weakest part, viz., in the neighbourhood of the canal of Schlemm. The wound is oblique, being farther forwards internally than externally, where it appears more or less concentric with the corneal margin and about 3 mm. behind it. The conjunctiva is often intact, but there are always severe injuries to other parts of the eye. The iris is generally prolapsed or torn away (iridodialysis) or retracted. The lens may be expelled from the eye or escape under the conjunctiva (subconjunctival dislocation of the lens) or be forced back into the vitreous, making the anterior chamber deep. The anterior chamber contains blood (hyphæmia), and there may be hæmorrhage into the vitreous. Detachment of the retina occurs, with or without subretinal or subchoroidal hæmorrhage. The eye usually shrinks and is lost.

Treatment.—The eye must be carefully examined with lid retractors, under an anæsthetic if necessary. In severe cases nothing remains but to excise the collapsed globe. In less severe cases, without extrusion of the contents of the globe, atropin may be instilled, cold compresses applied, and the patient kept in bed. Sometimes good results follow suture of the rupture. If the rupture involves the periphery of the cornea the iris alone may be prolapsed. It is then a good plan to insert the sutures in the sclerotic without tying them, before excising the prolapse ; they are then tied.

In subconjunctival dislocation of the lens it would seem a natural procedure to open the conjunctiva and let out the lens. This is, however, contraindicated in the early stage. It must be remembered that there is an opening directly into the vitreous, and that such a procedure will almost inevitably involve escape of vitreous, and possibly panophthalmitis. Atropin should be instilled and cold compresses applied. The lens will gradually become absorbed, but no harm will accrue if the remnants are removed after the scleral rupture has healed.

Iris.—Most injuries to the iris caused by contusion are due to sudden incurving of the cornea, whereby the aqueous is forced back against the iris and lens.

Traumatic mydriasis may follow a contusion. The pupil is large and immobile, and usually remains moderately dilated permanently. It is due probably to paralysis of the motor nerve fibres, though there are minute ruptures in the pupillary margin. There is usually also paralysis of accommodation. *Traumatic miosis* is rarer, and results from less severe injuries; it usually passes off. *Radiating lacerations* of the iris, sometimes extending to the ciliary margin, are rare (Fig. 209). *Iridodialysis* is commoner (Fig. 210). The iris is torn away from its ciliary attachment for a variable distance. On inspection a black biconvex area is seen at the periphery, and the pupillary edge bulges slightly inwards. With the ophthalmoscopic mirror a reflex can be obtained through the peripheral gap, and the fibres of suspensory ligament and edge of the lens may be visible. Uniocular diplopia may be produced by this injury. In extensive iridodialysis the detached portion of the iris may be completely rotated, so that the pigmented back of the iris faces forwards (*anteflexion of the iris*). The iris never becomes re-attached, but iridodialysis, apart from other injury, rarely causes serious results. In *traumatic aniridia* or *irideremia* the iris is completely torn away from its ciliary attachment, contracts into a minute ball, and sinks to the bottom of the anterior chamber, where it may be impossible to see it. Rarely the same appearance is caused by *total inversion* or *retroflexion* of the iris, the whole iris being doubled back into the ciliary region out of sight. More commonly inversion is partial, so that the appearance of a coloboma (*q.v.*) is obtained, but the fibres of the suspensory ligament cannot be seen. In all these cases there is usually hyphæmia; and other injuries, such as partial dislocation of the lens and so on, may be present.



Fig. 209. — Lacerations of the pupillary margin of the iris and dislocation of the lens, following a blow. (From a drawing by Holmes Spicer.)

The *treatment* consists in rest and the application of cold compresses. Atropin should be instilled in iridodialysis, but not in ruptures of the iris. If there is also subluxation of the lens neither mydriatic nor miotic should be used.

Ciliary Body.— Contusions may cause diminution in the amplitude or loss of accommodation, owing to paralysis of the ciliary muscle. The ciliary body may be involved in rupture

of the globe (rupture, prolapse, &c.), and plastic cyclitis may be induced.

Lens. Subconjunctival *dislocation* of the lens has already been described. The same mechanism which produces the various injuries of the iris may cause the lens to be forced back into the vitreous. The suspensory ligament is then ruptured. The rupture may be complete or partial. When complete the lens may sink to the bottom of the vitreous chamber, where it may be visible as a yellowish mass; occasionally it remains clear and cannot be seen. Partial rupture of the suspensory ligament occurs with subluxation of the lens, which may be displaced laterally with or without some degree of rotation. This leads to irregularity in the depth of the anterior chamber, which is deeper in the part unsupported by the lens. With the pupil dilated the edge of the lens may be seen, as a grey convex line by oblique illumination, but more readily and unmistakably as a black line with the ophthalmoscopic mirror (*vide* p. 104). The want of support to the iris causes tremulousness (iridodonesis) on the slightest movement of the eye, a tremulousness which is limited to the unsupported part.

FIG. 210.—Iridodialysis following a blow. (Nettleship.)

Blows upon the eye less directly from before backwards occasionally cause dislocation of the lens into the anterior chamber. It rarely occurs with a lens of normal size, but not infrequently by quite trivial injury when the lens is shrunken. The clear lens in the anterior chamber is not always easily recognised, but it does not long remain clear, and diagnosis is then easy. It is more globular than normal owing to its freedom from the restraint of the suspensory ligament. When still clear it looks like a globule of oil in the anterior chamber. With oblique illumination it has a golden rim, due to total reflection of the light: this is the exact opposite of the total reflection when the edge of the lens is seen with the mirror, the light being then totally reflected away from the observer's eye. The lens in the anterior chamber causes spasm of the sphincter iridis, which may occur at the moment when it is passing through the pupil. Intense iridocyclitis or secondary glaucoma is then set up. In most cases of dislocation forwards the lens is shrunken, and the suspensory ligament has become partially absorbed; dislocation into the anterior chamber

may occur in these cases spontaneously, without any contusion. Unless the lens is very small, extreme irritation is set up by its presence in the anterior chamber, and the eye is lost if it is allowed to remain.

Dislocation of the lens always causes considerable disturbance of vision. In partial rupture of the suspensory ligament there is astigmatism which is much increased by tilting of the lens. The slackening of the suspensory ligament causes increased curvature and myopia, which may be more than compensated by backward displacement. In total dislocation into the vitreous the effect is that of the old cataract operation of couching; the pupillary area is aphakic, the refraction is highly hypermetropic, requiring cataract glasses for its correction. Vision usually deteriorates gradually.

If the lens is displaced so much laterally that the edge crosses the pupil uniocular diplopia is present. Through the aphakic area of the pupil the eye is highly hypermetropic, through the phakic portion it may be myopic, in addition to which the periphery of the lens acts as a prism. Ophthalmoscopic examination under these conditions shows two images of the disc by the indirect method, differing considerably in size. By the direct method the fundus may be observed through the phakic or through the aphakic portion of the pupil, different lenses being required to correct the refraction in the two cases.

Subluxation of the lens may occur as a congenital condition (ectopia lentis) (*vide* p. 305).

Besides the immediate consequences of dislocation of the lens, very serious remote effects may follow. In subluxation the lens is very liable to become opaque, owing to malnutrition. The pressure of the edge of the lens on the back of the iris and on the ciliary body often sets up severe iridocyclitis, which may lead to the loss of the eye, and even endanger the other by sympathetic ophthalmia (*q.v.*). Secondary glaucoma is a very frequent sequel (*vide* p. 265).

Treatment.—In the absence of irritation vision may be improved in total luxation into the vitreous and in subluxation by suitable glasses. In the latter case it is usually impossible to correct the astigmatism, but sometimes the aphakic part of the pupil can be used. If iridocyclitis or secondary glaucoma is present the lens should be extracted if it is possible. It is imperative when the lens is in the anterior chamber. In all cases it is unusually difficult. There is always a considerable rupture of the suspensory ligament, so that vitreous presents as soon as the corneo-scleral section is completed, and the

delivery of the lens has usually to be effected with the scoop, much vitreous being lost (*vide* p. 452). If extraction is impossible an iridectomy or trephining may improve matters, but more usually fails. If the eye is blind and painful it should be excised. For the treatment of subconjunctival dislocation see p. 399.

Besides dislocation of the lens, *concussion cataract* occasionally follows a contusion. In most of these cases the capsule is ruptured, though the site of rupture, usually behind the equator, cannot be seen clinically. The lens gradually becomes opaque, a rosette-shaped opacity being usually first formed in the posterior cortex. Rarely the cataract remains in this condition, but much more commonly it spreads throughout the cortex until the appearance of a mature cataract is found. More or less absorption takes place, but it may be permanently incomplete. The condition should be treated in the same manner as traumatic cataract.

In some cases a circular ring or disc of faint or stippled opacity is seen at the anterior surface of the lens (Vossius' ring). It usually has about the same diameter as the pupil, and has been attributed to the impress of the iris on the lens, produced by the force of the blow driving the cornea and iris backwards. This view is supported by the not infrequent presence of iris pigment on the lens capsule, but is opposed by the variation in the size of the disc and the occasional presence of a second concentric ring. The opacity is probably due to changes in the anterior capsular epithelium, and often clears up.

Concussion cataract may also be caused by lightning and high tension electric discharges. It is possible that electrolytic changes play some part in these cataracts.

Vitreous.—Hæmorrhage into the vitreous is the commonest effect produced in it by contusion. The vitreous chamber may be filled with blood. In this case no reflex will be obtained with the ophthalmoscopic mirror. With oblique illumination a dull red hue may be seen, especially if the pupil is dilated. The blood may become almost completely absorbed, but cloudy opacities remain. In rare cases "retinitis" proliferans follows from organisation of the clot (*vide* p. 345).

Liquefaction of the vitreous and opacities in it may follow a blow owing to uveitis and defective nutrition, and without hæmorrhage.

Choroid.—*Rupture of the choroid* occurs as the result of contusion. Immediately after the injury the view is obscured by

extravasation of blood. When it has become absorbed the rupture, usually not far from the disc and concentric with it, is seen as a curved white streak over which the retinal vessels pass (Plate IX., Fig. 2). The retina may also be ruptured, but this is exceptional. The edges of the streak are pigmented in the later stages; in the earlier, remnants of blood may be seen. The white appearance is due to the sclerotic shining through. The rupture is generally to the outer side of the disc, and there are often two or three of different sizes, more or less concentric with each other. If the choroid is ruptured near the macula loss of central vision results. If the retina is ruptured also or becomes atrophied throughout its thickness, including the nerve-fibre layer, a large sector-shaped scotoma is produced. Simple ruptures of the choroid in which the macula is not involved cause little impairment of vision. The treatment consists of rest in bed until all extravasated blood is absorbed; atropin and dark glasses, and abstinence from reading, &c.

A contusion may cause choroidal hæmorrhage, which may be small, shown later by patches of choroido-retinal atrophy, or large, subretinal or subchoroidal. The latter can seldom be seen ophthalmoscopically, but are part of more extensive mischief.

Retina.—It has already been pointed out that *detachment of the retina* (*q.v.*) is often due to contusion. *Rupture of the retina* with rupture of the choroid is rare. *Hæmorrhages* into the retina occur; they are usually small, but large hæmorrhages into the vitreous (*vide supra*) are in part derived from retinal vessels.

Commotio retinæ is a frequent result of blows upon the eye. Instead of the normal bright red colour the retina shows a milk-white cloudiness, usually near the papilla and posterior pole and over a considerable area. It is probably due to œdema. It disappears after some days, and vision is usually restored to normal.

Serious changes, however, are apt to occur at the macula, and are easily overlooked immediately after the accident, or it may be impossible to obtain a good view. A "*hole*" may occur at the macula. It appears as a small circular or oval deep red patch, just as if a hole had been punched out. It is caused by cystic degeneration following œdema due to commotio retinæ, and in course of time a complete perforation of the retina develops. Similar ophthalmoscopic signs and pathological changes have been observed occasionally after

iridocyclitis, and associated with arteriosclerosis, albuminuric retinitis, and amaurotic family idiocy (*vide* p. 333). In other cases the macula looks deeper in colour than normal, and in the course of time it becomes pigmented. The spots of pigment are very fine, mostly aggregated near the fovea, with a few farther afield. This *pigmentation* is the sign of serious defect in central vision, which has a tendency to increase progressively.

Optic Nerve.—The optic nerve is not infrequently ruptured or injured in fractures of the base of the skull (*vide* p. 365). Injuries by sticks or knives, &c., penetrating the orbit are rare. Avulsion of the optic nerve is very rare in civil practice (cf. p. 422).

Perforating Injuries.

Perforating injuries may be caused by sharp instruments or by foreign bodies.

A wound with a sharp instrument may penetrate the cornea, the corneo-scleral junction, or the sclerotic; it may pass in for a variable distance, wounding the iris, lens, &c., or pass through the eye.

Wounds of the conjunctiva are common. They heal readily, but the process may be hastened, and the resulting adhesion to the sclerotic lessened by introducing one or more sutures. Polypoid masses of granulation tissue sometimes form on the surface; they should be snipped off with scissors after application of cocain.

Wounds of the cornea may be linear or lacerated. The margins swell up and become cloudy through imbibition of fluid. This facilitates closure of the wound and restoration of the anterior chamber. If small and limited to the centre they heal well unless they become infected. The eye is irrigated with boric lotion or sterile saline, atropin instilled, and a pad and bandage applied. A permanent dense opacity is left, and the contraction of the organising scar tissue causes irregular astigmatism. If the wound becomes infected it must be treated like a perforating ulcer.

The danger is greatly increased (1) if the wound is large, especially if it extends into the sclerotic, or (2) if the lens is also wounded. In the former case prolapse of the iris is almost certain to occur. The prolapsed iris should never be replaced, even if this is possible, since it may carry infection into the eye. It must be excised (*vide* p. 201), care being taken that the iris is quite free, though this is a counsel of perfection not always

to be attained. If in a few days it is found that there is an anterior synechia this should be divided as soon as the wound is sufficiently healed to permit of the necessary procedures without re-opening it. If it is not done the traction on the iris will keep up irritation in the eye, as shown by ciliary injection, &c., and such eyes are liable to cause sympathetic ophthalmia. Moreover, a broad anterior synechia tends to bring about secondary glaucoma or predispose to secondary infection and panophthalmitis. In some cases it is advisable to transplant a conjunctival flap (*vide* p. 201). Atropin should always be instilled (never eserin).

Wound of the lens very greatly increases the gravity of the case, especially in children. It may escape notice at first, especially if the wound is small, *e.g.*, that caused by a needle or thorn. The lens swells and keeps the iris in contact with the cornea, so that re-formation of the anterior chamber is much delayed. If at length it re-forms the aqueous becomes filled with swollen lens fibres, which also irritate the iris. The swelling is greater and more rapid the younger the patient. Infection, which is one of the greatest dangers of all perforating wounds, is particularly likely to occur in these cases. The excision of any prolapsed iris in such a manner as to free it completely from the wound is extremely difficult. The subsequent prolonged contact of the iris with the cornea facilitates the formation of broad adhesions of iris and often also of lens capsule which it may be found impossible to divide. Ciliary injection is kept up, and sympathetic ophthalmia is very liable to supervene if the eye is retained too long.

Occasionally in perforating wounds with a dirty implement pyogenic organisms are carried into the eye, multiply there and cause rapid necrosis of the whole cornea. In these cases a ring of deep infiltration appears in the cornea 2 or 3 mm. internal to and concentric with the corneo-scleral margin—so-called *ring abscess*. There is much chemosis of the conjunctiva and a greenish discharge. The organism has generally been found to be the bacillus pyocyaneus. Usually panophthalmitis is set up and the whole of the central part of the cornea is cast off. The only chance of saving such an eye is to do a paracentesis directly the infiltration is observed and to wash out the anterior chamber with hydrogen peroxide.

Wounds of the corneo-sclera are commonly held to be particularly liable to set up sympathetic ophthalmia (*q.v.*). In them the ciliary body is injured directly, and may be prolapsed. There is the usual danger of infection, and if it occurs panoph-

thalmitis is certain to follow on account of the ready communication with the vitreous. If this does not happen ciliary irritation is kept up. There can be no question that the sterility of the instrument which caused the accident is of the utmost importance, quite apart from actual suppurative infection. It is uncommon for a wound with a sterile instrument to cause sympathetic inflammation, though it cannot be positively denied that such cases may occur. If the lens is wounded the chances of saving the eye are greatly diminished.

Treatment. If the injury is strictly limited, and has been caused by an instrument which is probably sterile, an attempt should be made to save the eye. If the iris or ciliary body, or both, are prolapsed, they must be excised. Some vitreous will inevitably be lost during the operation. The following procedures may be adopted: A general anæsthetic is given, unless specially contra-indicated. The conjunctival sac is washed out with a large quantity of perchloride lotion. A speculum is inserted; it may be kept from pressing on the globe by an assistant. The wound is thoroughly inspected, the opening in the conjunctiva being enlarged if necessary. Fine silk sutures are then inserted in the sclerotic, but are not tied, great care being exercised not to press unduly upon the eye. The prolapsed iris and ciliary body are then excised as in the operation for simple prolapsed iris (*vide* p. 201). The scleral sutures are then tied. Finally the conjunctiva is sutured over the wound. By excising a small triangular flap of conjunctiva on one side of the wound the conjunctiva from the other side can be drawn across so as to cover the whole wound: by this device the wound in the conjunctiva is so placed that it does not coincide with that of the corneo-sclera (Lister). Atropin is instilled, the eye bandaged lightly, and the patient put to bed.

If the eye does not quiet down in the course of a week or ten days, as shown by diminution of ciliary injection and cessation of photophobia and lacrymation, it should be excised. In the interval the cornea is examined most carefully each day by oblique illumination with the loupe for precipitates ("k.p."). If they are seen the eye should be excised. Similar care is devoted to the discovery of spots of "k.p." in the other eye, and if they should be found there excision of the injured eye is still more imperative at the earliest possible moment. If the eye quiets down quickly and there is no evidence of iridocyclitis in either eye the case will probably do well, but

it should be kept under observation for a very prolonged period (*vide* p. 426).

If there is much prolapse of vitreous as well as of iris and ciliary body, or if the lens is wounded, there is little probability of saving the eye. If it is almost certain that useful vision will be lost the risks of sympathetic should not be run, but the eye should be promptly excised.

Wounds of the sclerotic are not always easily recognised. The eye may have been wounded through the lid. The bruising and laceration of the lid may make examination of the eye difficult. The lid should be raised from the globe and drawn back with a Desmarres' retractor, under an anæsthetic if necessary. Even when the eye is examined the effusion of blood under the conjunctiva may render the diagnosis uncertain, still more the question of perforation. When perforation has occurred, there is reduction in the intraocular tension. If the perforation is near the cornea, the anterior chamber is shallow or obliterated. If the wound is large, prolapse of some of the contents of the globe occurs. The uvea—iris, ciliary body, or choroid—are most easily recognised on account of their pigmentation. Very often the gelatinous vitreous can be seen hanging out of the wound. Hyphæma and vitreous hæmorrhage may be present with or without perforation.

Treatment.—If the injury is so severe that there is no likelihood of recovery of useful vision, the eye should be excised at once. If there is a chance of useful vision, or if permission to excise is withheld, the sclerotic may be sutured after excision of prolapsed vitreous, &c. Very small wounds do not require suturing, but the conjunctiva should be cleansed and stitched over them. Atropin should be instilled, and both eyes bandaged. Complete rest in bed is imperative. Such eyes usually shrink, unless indeed panophthalmitis ensues.

Wounds of the lens cause traumatic cataract. Usually the anterior capsule is wounded; when the posterior capsule is torn, the lymph in the vitreous acts in the same manner as the aqueous. A few hours after the injury the lens becomes cloudy in the vicinity of the wound owing to the action of the aqueous upon the fibres, causing them to swell. In most cases opacities rapidly form in the posterior cortex, irrespective of the exact site of the wound. They are at first feathery lines, which follow the natural distribution of the lens fibres. Later a rosette-like opacity is formed which gradually spreads until the whole cortex is opaque. Flocculent grey masses protrude through the opening in the capsule, some of which become free

and sink to the bottom of the anterior chamber. Sometimes the whole chamber is full of white flocculi, and the lens nucleus may escape entire. The masses are gradually dissolved by the aqueous, and pass out with it through the angle of the chamber. In this manner, in young patients, the whole lens, with the exception of the capsule, may be absorbed. Usually absorption ceases earlier through closure of the capsular wound. The enclosed lens fibres become opaque, and discission is necessary to cause their complete absorption ; in adults, with a hard nucleus, extraction may be necessary (*vide* p. 441). The smaller the wound in the capsule, the sooner it is likely to close. In rare cases, with very small, though it may be deep, wounds, the opacity may remain limited to the site of injury.

A certain amount of inflammatory reaction with ciliary injection is always set up, which may be excessive, even in the absence of infection. It will then be found that the intraocular tension is raised. The swelling of the lens forces the periphery of the iris against the cornea and secondary glaucoma supervenes. This tendency is increased by the difficulty with which the highly albuminous aqueous escapes from the eye, especially if the wound in the capsule is large and much of the lens substance has escaped into the anterior chamber. If the secondary glaucoma is not relieved, the sight will be lost by destruction of the optic nerve. The aqueous and swollen lens substance must therefore be let out by a curette evacuation (*vide* p. 444).

Traumatic cataract is deliberately induced in the operation of discission.

In the absence of secondary glaucoma, the condition is treated by rest in bed, atropin, and a bandage. It is of the utmost importance that the pupil should be kept well dilated. If it is not, adhesions will form between the iris and the lens capsule. The tendency to iritis is increased, a ring synechia may form, leading to a more serious form of secondary glaucoma, and in any case a subsequent needling will be made more difficult and dangerous. *Sterile* atropin ointment, 1 per cent., should be used three or four times daily. If the pupil does not dilate satisfactorily, hot bathings should be given every four hours, and in the case of adults leeches may be applied. It is usually necessary to needle the lens capsule in order to obtain an opening through which vision is made possible, and needling may have to be repeated (*cf.* Lamellar Cataract).

Perforating wounds with retention of foreign bodies have special features and dangers. Eyes containing foreign bodies

are particularly liable to set up sympathetic ophthalmia. In cases of perforating wound, the question often arises whether a foreign body has been retained within the eye. The foreign bodies most likely to penetrate the eye and be retained are minute chips of steel, stone, and particles of glass, lead pellets, copper percussion caps, less frequently spicules of wood. In chipping stone with an iron chisel, it is usually a chip of the chisel and not of the stone which enters the eye. Among war injuries penetration of the eye by fragments of the casing of rifle bullets, often containing nickel, of shells, &c., frequently occurs.

The *size and velocity* of the missile are of importance. If the foreign body is large so much damage is usually caused that the eye has to be removed. Very minute particles can, however, penetrate the cornea or sclerotic and lodge in the deeper parts of the eye. The velocity of these small particles must be very high, for the energy needed to penetrate the walls of the globe is considerable, and the relationship of energy to mass and velocity is given by the formula $E = \frac{1}{2} m v^2$. This is a point of more than academic interest. It has been shown that rifle bullets are sterilised by their rapid transit through the air. This fact may account to some extent for the sterility of minute intraocular foreign bodies, though not entirely; for they are not always sterile, and when they are it is often more probably due to being hot when emitted. Their irregularity of shape would render them less easily sterilised. Cæteris paribus, metallic foreign bodies appear to be more commonly sterile than those made of other substances.

The *nature of the foreign body* is very various and affects the diagnosis, pathological condition, treatment, and prognosis profoundly. As regards diagnosis it is only rarely that we can see the foreign body in the eye or have indubitable proof of retention. In the absence of irrefragable evidence our surest test is skiagraphy, but this method has severe limitations. Relatively few substances are opaque to X-rays. Fortunately, by far the greatest number of intraocular foreign bodies are composed of iron or steel, which gives a good shadow. The same applies to lead and metallic fragments from the casing of rifle bullets. Particles of glass commonly fail to reveal their presence in the skiagram, except heavy lead glass, such as some bottles are made of. Very accurate localisation is necessary. Various methods are in use; Mackenzie Davidson's method is usually employed in England.

Sweet's method, dependent upon the same principle, in America. When it is remembered that accuracy of localisation of the order of 1 mm. is essential and that inaccuracy of this amount may lead to the needless sacrifice of an eye it will be recognised how culpable is any carelessness in this respect.

The influence of the nature of the foreign body on the pathological condition set up in the eye is profound and varied. Thus, it was shown by Leber many years ago that copper causes suppuration even in the absence of pyogenic organisms, leucocytosis being set up by chemical action. Clinically, however, this result may certainly be delayed for a very considerable period, possibly owing to the copper—usually a fragment of a percussion cap—being surrounded by a wall of inflammatory material and encysted.

Copper in the lens may cause little reaction. This is further evidence in favour of the usual reaction being chemical, for such changes are very slow in the lens. Another fact supporting this explanation is that even in severe suppuration induced by copper within the eye the inflammation tends to cause shrinkage of the globe and not perforation. Not infrequently, however, the reaction is so severe as to cause expulsion of the foreign body from the eye. The pus formation retrogrades after a time and is not progressive. Only very rarely does a copper particle become encapsuled, with restoration of useful vision. Sympathetic ophthalmia is less likely to follow copper foreign bodies than others, probably owing to the intense reaction (Leber).

Iron is also dissolved by the tissue fluids and sets up the condition known as *siderosis bulbi*. The earliest clinical manifestation is the deposit of iron in the anterior capsular cells of the lens. This is not affected uniformly, but oval patches of the rusty deposit are arranged radially in a ring corresponding with the edge of the dilated pupil. This appearance is pathognomonic. At later stages the iris becomes characteristically stained, first greenish and later reddish-brown. The vision of these eyes, however little affected by the primary injury, gradually fails owing to degenerative changes in the retina and lens.

Siderosis bulbi (Bunge) has been exhaustively investigated. In a typical case brown granules are found in the corneal corpuscles, in the meshes of the ligamentum pectinatum iridis, on the inner surface of the ciliary body, and in the retina. The anterior layers of the iris are impregnated, and in addition to

subcapsular deposits in the lens, the fibres are also stained. The retina shows complete degeneration, and Perls' micro-chemical reaction shows the whole retinal vascular system marked out by blue coloration. There is always intense blue coloration immediately around the foreign body. The pigment epithelium of the ciliary processes, pars ciliaris retinæ, and retina, and sometimes the supporting tissues of the retina, show diffuse staining. The brown pigmented cells which give the blue reaction are found particularly in the angle of the anterior chamber and in the retina, less in the iris, and least in the choroid. They are not bleached by the ordinary methods for bleaching the normal pigment. Hæmorrhage associated with the injury introduces a complication, for it causes hæmatogenous pigmentation giving the iron reaction and distinguished with difficulty from the xenogenous pigmentation due to the foreign body.

The chemistry of siderosis bulbi is not yet fully understood. E. von Hippel says that the iron is dissolved by the carbon dioxide of the tissues and is fixed by cells which have a specific affinity for the metal; it then becomes oxidised. It has also been suggested that the iron is dissolved by acid phosphates in the intraocular fluid, or that iron may enter into solution in organic form as an albuminate or in combination with an organic acid. The brown precipitate in the tissues is almost certainly produced by oxidation, but it is not a simple oxide or hydroxide, as it is only very slightly soluble in oxalic acid (McMullen).

The characteristic ring of brown spots under the lens capsule is caused by deposition of iron in circumscribed aggregations of newly proliferated capsular epithelial cells. Leber showed experimentally that the introduction of a particle of iron into the vitreous causes extreme degeneration of the retina. Peculiar large granular cells are found which are derived for the most part from the retinal pigment epithelium.

Metals, other than iron and copper—such as lead, zinc, gold, silver—appear to cause little chemical reaction and usually remain quiescent, becoming more or less thoroughly encapsuled according to their position. Lead becomes coated with the carbonate.

Stone is chiefly dangerous from pyogenic infection, but chemical changes also occur, varying with the nature of the stone.

Glass and porcelain may cause remarkably little reaction, but iridocyclitis and disorganisation of the eye usually occur eventually.

With regard to wood, apart from infection the most characteristic feature is the local irritation produced, resulting

in the formation of dense granulation tissue, studded with so-called foreign body giant cells.

Eyelashes may be carried into the anterior chamber in perforating wounds of the cornea, whether accidental or operative: and caterpillar hairs may penetrate the globe (*vide* p. 178).

More important numerically than the chemical changes are those due to infection, and though these are not, strictly speaking, due to the nature of the foreign body, it is undoubtedly true that certain types of foreign body are more apt to give rise to suppuration than others. Much may be learnt in this connection from the analogy of hypopyon ulcer. The common hypopyon ulcer is due to pure or mixed infection with pneumococci, and it is notorious that it is more likely to be caused by fragments of stone, wounds with the leaves of plants or twigs of trees and so on, than by steel or other foreign bodies. Moreover, it admirably exemplifies the rôle played in such infections by the resistance of the tissues, for the patients are usually either old and debilitated or alcoholic. With regard to intraocular infections, it is to be borne in mind that the lens substance and the vitreous form excellent culture media, and further, that even saprophytic organisms, like bacillus subtilis, are capable of setting up a suppurative inflammation in the eye. Probably the commonest pyogenic organism in the interior of the eye is, however, the pneumococcus.

The foreign body may pass through the cornea or the sclerotic. The wound of entry may be extremely minute. The patient may even be unaware that a foreign body has penetrated the eye. If it has passed through the cornea, the minute wound or scar can always be found by careful examination with oblique illumination and a loupe. It may escape detection in the sclerotic.

The foreign body may be retained in the anterior chamber. Here it may fall to the bottom of the chamber, and if very small be hidden by the sclerotic. It is generally, however, caught in the iris, and can be recognised with a loupe. A piece of glass in the anterior chamber is exceptionally difficult to see, on account of its refractive index differing so little from that of the surrounding media.

The foreign body may pass into or through the lens, either by way of the iris or of the pupil. In each case a traumatic cataract is produced, which undergoes the usual changes (*vide* p. 408). If the particle has passed through the iris there

will be a hole in this structure. If the case is seen very early or very late, the hole looks black by oblique illumination, but shows a red reflex when illuminated by the ophthalmoscopic mirror. In the intermediate stage the cataractous lens behind the hole prevents a red reflex from being seen. A hole in the iris is of great diagnostic significance, since it rarely occurs except as the result of perforation by a foreign body. The foreign body may be visible in the lens, either before or after dilatation of the pupil. It is possible for a foreign body to pass through the iris and through the circumlental space without wounding the lens.

The foreign body may be retained in the vitreous. Access to the vitreous by the foreign body may be given by various routes: through the cornea, pupil, and lens; through the cornea, iris, and lens; through the cornea, iris, and zonule; or through the sclerotic. Hildebrand found these four routes represented in forty-three cases by 6, 16, 6, and 15 respectively. The particle may pass quite through the globe into the orbit, remain near the site of entry, become imbedded in the opposite wall, rebound from it, or be suspended in the vitreous. In the latter case it eventually sinks to the bottom of the vitreous chamber owing to degenerative changes in the humour, which lead to liquefaction, partial or complete. Sometimes air is carried in and appears as bubbles in the vitreous. If the particle is small, the lens clear, and there has been little hæmorrhage, the body may be seen ophthalmoscopically in the vitreous or retina. The track through the vitreous looks like a grey line. The foreign body, generally black, and often with a metallic lustre, is surrounded by white exudate and red blood-clot. If the particle has been long *in situ* it may become more or less encapsuled, a small white area of fibrous tissue being seen with dense masses of black pigment in and around it. Fine pigmentary disturbance at the macula may follow, irrespective of actual injury to this region, and indeed merely as the result of concussion. More extensive degenerative processes also occur in the retina, which may become detached. Encapsulation is often rapid with iron, and useful vision may persist for an indefinite time. Particles more than 1—2 mm. in size are almost certain to lead to the destruction of the eye. In the absence of sepsis, siderosis bulbi is the almost inevitable cause of destruction. An encapsuled foreign body may become free after a long period of quiescence. The encapsulation of foreign bodies in the retina depends largely upon their asepsis. The amount

of cicatricial tissue formed in the early stages is inversely proportional to the amount of necrosis, which depends chiefly upon bacterial invasion, though mechanical injury and chemical action must also be taken into account.

Retinal degeneration attacks the macula or is generalised. In the former group yellowish-white spots appear in the region of the fovea, and pigmentation may also occur. Serious disturbance of vision results and is not recovered from. Generalised retinal degeneration takes the form of pigmentation, resembling that of retinitis pigmentosa, and may be preceded by night-blindness.

The prognosis is always bad. It is least bad if the foreign body is in the anterior chamber and the lens is not wounded. The eye may be saved if it is in the lens, especially in young people, whose lenses are capable of becoming completely absorbed. The prognosis is far better when the lens is not wounded, and this is still more markedly the case the younger the patient. With a small foreign body the wound of entry is so small that prolapse of iris rarely occurs, but if the lens is wounded, and more particularly if the patient is young, so that the swelling of the lens is excessive, the iris is pressed forwards against the cornea, anterior synechia is readily formed, and the obliteration of the angle of the anterior chamber easily leads to secondary glaucoma. If the iris is incarcerated in the wound in these cases the division of the synechia presents considerable technical difficulties, the iris and ciliary body are kept in a state of irritation, and the dangers of sympathetic ophthalmia are greatly enhanced. The satisfactory evacuation of the lens substance is much more difficult than in ordinary discission of the lens, so that the continued apposition of the wounded and inflamed iris to the cornea is scarcely to be avoided. Moreover, in my opinion children are very decidedly more susceptible to sympathetic ophthalmia than adults.

Though the dangers of wound of the lens are diminished in older patients, there is no doubt that the prognosis is rendered graver by this complication. However satisfactory the result may be from the purely surgical point of view, an aphakic eye is materially less useful than one which retains a normal lens.

Treatment.—It is a rule that the foreign body should be removed. There are rare exceptions to this rule, more particularly if (1) the foreign body was probably sterile; (2) little damage has been done to vision; and (3) the process of removal will almost inevitably destroy sight. These condi-

ditions are most often fulfilled in the case of minute foreign bodies in the retina.

Magnetisable foreign bodies are more easily removed than others, since the small or the large electro-magnet can be brought to bear upon them.

A chip of steel free in the anterior chamber is removed in the following manner by the hand magnet (Fig. 211). A general anæsthetic should usually be given owing to the iritis already present. After washing out the eye a keratome incision is made above, 3 mm. from the limbus. The keratome is pushed straight on until the incision is of the required size, remembering that the internal wound in the cornea is smaller than the external. The keratome is removed moderately quickly but smoothly, so as to prevent loss of aqueous.



Fig. 211.—Small magnet, with poles of various shapes.

The pole of the small magnet is then placed over the foreign body, outside the cornea, and moved towards the wound, thus dragging the particle along the back of the cornea. The posterior lip of the wound is then depressed with the pole and the foreign body is drawn towards it, out of the wound. If much aqueous is lost a flat pole should be used, and it must be passed across the anterior chamber into contact with the steel, which is then withdrawn. In this case it is more likely to become entangled in the iris, and considerable difficulty may be experienced in freeing it. All the preparations for dealing with a prolapse of iris must be made (*vide* p. 201).

A foreign body on the iris may sometimes be removed with iris forceps through a similar incision. If it is entangled in the iris tissue, the iris must be drawn out of the wound and the part containing the foreign body excised with de Wecker's scissors, as in the operation for iridectomy (*vide* p. 431).

If the foreign body is in the lens, a few days should be allowed to elapse for the aqueous to act upon the lens fibres. A curette evacuation (*vide* p. 444) is then performed and the foreign body will probably be evacuated with the lens matter, or it may be removed by forceps, or if magnetisable, by the small magnet. In a young subject it may be advisable to increase the opening in the capsule by discission a few days before doing a curette evacuation. In these cases a subsequent needling will often be required to obtain a clear opening in the pupillary area of the capsule. It should be done in exactly the same manner as after discission for lamellar cataract. If the patient's other eye is normal he will not be able to wear the correcting glass for the aphakic eye, but his field of vision will be increased on this side. A more cogent reason for operating is found in the fact that it is easier to obtain a clear opening at this period than after months or years, when the capsule will have become thickened and very difficult to divide. In elderly patients it may be necessary to extract the lens by the operation for senile extraction (*vide* p. 445), but the large amount of soft lens matter will increase the dangers of the operation.



Fig. 212.—Haab's giant magnet.

If the foreign body is in the vitreous or retina it is practically impossible to remove it without very seriously damaging the eye unless it is magnetisable. The treatment then lies

between leaving it alone and watching the case, or excising the eye on account of the damage done or the danger of sympathetic.

If the substance is iron it may be possible to remove it with

Fig. 212*a*.—Removal of intraocular foreign body by Haab magnet: first stage. Drawing the foreign body into the anterior chamber, the eye looking forwards (Goulden & Whiting).

the large magnet (Haab's Giant Magnet or Mellinger's Ring Magnet). This will be facilitated by knowing its position, either by direct observation or by skiagraphy; some hint may be obtained from the position of the wound of entry and the probable direction in which the foreign body was travelling. If the patient is seen immediately after the accident it is best

to use the large magnet at once, without waiting for a skiagram. The longer the foreign body is left the more firmly it becomes imbedded in exudates or fibrous tissue, and the less the probability of its successful removal.

FIG. 212*b*.—Removal of intraocular foreign body by Haab magnet: second stage. Drawing the foreign body across the anterior chamber from behind the iris (Goulden & Whiting).

The patient is seated in front of the magnet if Haab's instrument is used (Fig. 212*a*). The eye is brought close to the magnet, the cornea touching the pole. This is important, since if there is a large piece of iron in the eye and the cornea is not in contact with the pole, the whole eye may be drawn forward out of the socket when the current is turned on. Moreover,

the direction of the core of the magnet should coincide as nearly as possible with the direction of entry of the foreign body. The current is then turned on: some pain is usually



FIG. 213.—Mellinger's ring magnet.

felt when a piece of steel is present within the eye. If the operation is successful the foreign body comes forward into the anterior chamber. It may be necessary to turn the current on and off several times before this happens. If the particle is

small and deeply imbedded the operation is likely to fail; hence it should be performed as soon as possible after the injury, before the chip has become firmly encapsuled or imbedded in exudates. The foreign body often passes round the lens, not through it. It will be seen bulging the iris forwards. The current should then be turned off and the eye adjusted so that the particle will be drawn towards the pupil (Fig. 212*b*). When it has fallen into the anterior chamber the patient is transferred to the operating table and the foreign body removed with the small magnet in the manner already described.

In some cases when the foreign body has been accurately localised and the large magnet has failed to bring it forward it can be removed by the small magnet introduced through a scleral incision made as nearly as possible over the site.

The immediate effect of extraction of foreign bodies with the large magnet is often good, but irreparable damage is often done to the eye. The tracks through the vitreous often become filled with fibrous tissue. As this organises and contracts the retina is pulled up, and total detachment destroys vision. Or more severe iridocyclitis may be set up and the eye shrinks.

The ring magnet has the advantage that the patient lies upon the operating table throughout. When a current passes round a solenoid a magnetic field is generated, its greatest saturation being in the central axis of the solenoid. The ring is placed over the patient's head (Fig. 213), which is arranged so that the affected eye is as near the centre as possible. When the current is turned on every steel instrument placed within the ring becomes a magnet, the force varying with the mass of metal in the instrument and its position in the ring. The foreign body is drawn forwards, as already described, by means of rods of soft iron of various sizes. The smallest should be used first, and the largest should be firmly grasped, otherwise they will be dragged out of the hand and thus do damage. As soon as the foreign body is seen to bulge the iris forwards the position of the rod is altered so as to draw it through the pupil. It may then be withdrawn by the small magnet or a keratome incision can be made, and the smallest rod or a steel spatula can be used in the same manner as the hand magnet.

War Injuries.

The injuries inflicted upon the eye and visual apparatus in war differ somewhat in character from those met with in civilian practice. They are due either to direct injury by a penetrating foreign body or by concussion: in many cases it is impossible

to determine the exact mode. Foreign bodies include bullets, shrapnel shots, fragments of shells, minute metallic particles, and earth, gunpowder, dust, &c. Bullets usually cause clean perforating wounds, boring a hole straight through the tissues, unless deflected by the stronger bones of the face and skull. They are responsible for many cases in which the orbits or face are perforated from side to side in various directions. If the eye is touched it is usually irretrievably damaged, the cornea being often turned downwards as a flap. The bullet not infrequently passes behind the eyes from side to side, dividing both optic nerves and ploughing through the nasal fossæ, or it may pass similarly below the eye, involving the maxillary antrum and nose. The eyeball itself is then usually injured by concussion, probably by the cone of compressed air which is driven forwards by the projectile. There are then usually extensive retinal hæmorrhages in the part nearest to the wound track, followed by scarring and pigmentation of the retina. Shrapnel shots have usually less velocity and are more liable to be retained in the orbit, eye, or neighbouring parts. The same applies to large jagged fragments of shells and to minute particles of metal. Since it is impossible to be certain that any eye injured in warfare does not contain a foreign body a skiagram should always be taken, and if any such body is found it must be accurately localised. Many of the smaller fragments are non-magnetic. The cases must be treated on the principles already enunciated.

Severe injuries are caused by concussion. Traumatic cataract from this cause is not uncommon, and subluxation of the lens, sometimes in both eyes, seems to be more frequent than in accidents in civil life. The ordinary rupture of the globe, concentric with the margin of the cornea, appears to be very rare. Some of the changes in the fundus are of great interest. Mention has already been made of hæmorrhages and subsequent retinal pigmentation. Ruptures of the choroid of the ordinary type are common. Ruptures of the choroid *and* retina, which are rare in civil life, are quite frequent. They are followed almost invariably by organisation of blood clot and exudate and formation of large masses of fibrous tissue. This proliferation of fibrous tissue is probably due to the clot being brought into contact with fibroblasts derived from the choroidal mesoblastic tissue (*vide* p. 345). Detachment of the retina is rarely seen as the result of war injury. Partial avulsion of the optic nerve, which is very rare in civil life, is not uncommon in cases where a projectile passes behind the eye. The disc is seen to be torn on one side and covered with hæmorrhage ; often the true nature of the case is revealed only when the blood has become absorbed. Extensive vitreous hæmorrhages are of course very common.

It is probable that the concussion in these cases is chiefly due to air pressure, either by the explosion of high explosive shells in the immediate neighbourhood or by the compressed air in the immediate vicinity of a rapidly travelling projectile. The effects of concussion on the brain are modified by the frightfulness of the conditions of modern warfare, and visual symptoms play a large part in the traumatic neurasthenia of "shell shock." The commonest symptom is functional blindness, lasting for a variable period, and often recovered from suddenly. Patients often complain of being able to see only a foot or two in front of them. In some cases there are nystagmoid jerks, and spasmodic convergence (kinetic strabismus) associated with spasm of accommodation is not uncommon.

Defects of movement from direct injury to the nerves and muscles in the orbit are common. Proptosis occurs from intra-orbital hæmorrhage and venous thrombosis, and enophthalmos may result from retraction of the scar tissue and fracture of the walls of the orbit. The projectile not infrequently passes through, or in close proximity to, the cavernous sinus; and wounds of the occipital lobes, causing homonymous hemianopia, uni- or bi-lateral, are common.

Panophthalmitis.

Panophthalmitis is generally caused by infected wounds, whether accidental or the result of operations, and ulcers. Less frequently it is metastatic, accompanying pyæmia and puerperal fever, meningitis, and orbital cellulitis (*vide* p. 318). In the exogenous form the vitreous is usually first affected; organisms grow in it as in a culture medium, and purulent cyclitis, retinitis, and choroiditis are set up (Fig. 214). In the endogenous forms there may be a septic embolism of a retinal artery or the choroid may be first affected. In this group it may be bilateral.

In both forms there is rise of temperature, headache, drowsiness, and sometimes vomiting. In the exogenous forms the edges of the wound become yellow and necrotic, hypopyon appears, there is great chemosis, with intense ciliary and conjunctival congestion, and the lids are swollen and red. There is severe pain in the eye, due at first to iritis, later to increased tension. The vitreous becomes purulent, as shown by a yellow reflex by oblique illumination. The anterior chamber soon becomes full of pus, and the cornea cloudy and yellow: ring infiltration may occur (*vide* p. 406). There may be exophthalmos and limitation of movement of the globe, due to extension

of the inflammation to Tenon's capsule. In the metastatic cases rapid failure of vision, a yellow reflex, and hypopyon are found in the early stages.

If the case is left to take its course the pus bursts through, usually just behind the limbus. The pain subsides, and after prolonged suppuration the eyeball shrinks.

The prognosis is bad, the eye being almost invariably lost. The condition is not likely to set up sympathetic ophthalmia.

The pneumococcus is responsible for most cases of panophthalmitis, but it is also caused by staphylococci, streptococci,



Fig. 214.—Section of the retina and choroid in panophthalmitis (× 60), following a perforating wound.

bacillus coli, bacillus pyocyaneus (*vide* p. 406), and even by saprophytic organisms such as bacillus subtilis.

Treatment.—In the early stages after operation wounds an attempt must be made to stay the process. The edges of the wound are cauterised with pure carbolic acid or the actual cautery. The anterior chamber may be washed out with hydrogen peroxide solution, and iodoform introduced. Hot fomentations and leeches are applied, and atropin instilled.

As soon as it is evident that the eye cannot be saved it

should be excised. This should not be left too long, and great care should be taken that no undue pressure is put upon the eye. If pus escapes there is danger of purulent meningitis being set up, the patient's life being thereby endangered. On this account many surgeons refuse to excise these eyes. They open the eye freely by a crucial incision and wait until all signs of suppuration have disappeared before excising the shrunken remnants of the globe.

If pus escapes during excision of an eye with panophthalmitis, the globe should be opened freely and the contents scooped out. The interior of the sclerotic is then swabbed out with perchloride lotion (1 in 2,000) and the conjunctival sac irrigated with a large quantity of weaker lotion (1 in 5,000). The sclerotic may then be excised at once, the risk of meningitis being small, or excision may be postponed.

Sympathetic Ophthalmia.

Sympathetic ophthalmia (or sympathetic ophthalmitis) is the much dreaded condition in which serious inflammation attacks the sound eye after injury of the other eye. In recent years sympathetic ophthalmia has become a rare disease, in spite of the fact that ophthalmic surgery has become more conservative. Though common in the American Civil and Franco-Prussian Wars, it was very rare during the Great War. This gratifying fact is due to increased skill in the treatment of perforating wounds, particularly in the application of antiseptic principles. A perforating wound, especially if a foreign body is retained within the eye, is, however, a source of great anxiety to the most experienced surgeon.

Sympathetic ophthalmia almost always results from a perforating wound, especially such as is caused by a foreign body which remains within the eye. Wounds in the ciliary region—the so-called "dangerous zone"—involving the ciliary body and leading to its incarceration in the scar, have always been considered specially dangerous; it is doubtful if, *per se*, they are more dangerous than others. On the other hand, it is certain that wounds in which iris, ciliary body or lens capsule is incarcerated are more likely to set up sympathetic ophthalmia than others. If suppuration supervenes sympathetic ophthalmia is very unlikely to follow; hence perforating ulcers very rarely cause it. It is also extremely rare without perforation, if indeed it ever occurs under these circumstances.

Children are particularly susceptible, but it occurs at any

age. It usually begins four to eight weeks after the injury to the first eye (the exciting eye) has taken place, rarely earlier, but the onset may be delayed for many months or even years —it is said as many as 40 years.

There is always iridocyclitis in the exciting eye. Usually it is a plastic iridocyclitis which has been set up by injury and has not subsided in the course of three or four weeks. Instead of quieting down the ciliary injection remains, there is lacrymation and the eye is tender: special attention should be directed to the presence or absence of precipitates ("k.p.") on the back of the cornea. In the rarer cases of delayed sympathetic ophthalmia the exciting eye has passed into a quiescent state. It may have shrunken completely. The onset of sympathetic ophthalmia in the second eye is then often ushered in by return of irritation—ciliary injection, tenderness, &c.—in the shrunken globe. The exciting eye, whilst showing evident traces of old iridocyclitis, may yet possess useful vision.

Sympathetic ophthalmia—the disease in the second or sympathising eye—is almost always a plastic iridocyclitis differing in no respect from this form of iridocyclitis due to other causes. In rare cases it manifests itself as a neuro-retinitis or choroiditis. In cases which the surgeon knows to be liable to the condition the first sign may be the presence of precipitates ("k.p.") on the back of the cornea, noticed at this early stage because they have been dreaded and carefully watched for. In other cases the patient first seeks advice for defective vision or inflammation in the uninjured eye (sympathetic irritation).

Prodromal symptoms are sensitiveness to light and transient indistinctness of objects. The latter is due to weakness of accommodation; objects become blurred when doing fine work, but after an interval of rest vision improves. On examination at this stage there may be lacrymation, slight ciliary injection, tenderness of the eyeball, as shown by the patient shrinking from an attempt at examination, precipitates on the back of the cornea, and vitreous opacities. The prodromal symptoms may occur in intermittent attacks, spread over a considerable period.

When fully developed all the signs and symptoms of iridocyclitis (*q.v.*) are present, varying in degree according to the severity of the case. The prognosis as to vision is always doubtful, but if there is much deposition of plastic exudates in the pupillary area it becomes extremely grave. Cases show-

ing little exudation ("serous iritis"), but a deep anterior chamber and "k.p.," have a more favourable prognosis, but they may at any moment develop into the severe plastic type. Tension, difficult to determine on account of tenderness, is moderately raised in the early stages. It may then pass into the condition of lowered tension with gradual shrinking of the globe; or the iridocyclitis may subside, the eye quieting down and retaining fair vision. In the worst cases a ring synechia forms and secondary glaucoma supervenes (*vide* p. 265), or both occlusio and seclusio pupillæ or total posterior synechia (*vide* p. 256) occur and the eye shrinks. Sympathetic ophthalmia sometimes takes two or more years to run its course.

The *pathology* of sympathetic ophthalmia is unknown. The microscopic features in both the exciting and the sympathising eye are the same. In the earliest stages examined there are nodular aggregations of small round cells scattered throughout the uveal tract. In later stages the infiltration becomes diffuse, and epithelioid and giant cells appear; in fact, the condition is scarcely distinguishable from tubercle of the uveal tract. These are merely the signs of reaction to a constant, relatively mild form of irritation. The ordinary signs of uveitis and its consequences are present.

The evidence which has accumulated in modern times tends to show that sympathetic ophthalmia is an infective disease. It is least liable to occur in otherwise likely cases if the wound or the retained foreign body is sterile. On the other hand, it very rarely occurs if actual suppuration has taken place in the exciting eye; possibly this may be due to some specific organism being destroyed by the superabundant growth of pyogenic organisms. It is more likely to occur from retention of shot, a chip of stone, glass, china, &c., than from that of a particle of hot steel, probably because the latter is sterile.

Sterility of an ocular wound is usually judged by a satisfactory course of healing and the absence of suppuration. It is by no means certain that all such wounds are, strictly speaking, sterile. The resistance of the patient's tissues has to be taken into account. With the same precautions a cataract wound may heal readily in a healthy man, but only after prolonged subacute iridocyclitis in a weakly patient. In many of these cases there are reasons for delayed cicatrisation, such as incarceration of the iris, synechia of lens capsule, &c., but the exact mode in which they act is a matter of conjecture.

Various theories have been brought forward to explain the occurrence of inflammation in the sympathising eye. It has

been suggested that severe inflammation in one eye produces a tendency to ciliary irritation in the other eye by some occult means connected with their anatomical and physiological symmetry; there is no evidence to support this conjecture. More probable, *à priori*, is the view that infection travels along the optic nerve *viâ* the chiasma. On this theory one would anticipate neuro-retinitis in the sympathising eye as the most frequent manifestation of the disease, but it is extremely rare. The experiments supporting this theory fail to substantiate it. The most probable theory is that there is a specific organism, which has as yet escaped observation, possibly because it is so small as to be ultra-visible by the microscope, but one which causes general infection through the blood stream. Possibly it is a protozoön which is not stained by any of the methods at present devised: evidence derived from examination of the blood—increase in the number of large mononuclear lymphocytes—has recently been brought forward in favour of this view. It may be conjectured that the organism is harmless to other organs of the body, and that it finds a suitable nidus only in the other eye, and even then only under favourable circumstances. This theory explains best the facts of both ordinary and anomalous cases. When sympathetic ophthalmia supervenes after the injured eye has long been shrunken it may be conjectured that the organism has lain quiescent and encapsuled. The fact that in such cases the shrunken eye is again injured or becomes spontaneously irritable and inflamed shortly before the outbreak of inflammation in the other eye lends colour to this view. Cases in which the injured eye is excised and sympathetic ophthalmia is said to supervene many years afterwards are best explained as ordinary iridocyclitis—a by no means rare disease—occurring quite independently of the injury.

The *treatment* of sympathetic ophthalmia is one of the most difficult problems in ophthalmology, and often demands the exercise of great judgment.

It is, in the first place, prophylactic. In every case of perforating wound, with or without the retention of a foreign body, the question of excision of the eye on account of danger to its fellow arises. It may be assumed as an axiom that *sympathetic ophthalmia never occurs after the excision of an injured eye unless it has already commenced at the time of operation.* Hence, early excision is a positive safeguard against the disease. The injury to the eye may, however, be otherwise trivial, so that restoration of good sight may be possible. The

rule should be to excise any eye which is so injured that it is improbable that useful vision will be regained. In cases where this is doubtful expectant treatment may be adopted for a time. If the eye quiets down quickly it is unlikely to set up sympathetic. What, then, are the chief causes which keep up irritation ? The most important are entanglement of the iris or ciliary body or lens capsule in the wound, and the presence of a retained foreign body. Every effort must therefore be made to free the iris or ciliary body from the wound by excision of any prolapse, followed, if necessary, by division of anterior synechiæ. Upon the success of these efforts the retention of the eye may depend. If they fail, which is most likely to be the case if the lens is also wounded, ciliary injection is certain to continue.

During this expectant period the most careful watch is kept for " k.p." If the eye continues irritable, with ciliary injection, photophobia, and lacrymation, and above all if " k.p." appears, the eye should be excised. It is seldom wise to wait longer than a fortnight unless there are undoubted signs of amelioration. The slightest sign of ciliary irritation or " k.p." in the other eye indicates the necessity for immediate excision of the injured eye. It must be remembered that children are more susceptible than adults. Care must be taken not to confuse a simple conjunctivitis with ciliary irritation.

Even more difficult to decide is the treatment in those cases in which sympathetic ophthalmia has already supervened. If the case is seen early, shortly after the onset of inflammation in the sympathising eye, and if the injured or exciting eye has no useful vision, this useless eye should be excised at once. There is no question that the excision of the exciting eye has a good effect upon the process in the sympathising eye if performed early. At a later stage there is no evidence to show that it exerts any influence at all.

The chief difficulty arises when the exciting eye has useful vision and the inflammation in the sympathising eye is severe. If this is the condition soon after the injury it may be wise to excise the injured eye. If, however, a considerable time has elapsed since the injury, excision of the exciting eye is likely to have little or no influence upon the process. Moreover, in the end the injured eye may have better vision than the sympathising one, for if the iridocyclitis is severe the sympathising eye may be lost in spite of all efforts. Under these conditions, therefore, the injured eye should be retained.

The treatment of the sympathetic iridocyclitis is that of

iridocyclitis in general (*vide* p. 259). In addition to atropin, hot bathings, rest in a dark room, leeches, &c., the patient should be brought rapidly and thoroughly under the influence of mercury. Mercurial inunctions should be pushed, so that salivation occurs within a week, and the patient should then be kept on the border line of mercurialisation for a considerable period. Massive doses of sodium salicylate have proved beneficial in some cases. On the theory that the disease is a protozoal infection the intravenous injection of salvarsan has been advocated : the results have been encouraging. In later stages pilocarpin injections and the administration of iodides may assist in the absorption of exudates, and lead to improvement of vision. Perseverance in these measures, aided by general tonic treatment, is of the utmost importance.

Recently an old procedure, the formation of a "fixation abscess," has been advocated for sympathetic ophthalmia (van Lint and Coppez). One c.c. of pure oil of turpentine is injected subcutaneously in the flank. An aseptic abscess develops and is opened on the seventh day.

Improvement of vision may occasionally be obtained by operation, but no such interference is to be contemplated until all inflammation has subsided and the eye has been quiet for many weeks or even months. In the milder cases an optical iridectomy may do good. In the worse cases, so long as there is perception of light and moderately good projection of light, more desperate operations, such as extraction of the lens, &c., may be justifiably undertaken if the other eye is blind or has been removed.

CHAPTER XXII.

OPERATIONS UPON THE EYEBALL.

Operations upon the Cornea.

Paracentesis for hypopyon ulcer has already been described (*vide* p. 199). Paracentesis for cyclitis is performed in identically the same manner as in the second method there described.

Operations upon the Iris.

Iridectomy, which consists in the excision of a portion of the iris, is performed for the following conditions :—(1) Prolapsed iris ; (2) corneal or lenticular opacities (optical iridectomy) ; (3) glaucoma ; (4) as a preliminary or as part of cataract extraction ; (5) threatening ring synechia (*vide* p. 256) ; (6) ectatic corneal cicatrices ; (7) foreign bodies in or small cysts or tumours of the iris.

Iridectomy for prolapsed iris has already been described (*vide* p. 201).

Optical iridectomy is indicated in some cases of localised opacities of the cornea or lens, very rarely for occlusion of the pupil or subluxation of the lens. In all cases there must be proof or good reason to believe that the light percipient structures are capable of performing their functions. If the patient is old enough to have the vision tested this should show improvement when the pupil is dilated by a mydriatic. The opacities must be localised, and in the case of lenticular opacities there must be good reason to assume that they are stationary.

An optical iridectomy should be as narrow as possible, in order to avoid dazzling and to obtain an approximation to stenopæic vision (vision through a narrow slit, *vide* p. 20). It should not extend to the ciliary border. The site of election is down and in (Fig. 215), but in the case of corneal opacities the clearest region of the cornea must be chosen, unless this happens to be above, in which case the coloboma would be covered by the lid and useless for vision.

Instruments required: speculum, two pairs of fixation forceps, bent keratome, iris forceps or iris hook, de Wecker's scissors, iris repositor. General anæsthesia is only necessary in very young or neurotic patients.

The keratome is inserted at or just inside the apparent corneo-scleral margin, the blade being kept parallel to the plane of the iris. It is pushed on until the incision is sufficiently long. The handle is then depressed, so that the blade lies against the back of the cornea; the danger of pricking the lens with the point is thus reduced to a minimum. The keratome is then slowly withdrawn. The iris forceps are inserted closed, then opened very slightly and the iris seized just outside the pupillary margin: or the iris may be drawn out with an iris hook, which allows a narrower coloboma to be made. The iris is drawn out of the wound and a portion excised with de Wecker's scissors. A slit-like coloboma is made by holding the scissors so that the blades are in the direction of a radius of the iris (Fig. 218). The iris is freed from the wound, unless already free, by the repositor. Sterile atropin ointment is introduced into the conjunctival sac, and the eye is bandaged.



FIG. 215.—Diagram of wound and coloboma in optical iridectomy at the site of election.

Iridectomy for glaucoma has for its object the opening up of a sufficiency of the angle of the anterior chamber to permit of efficient filtration of lymph. It is essential, therefore, that the coloboma shall extend to the ciliary attachment of the iris, and that it shall be broad at the periphery. It has already been pointed out that when the iris is torn away the fracture occurs at the thinnest part, viz., at the ciliary attachment. This will generally happen in iridectomy for acute glaucoma, if the attack is the first or early in the history of the disease. In chronic glaucoma, however—and the same applies to an acute exacerbation occurring in the course of chronic or subacute glaucoma—the periphery of the iris is firmly adherent to the corneo-sclera. When the iris is torn away the fracture will be at the false angle, and filtration of lymph will not be facilitated. The rules usually given as guidance to the correct performance of iridectomy for glaucoma are that the section shall be peripheral and the coloboma wide. From the above remarks it will be seen that a very peripheral section is not very important

in true acute glaucoma, but that it has usually been thought of importance in chronic glaucoma.

In acute glaucoma and in acute exacerbations of the chronic form a general anæsthetic is required, chiefly because the high tension prevents sufficient absorption of cocain to render the cornea—much less the iris—anæsthetic, partly also because the patient's self-control has been shattered by pain and anxiety. In chronic glaucoma, if the tension is not very high, local anæsthesia may suffice.

Instruments required: speculum (Fig. 216), two pairs of fixation forceps, narrow Graefe cataract knife, iris forceps, de Wecker's scissors, iris repositor. Lawford's speculum is preferable to Lang's for iridectomy upwards, cataract extraction, &c., as it interferes less with the field of operation.



Fig. 216.—Lawford's speculum for the left eye.

The surgeon stands above the patient, using his right hand to make the section for the right eye, the left hand for the left eye. After the lids have received a final cleansing with ether soap, &c., and the conjunctival sac has been douched with boric lotion or saline, the speculum is inserted. The fixation forceps are held in the left hand, the knife in the right, care being taken that the back of the knife is away from the surgeon. (It is an extremely awkward accident to introduce the knife with the back upwards.) The eye is fixed close to the limbus below, or down and in for the right eye, down and out for the left; if the latter, care must be taken not to rotate the globe. The knife is held with its plane parallel to the plane of the iris. The point is introduced at least 1 mm.—in chronic glaucoma 2 mm.—behind the apparent corneo-scleral margin. It is inserted at the point corresponding with seven minutes

to twelve on a clock face (Fig. 217). As soon as the point is in the anterior chamber it will look much brighter than the part in the corneo-sclera ; if this is not noticeable it is probable that the knife is badly directed and is burrowing in the cornea. It is passed steadily onwards across the anterior chamber to a spot corresponding with seven minutes past twelve on the dial of a clock, where the counter-puncture is made. In glaucoma, especially acute glaucoma, the anterior chamber is very shallow, so that it may be very difficult to pass the knife across without catching in the iris and wounding the lens, a most disastrous accident. The deepest part of the chamber is at the periphery, and it will usually be found easiest to coax the point of the knife round the periphery, gently pushing the iris away with the back.

As soon as the counter-puncture is made the knife is pushed on until 5 or 6 mm. are exposed. Cutting out is performed by a series of small sawing movements, little pressure upwards being required with the very sharp knife. It is very necessary to use these sawing movements properly, as the sharpest knife fails to cut if it is simply pressed hard against a surface.



Fig. 217.—Diagram of wound and coloboma in glaucoma iridectomy.

In this manner the section through the corneo-sclera is kept at a uniform distance of 1 or 2 mm. behind the apparent corneo-scleral margin. Some conjunctival flap has already been cut at the sides, but the middle of the knife blade is still under the conjunctiva. The edge of the knife is then directed forwards and the conjunctiva cut through by one or two sawing movements.

During these manœuvres care must be taken that the points of the fixation forceps are not pressing into the globe, which is very likely to occur owing to the attention being concentrated upon the section. It may lead to subluxation or complete extrusion of the lens.

The eye is now fixed with fixation forceps, just below the cornea, by an assistant, who also draws the eye downwards. This requires a little skill. If the eye is simply pulled down the wound will gape. It should be rather pushed down and back by the points of the fixation forceps. The surgeon takes the iris forceps in his left hand and de Wecker's scissors in his right. With a sterile swab the assistant turns down the con-

junctival flap so that it lies over the cornea, unless this has been done previously with the back of the knife after completion of the section. The points of the closed forceps are then inserted in the wound and carried to a point half-way between the pupillary and ciliary margins of the iris at a point slightly to the right of the vertical meridian of the cornea. The forceps are opened slightly, and the iris gripped. The iris is pulled out and cut to the right side of the forceps. This stroke makes the right limb of the coloboma. The iris is then drawn across towards the left. By this movement it is torn from its attachment for the whole width of the section. It is then drawn a little back towards the right, so that it may not be jammed into the left angle of the incision. The freed

FIG. 218.



FIG. 219.

FIG. 218, iridectomy with blades of de Wecker's scissors held radially; FIG. 219, with blades held at right angles to the previous position. The former method results in a narrower coloboma, such as is preferable in optical iridectomy and in extraction of senile cataract with iridectomy.

part is then cut off by a second snip of the scissors. This stroke makes the left limb of the coloboma.

The iris repositor is then taken by the surgeon, the eye still being fixed as before. The tip of the repositor is introduced into the wound and insinuated between the cornea and the iris on one side. By a radially directed movement the iris is smoothed out towards the centre of the pupil so that if the edge of the coloboma is caught in the angle of the wound it will be freed. The same manœuvre is repeated on the other side of the wound. This part of the operation is very important, and may be very difficult. Only when the edge of the pupil is in its natural position and looks circular but for the small defect in the upper part is the surgeon convinced that the pillars of the coloboma are free from entanglement in the wound. The conjunctival flap is then turned back over the wound with the iris repositor, care being taken that it is not folded upon itself. The assistant removes the fixation forceps.

Any blood clot is carefully removed with forceps or a swab. The surgeon lifts the speculum away from the eye, at the same time seizing the upper lashes in the middle of the lid. Removing the speculum, he lifts the upper lid over the wound in such a manner as to prevent the lid from displacing the conjunctival flap.



Fig. 220.—Drawing up of the iris after extraction of cataract with incarceration of the pillars of the coloboma.

Both eyes are bandaged. The unoperated eye may be uncovered after two days.

The chief complications which may arise during the operation are hæmorrhage into the anterior chamber (not usually serious, but inconvenient); wound of the lens (often not discovered until opacity develops); severe intraocular hæmorrhage, leading sometimes to extrusion of the lens, vitreous, and even retina.

It may be mentioned that some surgeons use a keratome in this operation. Some also cut off the iris with one snip of the scissors, the blades being directed at right angles to the direction of the forceps (Fig. 219). It is difficult to imagine how this can produce an absolutely peripheral coloboma.

The after-treatment consists in complete rest in bed. The eye is dressed once daily; neither mydriatic nor miotic is instilled unless complications supervene.

The chief complications arising after the operation are extrusion of the lens (due to too large a section), injury to the eye by the patient (usually during sleep) (*vide* p. 454), severe intraocular hæmorrhage, &c. Delay in re-formation of the anterior chamber for several days may happen; though undesirable, it may lead to no ill effects. The wound may bulge, with or without prolapse of iris or incarceration of the angles of the coloboma in the wound. A cystoid cicatrix may result, not altogether undesirable from the point of view of filtration in chronic glaucoma, but liable to arouse iridocyclitis or permit infection and panophthalmitis. Bulging of the wound may be due to partial subluxation of the lens, which may necessitate extraction under grave technical difficulties. Wound of the lens during the operation leads to traumatic cataract, also demanding extraction.



Fig. 221.—Diagram of wounds in anterior sclerotomy.

Iridotomy is section of the iris without excision of any portion. It is employed for making a new pupil when the normal pupil is closed or has been drawn up to the wound of a faulty cataract extraction with incarceration of the pillars of the coloboma (Fig. 220). In such a case it is usually done as follows. A keratome incision 3 or 4 mm. long is made near the periphery of the cornea at the most suitable part, usually the temporal side. The direction of the section should correspond with the position of the proposed puncture in the iris, *i.e.*, it will be approximately radial. This facilitates the opening and shutting of the iris scissors, and minimises the bruising of the lips of the wound. The closed blades, one of which is pointed, of de Wecker's scissors are passed into the anterior chamber. The pointed blade is forced through the iris and passed on horizontally. The blades are then closed, a horizontal slit being made in the iris. This cuts across the stretched fibres, which retract, leaving an oval artificial pupil.

Iridotomy may also be performed by sawing movements with Ziegler's sickle-knife (Fig. 231).

Sometimes the iris can be hooked out through a keratome incision by means of an iris hook, and a piece cut off. This gives a good pupil, but is of course strictly speaking an iridectomy.

The results of iridotomy for artificial pupil are often disappointing.

Iridotomy may be a necessary preliminary to iridectomy in cases of bombé iris. It is then usually done by passing a Graefe knife across the anterior chamber, puncturing and counter-puncturing both cornea and iris. The iridectomy is done before the punctures become closed with exudate, which usually soon occurs.

Division of Anterior Synechiæ is a form of iridotomy. The operation is too technical to be described in detail here.

Operations upon the Sclerotic.

Sclerotomy.—*Anterior Sclerotomy* is performed by some surgeons instead of iridectomy for certain cases of glaucoma. It is more frequently indicated after iridectomy has been performed and has failed permanently to relieve the increased tension. It may also be done in buphthalmia. The section is commenced in exactly the same manner as in iridectomy for glaucoma, though if an iridectomy has already been done it may be advisable to choose some other part of the corneal

limbus. The counter-puncture is made in the same manner, but instead of cutting completely out a bridge of sclera is left (Fig. 212). If the iris presents in either wound it is replaced with the repositor ; no part of the iris is excised.

The object of the operation is to open up the filtration angle. The section must therefore be as peripheral as is consistent with the safety of the ciliary body.

Posterior Sclerotomy is employed by some surgeons as a preliminary to iridectomy for glaucoma. It is especially

FIG. 222.—Beer's discission needle, with lance-shaped point.

FIG. 223.—Discission needle with sickle-shaped edge.

FIG. 224.—Bowman's discission needle, with stop.

FIG. 225.—Saunder's discission needle, with long cutting edge.

indicated as a temporary measure for relieving pressure while preparations are being made for the iridectomy. It has the advantage of causing the anterior chamber to become deeper, so that the subsequent operation is facilitated. Posterior sclerotomy is also indicated in cases of early detachment of the retina, the site of operation being chosen according to the situation of the detachment. In this case it is essential that the retina should be perforated. The operation is sometimes used for extraction of a foreign body in the vitreous with the small magnet (*vide* p. 421).

The operation as performed for detachment of the retina has

already been described (*vide* p. 353). When performed as a preliminary to a glaucoma iridectomy the wound should pass through the sclera between the external and inferior recti.

Sclerectomy.—*Anterior Sclerectomy* is the name given to various operations for chronic glaucoma in which a fragment of the sclerotic is excised. Removal by scissors, as in

Fig. 226.—
Lang's knife.



Fig. 227.—
Trephine.

Fig. 228.—
Elliot's disc forceps.

Lagrange's operation, is now almost entirely replaced by trephining.

Trephining was first used for glaucoma by Argyll Robertson (1876), and reintroduced in 1909 by Freeland Fergus and R. H. Elliot. The details of the modern method have been elaborated chiefly by Elliot. The operation is indicated in chronic glaucoma and infantile glaucoma, and has been recom-

mended in some forms of secondary glaucoma, *e.g.*, that following cataract extraction, anterior staphyloma, &c.

Instruments required: speculum (Fig. 216), two pairs of fixation forceps (ordinary and claw), tenotomy scissors, iris repositor, Lang's knife (Fig. 226), trephine (Fig. 227), straight iris forceps or Elliot's disc forceps (Fig. 228), de Wecker's scissors, needles, and needle-holder. The eye having been cocained and a drop of adrenalin instilled, the conjunctival sac having been douched, the speculum is inserted. The patient is told to look towards his feet, and the conjunctiva is seized with ordinary fixation forceps 8 or 9 mm. above the cornea. A large conjunctival flap is made, almost concentric with the margin of the cornea (Fig. 229): the lower ends of the wound should be well away from the limbus, otherwise filtration is likely to be impeded owing to cicatricial tissue. The flap is dissected down to the upper part of the corneal margin and turned down over the cornea. It is kept stretched in this position with the iris repositor laid horizontally upon it or with the closed fixation forceps. The subconjunctival tissue is divided with the Lang's knife, the utmost care being taken to avoid button-holing the flap. The edge of the cornea is thus clearly defined, and the dissection is carried into the cornea so that the superficial lamellæ are dissected up with the flap for 1—2 mm. The trephine is then applied, so that half the aperture lies on the cornea, the other half on the sclera. The corneo-scleral disc is cut by a few rotatory movements. When the anterior chamber is entered the patient experiences some pain, aqueous escapes, and the pupil is displaced upwards. The trephine is removed, a knuckle of iris protrudes from the wound and the disc is forced out. It usually remains attached by a small hinge. By tilting the trephine slightly forwards, so that the corneal side of the disc is cut rather more deeply than the scleral, it is generally possible to insure that the hinge shall be on the scleral side. Iris and disc are seized with fine iris forceps or Elliot's disc forceps, and both are cut off with de Wecker's scissors. The iridectomy is usually only a peripheral button-hole. During these manœuvres the assistant must keep the flap well stretched downwards so as to avoid it also being button-holed—a rather serious accident. The flap is then replaced and is fixed at the summit by a single suture. The



Fig. 229.

flap is smoothed down, the speculum removed, and the lid lifted over the wound by means of the lashes.

There is a great tendency for iritis to develop immediately after trephining. Hence atropin should always be used on the following day and for several days afterwards.

The chief complications at the operation are making the trephine hole too peripheral, often with consequent escape of vitreous and incarceration of the ciliary processes; button-holing the flap either with the trephine or de Wecker's scissors; escape of the corneo-scleral disc into the anterior chamber; and non-presentation of iris in the wound. The last complication is usually due to slow escape of aqueous owing to the trephine being blunt.

Subsequent complications are iritis, detachment of the choroid (*vide* p. 327), blockage of the wound with iris, ciliary body, lens or vitreous, failure of filtration from dense cicatrisation, &c.

Anterior sclerectomy may be done with punch forceps. This instrument and the technique of the operation have been much improved of late by Holth.

Posterior Sclerectomy with the trephine has been employed instead of posterior sclerotomy in detachment of the retina.

Operations upon the Lens.

Discission or **Needling** of the intact lens should rarely be performed after fifteen years of age; it may be employed up to thirty or even thirty-five, but the nucleus of the lens is then likely to give trouble. It is indicated in most cases of dense lamellar cataract, some cases of congenital cataract, and some cases of high myopia. Discission is used at any age for the division of dense secondary cataract (after-cataract).



Fig. 230.—Diagram of discission with one needle.

Needling of the soft lens in young patients usually requires a general anæsthetic, though it is quite painless under cocain. The pupil must be fully dilated with atropin.

Instruments required: speculum (Fig. 216), fixation forceps, cataract needle (Figs. 222—225). I prefer a needle with a fairly long cutting edge. If the lenticular opacity is scarcely visible in diffuse daylight, it is best to perform the operation in a darkened room with oblique

illumination. The surgeon stands above the patient. The conjunctival sac having been douched and the speculum inserted, the eye is fixed down and in (right eye) with fixation forceps held in the left hand. The needle is introduced just outside the limbus (Fig. 230), *i.e.*, through the conjunctiva and sclero-cornea, in a plane parallel to that of the iris, at a point just above the horizontal meridian of the cornea. It is carried through the anterior chamber until the point reaches the lower part of the pupil. The handle is then slightly raised, so that the point just perforates the lens capsule. The handle is then moved so that it and the point move through arcs of circles which have their centre at the spot where the shaft is engaged in the corneo-sclera. Having thus made a curved, more or less vertical incision in the capsule, a second incision is made at right angles to it. This is done by very slightly withdrawing the needle so as to disengage it. It is then passed farther on towards the left side of the pupil. The handle is again slightly raised, and at the same time rotated, so that the cutting edge is brought in contact with the capsule. As the needle is slowly withdrawn a straight incision is made in it in a horizontal direction. When this is sufficiently large the handle is depressed. The handle is rotated so that the plane of the blade faces upwards, and the needle is quickly withdrawn from the eye. By withdrawing it quickly no aqueous should be lost. If much aqueous is lost, anterior synechia may result. Sterile atropin ointment is introduced into the conjunctival sac, and both eyes are bandaged.

FIG. 231.—Ziegler's knife.

The most important point about after-treatment is keeping the pupil well dilated, which is done by atropin ointment three or four times a day. There is always some ciliary reaction. The amount of swelling of the lens fibres depends upon the size of the incisions in the capsule, but also varies with different lenses. The reaction is often very slight in cases of true congenital cataract; in these cases the iris scarcely responds at all to atropin. If it is particularly desired to avoid the necessity of a subsequent curette evacuation the incisions should be quite small in the first operation. If it is intended to perform a curette evacuation (*vide* p. 444) the incisions may be as large as possible, and the needle may even

be introduced moderately deeply into the lens and the fibres broken up. In these cases, and occasionally when it is not anticipated, there is great swelling of the lens; the anterior chamber becomes filled with flocculent masses; there is intense ciliary injection accompanied by raised tension and pain. The tension sometimes subsides in a few hours with leeching, but if not must be relieved at once, or the sight will be seriously damaged by secondary glaucoma. A curette evacuation is then done (*vide* p. 444). In cases in which no curette evacuation is performed a second and third needling will usually be necessary before a clear opening is obtained. Intervals of several weeks or even months may elapse until the absorption set up by the previous operation is complete. The final needling will be of the type of a discission for secondary cataract.

Since time is of little importance in the case of children



Fig. 232.—Diagram of discission with two needles. The needles have been inserted, but not yet rotated.

compared with safety to the eye, I prefer to avoid curette evacuation in these cases if possible. Simple needling causes little reaction, and septic infection is very rare if a subconjunctival puncture is made, as recommended above. Excessive swelling of the lens and curette evacuation cause much more disturbance, and the risks of sepsis are greater, for the swollen lens substance is an excellent culture medium.

Discission of Secondary Cataract (after-cataract) is performed in exactly the same manner as discission of the soft lens if the after-cataract is not too dense to be divided by a single needle. Some surgeons advise needling every case of senile extraction as soon as the eye has quieted down, *i.e.*, in a fortnight or three weeks: the capsule is then soft and easily divided. If there has been iritis or iridocyclitis needling is contraindicated until all signs of inflammation have passed off; hence a careful search for "k.p." must be made in every case.

If the capsule is dense and thickened a Ziegler's knife may be used (Fig. 231), or two needles may be employed. By the latter method no undue strain is thrown upon the ciliary body, and the membrane is prevented from tearing away from the ciliary body instead of being torn in the centre. The same instruments are required together with an extra needle. The needles may be "stop" needles, though this should not be necessary. The eye is fixed by an assistant. The needles are introduced with the flat surfaces upwards through the cornea at opposite sides of it in the horizontal meridian, 3 or 4 mm. internal to the apparent margin (Fig. 232). They are then rotated through a right angle, so that the cutting edges are applied to the membrane. They are passed through the centre of the membrane close to each other. The handles are then simultaneously raised, so that each needle rotates around the spot on the shaft which is engaged in the cornea. In this manner the points are drawn apart, and the membrane is cut. A further attempt may be made to tear it in the direction at right angles. The needles are rotated so that the flat surfaces are again upwards, and are withdrawn quickly so that aqueous may not be lost. Atropin is instilled, and the eye is bandaged. There is seldom much reaction, unless the previous operation has been faulty; in such cases the additional irritation may precipitate an attack of sympathetic ophthalmia.

When the membrane is very dense it is often best to divide it with scissors, as in iridotomy (*vide* p. 437); or the very ingenious canula-scissors may be employed. They were invented many years ago, but fell into disrepute owing to the loss of eyes from sepsis. They should be kept in alcohol when not in use. A small peripheral puncture with a broad needle is first made, and through this the canula-scissors are introduced.

It may be mentioned that needling operations are by no means so simple as they appear in the hands of an experienced operator. Every movement of the needle must be made round the spot where the shaft penetrates the globe.

Curette Evacuation or Linear Extraction is the operation whereby after discission, whether accidental (traumatic cataract) or intentional, the softened lens matter is let out of the anterior chamber.

Instruments required: speculum (Fig. 216), fixation forceps, bent keratome, curette (Fig. 234), iris repositor. Local

anæsthesia suffices, except in the case of children or unruly patients.

The pupil must be fully dilated with atropin. The position of the section is of little importance: some surgeons place it above, where it is under the upper lid, others below. The surgeon stands accordingly either above or at the side of the patient.

The conjunctival sac having been douched and the speculum inserted, the eye is fixed with fixation forceps. The keratome is passed through the cornea, 1 mm. internal to the apparent margin (cf. wound in Fig. 215), with the blade parallel to the plane of the iris. It is pushed on until the incision is about 5 mm. long. The point may be dipped so as to pass into the lens without disadvantage, and the incision may be increased as the keratome is being slowly withdrawn by extending each angle, using the two edges of the keratome like knives. The tip of the curette is then gently insinuated just within the edges of the wound, not quite into the anterior chamber. Slight pressure is exerted upon the peripheral lip, and the soft lens matter travels along the groove of the curette. No attempt should be made to remove the whole of the lens matter, on account of danger of rupture of the suspensory ligament and escape of vitreous. The remnants of lens matter will be absorbed. There is no probability of the iris prolapsing or becoming incarcerated in the wound if it is properly dilated, but the repositor should be at hand in case this accident occur. Sterilised atropin ointment is inserted, and both eyes are bandaged.

Usually the ciliary irritation is greatly benefited by the operation. The greatest care must be exercised as to surgical cleanliness, for most of the accidents from infection in treating soft cataract occur, not from needling, but from curette evacuation. Anterior synechia of iris or capsule, iridocyclitis, or sympathetic ophthalmia may occur, but are happily rare.

Some surgeons extract soft cataracts by the linear method without previous discission. This mode of treatment is not advocated.

Extraction of senile cataract and of lenses containing a hard nucleus may be performed with or without iridectomy. The latter method should seldom be adopted. The iridectomy is performed by some surgeons six weeks or more before extraction. I advise this method only in cases of immature cataract (*vide* p. 294), and in cases where it is necessary to operate, though there is some doubt as to the sterility of the con-

junctiva, &c., in spite of thorough preliminary treatment. Infection is likely to be more under control after iridectomy than after extraction, and the behaviour of the eye in the first operation is a guide to its probable behaviour after extraction.

In every case of cataract, before operating the condition of the patient's conjunctiva and lacrymal apparatus is

FIG. 233. — Graefe cataract knife. A slightly broader knife is used in cataract extraction than in glaucoma iridectomy; it has the advantage of being less resilient. The knives used for glaucoma iridectomy are generally re-ground cataract knives.



FIG. 234. — Cystotome (Moorfields pattern) and curette. (These should *not* be at the two ends of the same instrument.)

thoroughly investigated. If there is any regurgitation of tears, mucus, or pus on pressure over the lacrymal sac, the condition must be cured (*vide* p. 595) before operation. If there is any conjunctivitis, it must be treated and cured, as shown by the absence of pathogenic organisms on bacteriological examination. Some of these cases are very troublesome (*vide supra*); in case of doubt, the patency of the nasal duct should be demonstrated by syringing.

The usual preparation of the patient—bath, purgative, and so on—is made as for any major operation. The lids are cleansed with ether soap on the previous night.

Cataract Extraction with Iridectomy ("*Combined Extraction*").—Local anæsthesia suffices, unless the patient has lack of self-control. Sterile 2 per cent. solution of cocain is instilled into the eye three times at intervals of five minutes, the eyelids being kept closed between the applications of the drops. At the last instillation a drop is also instilled into the other eye. This is necessary, since, if fluid happens to drop into this eye during the operation, the patient is liable to squeeze the lids together reflexly, unless the eye is cocained: moreover, it helps the patient to keep both eyes open, diminishing the risk of winking. A drop of adrenalin solution (1 in 1,000) may with advantage be put into the eye to be operated upon. The surgeon stands over the patient, making the section with his right hand on the right eye, and with his left hand on the left eye. (Some surgeons stand below and at the side for the left eye, and cut away from themselves, using the right hand.)

Fig. 235.—Maunoir's angular iris scissors.

Instruments required: speculum (Fig. 216), two pairs of fixation forceps, Graefe cataract knife (Fig. 233), bent iris forceps, de Wecker's scissors, cystotome (Fig. 234), curette (Fig. 234), iris repositor [probe-pointed bent scissors (Maunoir's) (Fig. 235), lens spoon (Fig. 236)]. The instruments in brackets are not required for uncomplicated extraction, but must always be ready.

The following description applies to operation upon the right eye.

The conjunctival sac having been douched and the speculum inserted, the patient is told to look towards his feet and a little outwards, and the eye is fixed with fixation forceps applied just outside the limbus directly downwards or preferably down and in below the horizontal meridian of the cornea. The point of the Graefe knife is inserted in the apparent corneo scleral margin at a point corresponding with ten or twelve minutes to twelve on the dial of a clock (Fig. 237). It is passed across the anterior chamber to the corresponding spot on the opposite side, ten or twelve minutes past twelve respectively. The knife is made to cut out in exactly the same manner as in iridectomy for glaucoma (*q.v.*), but following the corneo-scleral margin. A conjunctival flap should be made as in that operation, but it will be narrower at the sides, since the section is farther forwards. It may be made slightly broader above by carrying the conjunctival incision a little upwards after the corneo-sclera has been divided before turning the knife to cut directly forwards. The conjunctival flap is then turned down over the cornea by means of the back of the knife.



Fig. 236. Critchett's lens spoon.

The fixation forceps are then removed. If the patient is unsteady, they are handed over to an assistant, who applies them below the cornea and maintains fixation. If the patient is submissive, he is simply told to keep looking towards his feet all the time. He is warned that he may feel a little pain during the next stage, but that he must keep still, or the operation may be unsuccessful.

The iris forceps are taken in the left hand and the de Wecker's scissors in the right (irrespective of the eye which is being operated upon). The closed points of the forceps are inserted at the centre of the wound and passed straight downwards to just above the pupillary margin of the iris. The blades are opened slightly and the iris seized and drawn out of the wound. The piece of iris grasped is cut off with one snip of the de Wecker's scissors, the blades of which are held radially to the iris, *i.e.*, with the points directed upwards (Fig. 218).

The iris repositor is now taken and the iris replaced as in the

operation for iridectomy (*q.v.*). Particular care is taken to free it from the angles of the wound, usually an easy task at this stage owing to the support afforded by the lens and the smoothness of the surface of the capsule.

The patient still looking towards his feet, the cystotome is introduced with the cutting edge directed towards the left and slightly upwards. It is passed on until the point is near the lowest part of the pupillary margin. The cutting edge is then directed backwards, the handle is slightly raised, and the lens capsule is incised vertically as the instrument is slowly withdrawn.

The cystotome is then taken in the left hand, and the curette in the right. The back of the curette is placed horizontally upon the lower part of the cornea (from about twenty



FIG. 237.—Diagram of wound in extraction of senile cataract.



FIG. 238.—Diagram of coloboma in cataract extraction with iridectomy.

minutes to twelve to about twenty minutes past twelve). Gentle but firm pressure is made upon the cornea in a direction backwards and slightly upwards. This causes the nucleus of the lens to be tilted so that the upper edge appears presenting in the wound. The lens nucleus is coaxed out of the wound by repeating the pressure with the curette, but more and more in an upward direction. Meanwhile the lens may be gently helped out by the cystotome in the left hand. As soon as the diameter of the nucleus has passed through the wound, pressure upon the cornea is immediately relaxed. The nucleus is directed down over the cornea into the curette.

If there is much clear soft lens matter, this is coaxed out of the wound by gently repeating the stroking movements with the curette on the cornea. Too much pressure must not be exerted. Any soft matter protruding from the wound is removed with the curette.

The iris repositor is now again used to free the iris from the

angles of the wound. If the iris has been carefully replaced after the iridectomy, as recommended above, little trouble will probably be experienced at this stage. If this step has been omitted it is likely that the nucleus may have jammed the iris into the angles of the wound during its expulsion and there may be great difficulty in freeing it. In addition, the repositor is swept over the whole line of the wound, so as to push back into the anterior chamber any tags of capsule which may be presenting. These are so transparent as to be invisible, and it is extremely important that they should not become incarcerated in the wound.

The conjunctival flap is then turned back into its natural position by the repositor, care being taken that it is not doubled up.

The removal of the speculum and closure of the lids are carried out as in iridectomy (*q.v.*). Both eyes are bandaged.



Fig. 239.—Diagram of Moorfields' cataract bandage. The semicircular opening is for the bridge of the nose. The upper and lower tapes are passed above and below the ears. The long single tapes pass behind the head and are brought forward and tied over a pad of wool placed at the upper part of the bandage over the forehead.

and the patient is put to bed without altering his recumbent posture. It is a good plan to operate with the patient upon a suitable bed, so that it is unnecessary to move him.

The dressings should consist of two large round pads of sterilised gauze covered by similar pads of sterilised cotton wool. They are kept in position firmly, but without undue pressure, by a cataract bandage. Of the many types of the latter the Moorfields' pattern (Fig. 239) is probably the best. With aged or unruly patients a wire or "cartella" shield (Fig. 240) should be placed over the dressing.

Remarks upon the Operation.—The size of the section depends upon the probable size of the nucleus of the lens. It must be remembered that the cortex is soft and broken up: the width of the incision must be slightly larger than the diameter of the nucleus. In black cataract the whole lens is sclerosed, so that a very large section must be made: in these cases it should involve half the circumference of the cornea. More harm is done by bruising the edges of the wound than by

having a wound which is unnecessarily large. If the nucleus does not come forward through the wound with moderate pressure of the curette it is probably because the section is too small. The wound should then be enlarged with the probe-pointed bent scissors (Fig. 235). The probe-point of one blade is inserted between the iris and cornea at one angle of the wound, which is then extended by a single snip. The same manœuvre may be repeated at the other angle.

In making the section the aqueous may escape prematurely, so that the iris floats up in front of the knife. If this happens the section must be completed in the usual manner, though the iris is wounded, either a hole or a complete coloboma being cut out of it. Cutting the iris causes pain and is likely to make the patient flinch, screw up the eye, or completely lose self-control. Moreover, the coloboma is generally irregular. The accident is usually due to hesitation in pushing the Graefe knife steadily forward or to pressure —often unconscious—exerted on the eye by the fixation forceps. It is least likely to occur if the passage of the knife through the anterior chamber and the commencement of the section are all part of a single steady forward sweep of the blade, the handle of the knife being depressed directly the counter-puncture is made. In this manner the broad part of the blade is brought over the iris as quickly as possible. There should, however, be no haste, every movement being made with deliberation but not too slowly.



Fig. 240.—"Cartella" shield for right eye.

Old people sometimes have very rigid sclerotics. In these cases the cornea collapses and becomes saucer-shaped. This is of no consequence and requires no special treatment.

Sometimes a bubble of air enters the anterior chamber. This also is of no consequence, as it soon becomes absorbed.

The iris is more easily cut off by holding the de Wecker's scissors at right angles to the direction of the forceps, *i.e.*, horizontally (Fig. 219). This makes a larger coloboma, which is unnecessary and has some disadvantages.

Hæmorrhage may occur into the anterior chamber. The blood is derived from the conjunctival flap or from an hyperæmic iris. An attempt may be made to drive it out of the

wound by stroking the cornea upwards with a spatula, but too much time should not be devoted to this endeavour. If it fails the capsulotomy must be done with great care to avoid catching the iris.

The capsule is divided in all sorts of different directions by different operators. Probably the most satisfactory is a circumferential incision just inside and parallel with the wound. It is less easy to do than the method recommended. The other methods have no advantage. Some surgeons remove the anterior capsule by capsule forceps (Fig. 127), which has very decided advantages, but has also its own special dangers.

The most serious accident which may occur during extraction is prolapse of vitreous at an early stage. It may be due to inherent weakness of the suspensory ligament, which gives way while the section is being made or the iridectomy done. This cause is most likely to be present in complicated cataracts. In such cases it is a good plan to discard a speculum, an assistant holding the lids apart and raised off the eye by means of a pair of Desmarres' or similar lid-retractors. More commonly loss of vitreous is due to undue pressure on the eye by the fixation forceps. In concentrating all their attention on the section beginners often allow the left hand to dig the fixation forceps into the globe. Special attention must therefore be devoted to the avoidance of this mistake, which has also the lesser disadvantage of forcing out the aqueous and allowing the iris to float up in front of the knife. Should this occur the iris will be cut and pain caused, often leading the patient to flinch. The section must be completed, however, in the usual manner. Escape of vitreous may also be due to pressure with the curette in the attempt to expel the lens. The necessity of such great pressure is probably owing to the wound being too small or to the capsulotomy having been inefficiently performed. The former contingency has already been dealt with. The latter is overcome by more careful repetition of the capsulotomy. If pressure with the curette causes the vitreous to appear without any sign of the engagement of the edge of the lens in the wound resort must be made at once to scoop extraction. Hence the importance of having the spoon always in readiness in every case of extraction. The spoon is passed directly backwards into the vitreous so as to make certain that it passes behind the lens. It is then rotated forwards so that the lens is caught between the spoon and the back of the cornea. The lens is kept pressed up against the cornea and is removed by a rapid further rotation of

the spoon. Some vitreous is certain to be lost, but it is imperative that the lens should be delivered, or the eye will almost inevitably be lost. The eye is dressed at once in these cases without any endeavour being made to replace the iris, as any such manœuvre is likely to lead to futher loss of vitreous. In spontaneous rupture of the suspensory ligament and in other cases badly managed the lens may sink back into the vitreous. In such cases it is usually futile to attempt to remove it. The eye should be dressed at once, and if the lens floats up into the pupillary area at a later date a further attempt may be made to remove it.

Prolapse of the vitreous after delivery of the lens is less serious, though it increases the tendency to cyclitis, with opacities in the media, and may be followed by detachment of the retina. If much vitreous is lost the iris is always gradually drawn upwards, so that in course of weeks or months the pupil is much displaced and the lower part of the iris stretched (Fig. 220). This condition may also occur from incarceration of the pillars of the coloboma in the wound. If may be necessary to do an iridotomy or some such operation to make an artificial pupil so that vision may be restored.

When the cataract is immature some of the soft lens cortex remains in the eye, so that the pupil is not black but contains greyish masses. Much of the retained lens substance can be removed by stroking the cornea upwards with the spatula, repeating the movements used for delivering the lens, but with less pressure. It is usually impossible to remove it entirely in this manner. If it is left in the eye it gradually becomes absorbed, but it has the disadvantages of tending to irritate the eye and cause slight iridocyclitis, and of leading to the formation of a denser after-cataract. Some surgeons irrigate the anterior chamber with normal saline solution. This method gets rid of the lens substance, but in my experience is very liable to set up a mild iridocyclitis, in spite of the strictest antiseptic precautions. Irrigation may be performed with an undine to the nozzle of which an india-rubber tube is attached, having a flattened canula at the other end. The tip of the canula should be introduced just inside the lips of the wound, and the undine should not be held too high, only a gentle stream of fluid being used. I use the irrigator only in exceptional cases.

After-treatment.—There is usually some aching and smarting in the eye as soon as the effects of the cocain wear off. It lasts for four or five hours: hence it is best to operate in the

morning so that the patient may have a good night's rest. If the pain interferes with sleep a dose of aspirin will usually relieve it, or a mild bromide draught may be given.

The patient lies quietly upon his back. He is directed to avoid all straining. A sneeze may be inhibited by pressure with the finger on the upper lip close to the septum of the nose. *All* patients should have their hands loosely tied to the bed at night, so that they are unable to touch the eyes. Many eyes are lost from neglect of this precaution, for patients often knock or rub their eyes when they are half asleep.

The food must be fluid during the first few days; no aperient is given for three or four days.

On the day following the operation the bandage is removed, the lids are bathed with warm boric lotion, gently separated, and a drop of sterile 1 per cent. atropin solution instilled. The wound may be inspected, but should not be disturbed more than is absolutely necessary.

On the second day it is examined more thoroughly. The cornea should be bright and the pupil round and well dilated. Faint greyness in the cornea above (striate opacity, *vide* p. 234) need cause no alarm. Another drop of atropin is instilled. If the pupil is not well dilated on the third day there is probably some trace of iritis, and the atropin should be instilled more frequently, and it may be advisable to resort to hot bathings. In most cases there is no iritis, and after a transient ciliary injection the eye quiets down, so that it is almost free from injection in a week or ten days.

On the fourth or fifth day the unoperated eye may be left unbandaged. If both eyes are kept bandaged too long old people often become delirious. On the slightest sign of wandering in speech the unoperated eye should be uncovered at once. If this eye is blind or has very defective vision the dressings should be taken off the operated eye, and dark protective goggles worn during the daytime.

It is wise to keep healthy patients in bed for a week. Most cataract patients, however, are old, and extremely liable to hypostatic congestion of the lungs. Such patients should be propped up in bed on the third or fourth day, and allowed to sit up out of bed soon after. A light dressing is kept on for ten days or a fortnight; afterwards smoked glasses are worn until cataract glasses can be ordered, *i.e.*, in about six weeks. It is very important that smoked glasses should be worn continually, otherwise the patient is quite likely to develop red vision or erythropsia (*q.v.*).

Cataract Extraction without Iridectomy ("*Simple Extraction*"). Many surgeons advocate extraction of cataract without iridectomy in suitable cases, but opinion differs much as to the criteria of suitability for this operation. It is performed in the same manner as the combined operation, except that the iridectomy is omitted. The lens is delivered through the pupil.

Comparing the two operations, it may be succinctly stated that the chief advantages of *simple extraction* are: Simplicity of the operation, including especially minimum of mutilation, of instruments required, and of instruments introduced into the interior of the eye; optical advantages of a round pupil—minimal dazzling, best visual acuity, and best field; cosmetic advantage of a round pupil; ease of reposition of the iris; minimal danger of incarceration of capsule in the wound; infrequency of prolapse of vitreous; greater protection of deeper parts of the eye from infection; absence of pain and bleeding from cutting the iris. Of these it may be remarked that the optical advantages are not overwhelmingly manifest. The cosmetic effect is rarely of sufficient importance to outweigh the risks, though occasionally it is a justifiable argument, as in the case of an unsightly unilateral cataract in a young woman, or in a man in whom it forms a bar to obtaining employment. It is doubtful whether prolapse of vitreous is more frequent in combined than in simple extraction. Bleeding from the iris is only very exceptionally of any importance; when there is much bleeding in cataract extraction the blood is derived from the conjunctival flap or from a too peripheral section. The chief disadvantages of simple extraction are: risks of prolapse of the iris; less efficient treatment of the anterior capsule; greater difficulty in removal of soft lens matter; greater danger of ring synechia and secondary glaucoma if iritis should occur.

The chief advantages of *combined extraction* are: greater ease in expression of the nucleus; greater ease in removal of soft lens matter; increased facility in dealing with the anterior capsule, and therefore diminished necessity for subsequent discission; diminished risk of prolapse of the iris; diminished risk of secondary glaucoma. The chief disadvantages are: greater complexity of the operation, including especially need of more instruments and of the introduction of more instruments into the eye, and greater duration of the operation; optical and cosmetic diadvantages; greater danger of incarceration of iris and capsule in the wound; greater danger

of post-operative glaucoma; pain and bleeding from the iris.

The frequency of prolapse of the iris and the dangers resulting therefrom are such serious risks that no surgeon should perform simple extraction until he has obtained exhaustive experience of the combined operation. If there is any difficulty in replacing the iris after a simple extraction, or if the pupil, when the iris is reposed, is not quite circular, an iridectomy should be done at once. Prolapse is liable to occur within the first twenty-four hours of the operation. If it is not treated by an immediate iridectomy a very serious condition will result. The incarcerated iris will fail to withstand the intraocular pressure and a "cystoid cicatrix" will be formed. Visual acuity will be diminished by excessive astigmatism and the eye exposed to grave danger from iridocyclitis, and even panophthalmitis or sympathetic ophthalmia.

Many of the disadvantages of both operations, including the danger of prolapse of iris, are obviated by a modification in which after simple extraction has been performed a small button-hole is made in the periphery of the iris (*peripheral iridectomy*). After the lens has been extracted, as in simple extraction, the periphery of the iris is gently drawn out of the wound by iris forceps and the smallest possible fold is snipped off with de Wecker's scissors. The aperture in the iris is peripheral, so that it is almost entirely hidden by the sclerotic, and in any case is completely covered by the upper lid, while at the same time sufficient drainage from the posterior into the anterior chamber is provided for. Prolapse of iris is less likely to occur than in simple extraction, and it is quite unlikely that the capsule will be left entangled in the wound, a decided drawback to extraction with the ordinary form of iridectomy.

The chief complications arising after cataract extraction are striate "keratitis," incarceration of the iris in the angles of the wound, prolapse of the iris, iritis, iridocyclitis, sympathetic ophthalmia, secondary glaucoma, intraocular hæmorrhage, infection of the wound, panophthalmitis, &c.

Prolapse of Iris is most apt to occur after simple extraction, but may affect either pillar of the coloboma in combined extraction. It usually occurs in the first day or two, but may result later from injury to the eye by rubbing or knocking it, straining, coughing, &c. It must be treated *at once* by excision of the prolapse. As the iris is irritable the operation is painful, and a general anæsthetic is generally necessary or advisable. The wound is re-opened by insinuating the tip of an iris repositor

under the conjunctival flap and gently uncovering the prolapse. The flap is turned down over the cornea, the iris pulled out with iris forceps and snipped off with de Wecker's scissors. The iris is then replaced with a clean repositor and the conjunctival flap brought back into position.

Delayed Re-formation of the Anterior Chamber may be due to a jagged section, over-riding of the lower lip of the wound, or to no apparent cause. It is much less common in cataract extraction than after glaucoma iridectomy. In these cases the bandage should be very lightly applied or discarded, dark protective goggles being worn in the daytime, and a light bandage with wire or cartella shield at night.

Delayed Healing of the Wound is more likely to occur with a purely corneal section, such as some surgeons prefer. The patient should be kept in bed until it is firmly healed, unless this is specifically contraindicated (*vide* p. 454). Delay in healing is, however, generally due to incarceration of iris or capsule in the wound. If this amounts to an actual prolapse it must be operated upon (*vide supra*), but it may be very insidious. In either case the result may be the formation of a *cystoid cicatrix*, part or the whole of the scar slowly and gradually becoming more and more ectatic. The eye should be carefully examined to see if the pupil or either pillar of the coloboma is drawn up, or if capsule can be seen in the wound. A fully developed cystoid scar should be left alone, though it causes much astigmatism and is liable to give rise to secondary infection, or iridocyclitis and even sympathetic ophthalmia.

Expulsive Hæmorrhage is fortunately rare. It occurs during or soon after operation in old people with arteriosclerosis or some diathesis, such as diabetes. There is sudden severe pain, and on removal of the dressings the wound is found to be gaping and filled with blood clot, vitreous, &c. The eye is always lost and should be excised. This may be necessary in order to stop the bleeding, the socket being then packed and firmly bandaged.

Septic Infection may occur in spite of all precautions, especially in diabetic patients. It is most commonly due to the pneumococcus, but may be caused by the streptococcus, staphylococcus aureus or even albus, and many other organisms. It usually occurs from the twelfth to thirty-sixth hour after operation. There is severe aching pain, due to the accompanying acute iritis. On removing the dressings the upper lid is œdematous. When the lids are separated tears gush out and there is muco-pus in the conjunctival sac. The lids should be separated gently, if necessary with retractors. The cornea is then seen to be dull and hazy, especially in the upper part, the lip of the wound being yellow. Almost invariably the infection spreads rapidly. Intense iritis

is set up, the pupil and coloboma become filled with exudate and an hypopyon appears. Finally the vitreous becomes infected and panophthalmitis leads to the destruction of the eye.

Treatment is seldom of any avail, but must be applied quickly and energetically. The anterior chamber should be irrigated (*vide* p. 453) with hydrogen peroxide solution. The lips of the wound should be cauterised with the galvano-cautery. Sub-conjunctival injections of perchloride (1 in 2,000) or oxycyanide (1 in 5,000) of mercury may be given, but add much to the pain and are seldom of use. A staphylococcic or polyvalent vaccine should be given and an autogenous vaccine prepared and administered as soon as possible.

Iritis in mild degree probably occurs in all cases of cataract extraction. In more pronounced form it is specially associated with retained lens matter (*vide* p. 453) and diathetic states, such as diabetes, rheumatism, gout, &c. The worst cases occur with acute septic infection. Intermediate in severity are cases of plastic iritis due to infection by less virulent organisms or in patients with greater resistance to bacterial invasion. More insidious than any are cases of continued irritability of the eye with mild iritis. In these, spots of "k.p." are found upon the back of the cornea, so that there is also cyclitis. Both in these cases and in the cases of plastic iritis there is grave danger that the condition is really sympathetic ophthalmia. Hence it is very essential in all cases of cataract extraction to inspect the cornea most carefully with oblique illumination and the loupe and to do so frequently, especially if there is an unusual degree of flushing and lacrymation on exposure to light. The other eye must also be carefully watched. If there is "k.p." no needling operation must be undertaken until the eye has quieted down and remained quiet for many weeks.

Detachment of the Choroid.—*Vide* p. 327.

Secondary Glaucoma may set in after cataract extraction. It is probably usually due to peripheral anterior synechia and incarceration of capsule in the wound (*vide* p. 264). Sometimes it is due to the anterior chamber being lined with epithelium. In these cases there has been delay in healing and the conjunctival epithelium has grown down into the anterior chamber and spread over the surface of the iris, lens capsule and cornea. These cases are practically hopeless and cannot be diagnosed clinically with certainty. Sometimes secondary glaucoma follows needling of the after-cataract. It is usually then attributed to vitreous extending into the anterior chamber and interfering with filtration. It is doubtful if this is the true explanation. These cases usually do badly. As regards treatment, where there are definite adhesions of capsule an attempt may be made to divide them.

These cases afford the best prognosis. In the more obscure cases the eye should be trephined.

Excision of the Eyeball.

A general anæsthetic is required. The operation can be performed almost painlessly under novocain or eucain, and this method is sometimes advisable in old people with diseased arteries, or in patients with heart disease. Two c.c. of the solution (*vide* p. 629) are injected deep into the orbit on the nasal and temporal sides of the globe below the horizontal meridian.

Instruments required: speculum, two pairs of fixation forceps, tenotomy scissors (Fig. 241), strabismus hook (Fig. 242), blunt-pointed scissors — straight or curved on the flat (Fig. 243).

The surgeon stands above the patient.

The conjunctival sac having been douched and the speculum inserted, the surgeon seizes the conjunctiva just outside the limbus at the highest part of the cornea. The conjunctiva is incised here with the tenotomy scissors. The point of one blade of the scissors is passed under the conjunctiva and pushed on as far as possible round the cornea. By carrying the point out under the looser bulbar conjunctiva it may be taken a third of the distance round the circumference; the edge is then brought close up to the limbus before the conjunctiva is divided. Still fixing the eye in the same position the manœuvre is repeated on the other side of the cornea. Finally, the portion below the cornea is divided. The conjunctiva should be divided com-



Fig. 241.—Tenotomy scissors.

pletely round the cornea, and close to it in three or four cuts.

The peripheral edge of the cut conjunctiva is then taken up by the forceps, and the bulbar conjunctiva is separated from the globe as far back as the equator in all directions by a series of small snips, the blades of the scissors being kept flat in close contact with the eyeball. In this manner the capsule of Tenon is simultaneously opened.

The tenotomy hook is then taken in the left hand, the scissors being retained in the right. The recti muscles are taken up one by one and divided close to the globe. It is well to begin with the superior rectus, since it is the most difficult to get at, especially if the other recti have been previously divided. The obliques are found by passing the hook farther back and carrying it round close to the globe.

Fig. 242.—Strabismus hook, flat Moorfields' pattern.

The speculum is then taken and held widely open and pressed back into the orbit. If the muscles have been properly divided the globe springs forwards between the blades of the speculum. The other pair of scissors is now taken in the right hand. The globe is seized with the fingers of the left hand. The points of the closed scissors are passed into the orbit—to the outer side of the eye on the right side, to the inner on the left. The optic nerve is felt for with the closed scissors: it is easily recognised. The scissors are withdrawn a short distance, opened, and the blades pushed down, one on each side of the nerve, which is then divided. The sensation of dividing the nerve is unmistakable. The eyeball can then be freely drawn forwards. There are probably some remnants of the obliques still attached to the globe. These are divided close to the eye. The bleeding is profuse for a moment, but may be controlled by a little pressure with a pad of cotton wool. The edges of the conjunctiva are then pulled together with the fixation forceps, the lids are closed, and the dressing is applied. The latter should consist of a small spherical pad of cotton wool, then a round flat pad of sterilised or cyanide gauze, then a thick round pad of sterilised wool. The bandage

is applied with firm pressure. The patient is kept in bed for one or two days.

If the globe is perforated and collapsed excision is more difficult. The eyeball should be held forward with large strong vulsellum forceps. In children also excision is difficult. The orbit is small in proportion to the size of the globe. Here the manœuvre with the speculum to dislocate the ball forwards is often unsuccessful, and strong pressure may burst the eye. It is best levered out with the points of the excision scissors.



Fig. 243.—Excision scissors.

It is very easy to cut the sclera instead of the nerve, especially with curved excision scissors. I always prefer straight scissors; indeed, straight instruments should always be used in preference to curved whenever it is possible, because it is much easier to judge the position of the point. Straight scissors are particularly indicated when the nerve has to be cut long, as in excision for glioma retinæ and sarcoma of the choroid. Special precautions must be taken in excision for panophthalmitis (*vide* p. 425).

Evisceration of the eyeball is recommended only in some cases of panophthalmitis (*q.v.*). Some surgeons use it in anterior staphyloma and other conditions. Although the stump affords a good foundation for an artificial eye, the operation is not to be recommended in these cases: it has been followed by sympathetic ophthalmia.

Evisceration with insertion of an artificial vitreous—*Mules's operation*—is even more objectionable on theoretical grounds,

but is still performed by some surgeons. In this operation a glass sphere is inserted inside the sclerotic, which is stitched over it. The globe must be much smaller than the normal vitreous chamber. There is very severe chemosis and pain after the operation, and the glass globe may be extruded. Sympathetic ophthalmia has followed this procedure.

Excision, with the introduction of a glass globe in Tenon's capsule, is much less objectionable than Mules's operation. In carefully selected cases it is a good method, and affords admirable support to an artificial eye. The eye is much more movable than after simple excision, so that the simulation of a real eye is more exact. Care must be taken during the excision to keep close to the eyeball and to injure Tenon's capsule as little as possible. The recti muscles are drawn together over the small glass globe by a buried purse-string suture. The conjunctiva is sutured by a superficial purse-string suture. There is some inflammatory reaction, with chemosis, but much less than after Mules's operation, and little or no pain. If the operation has not been well performed, the globe may slip into the orbit, becoming loose, usually beneath the lower lid. If this occurs it must be removed and the case treated as an ordinary excision.

An *artificial eye* should not be worn less than six weeks after excision. A small eye is first worn for an hour or two a day until the conjunctiva becomes used to the foreign body. Eight or nine weeks after the operation a full-sized eye may be worn. It is taken out at night, carefully washed, and kept in water. If it becomes in the slightest degree rough at the edges it must be discarded at once. This usually occurs from chemical action of the secretion upon the enamel in about a year, so that in any case a new eye should be obtained every year. Various types of artificial eye are made. Snellen's "reform" eye gives a better cosmetic effect after the ordinary excision than the old shell, which may be used when there is good support, as after insertion of a globe in Tenon's capsule. The methods of insertion and removal of the artificial eye should be learnt by every surgeon by seeing it done.

Contracted Socket is the result of injury, faulty excision, cellulitis in the orbital tissues, or the continued wearing of a rough artificial eye. The first three causes lead to the formation of dense cicatricial bands across the socket, rendering the wearing of a prothesis impossible. The last cause usually results in obliteration of the lower fornix, so that the eye cannot be kept in place.

Contracted sockets are difficult to remedy. It is easy to divide the bands and make a new groove to hold the eye in position, but unless the wounds become covered with epithelium the edges heal together and no improvement is produced. It is therefore necessary to perform some plastic operation, whereby the wounds made are covered with an epithelial graft or skin flap. Grafting the wound with a Tiersch graft, mucous membrane from the mouth of the frog, or mucous membrane from the lip may suffice. Grafts may be kept in place by gutta percha, lead foil, or a glass shell, but this method usually fails. A better method is to divide the outer canthus, fix the everted lids to the brow and cheek by sutures, and cover the exposed surfaces with grafts. The lids are replaced when the grafts have taken satisfactorily.

The greatest difficulty is to restore the lower fornix. Maxwell's operation is the best for this purpose if the lower lid is uninjured. It is simple, but difficult to describe. An elliptical area of skin is marked out on the lower lid. The upper incision is carried through into the socket, in the position of the new fornix. The flap is about 8 mm. broad in the centre. It is dissected up at the edges all round, but a central elliptical pedicle is carefully retained. The flap is tucked through into the orbit; the upper edge is sutured to the posterior lip of the conjunctival wound, and the lower edge to the anterior lip. The gap in the skin is then closed with sutures, and a glass shell is inserted in the socket. The pedicle ensures the vitality of the flap and also keeps the new fornix depressed to the level of the orbital margin. If much ectropion results the scar can be re-opened at a later date and a Wolfe graft of sufficient size inserted. Wolfe grafts must not be sutured in position, but simply laid on the surface and kept in position by a network of stitches passing from one edge to the other of the wound. Further, they must be kept dry, and should therefore be powdered over with dry boracic acid.

Weeks inserts a large Wolfe graft into the orbit, suturing it at the apex of the fold to the periosteum of the lower orbital margin.

SECTION IV.

ERRORS OF REFRACTION AND ANOMALIES OF ACCOMMODATION.

CHAPTER XXIII.

RETINOSCOPY.

(Before reading this section the student should revise his knowledge of the optical conditions of the eye and the methods of testing visual acuity by again reading Chapters III., IV., VII., and IX.)

It has been already pointed out that the condition of the refraction of an eye can be estimated in various manners. The systematic examination of the visual acuity will in most cases *indicate* the absence or the nature of any error of refraction. The examination with the mirror at a distance of 1 metre also indicates the refractive condition by the visibility of retinal vessels and the direction of parallactic displacement; as will be shown below, this method may be made to give very accurate estimation of the exact refraction. The indirect method also indicates the refractive condition by the apparent change in size of the disc when the large lens is moved away from the eye. By the direct method the condition of the refraction can be accurately measured if the surgeon has acquired the ability completely to relax his accommodation.

Retinoscopy, or, more correctly, *skiascopy* or *the shadow test*, is the most accurate means at our disposal of estimating the condition of the refraction objectively. It depends upon the fact, first pointed out by Bowman, that when light is reflected from a mirror into the eye the direction in which the light travels across the pupil varies with the condition of refraction of the eye. If the light is thrown into a myopic eye from a concave mirror at a distance of 1 metre the light,

or what is easier to observe, the shadow, moves across the pupil when the mirror is slowly tilted in the same direction as that in which the mirror is moved (Fig. 244). If a plane mirror is used, the other conditions remaining the same, the shadow will be seen to move in the opposite direction to the movement of the mirror. If the eye is hypermetropic the direction in which the shadow moves is the opposite of that with the myopic eye. If the eye is emmetropic or has only a very low degree of myopia no shadow will be visible; the pupil will be either completely illuminated or completely dark.

The light seen in the pupil is the blurred image of the illuminated area of the fundus as seen by the observer when



FIG. 244.—Diagram of retinoscopy with a concave mirror. O_1 the observed eye; O_2, the observer's eye. The image of the source of light is formed at l_1 (the immediate source of light) by the mirror. If O_1 is hypermetropic a virtual image of l_1 is formed on the line l_1 n, passing through the nodal point n, as at h_1. If O_1 is myopic a real inverted image is formed as at m_1. If the mirror is tilted downwards, as shown by the dotted line, l_1 moves to l_2, h_1 to h_2, and m_1 to m_2. This shows that the shadow moves in opposite directions in hypermetropia and myopia.

he accommodates for the observed pupil. The shadow is merely the image of the edge of the illuminated area.

Imagine a point of light in front of an eye, the pupil being dilated and the accommodation paralysed by atropin (Fig. 245). The divergent rays which enter the eye are made convergent by the refractive media, so that a circular area of the fundus, varying in size according to the refraction of the eye, is illuminated. If the point of light moves upwards, the light on the retina will move downwards.

Now consider the rays of light which are reflected from the illuminated area. In the hypermetropic eye they will be divergent, as if they came from a point behind the eye. This far point, corresponding with the illuminated area, will move in the same direction, *i.e.*, downwards. Now imagine an observer, placed in front of the eye to look towards a point of

light situated at the position of the far point, but to accommodate for the position of the observed pupil. He will see a circle of light with a blurred margin, not a point, because he is not accommodating accurately for the far point. When the illumination on the retina moves down the circle of light which the observer sees will appear to move down also (Fig. 245).



Fig. 245.—Showing the course of incident rays and field of illumination of the fundus in hypermetropia. l_1 forms a virtual image at λ_1, l_2 at λ_2. The field of illumination is determined by the pupil of O_1.

Again, consider the rays of light reflected from the illuminated area on the fundus of a highly myopic eye. They will be convergent and will cross at a real point in front of the eye. This far point, corresponding with the illuminated area, will move upwards when the illuminated area moves downwards. An observer placed in front of the eye and farther from it than the far point, if he looks towards the far point but accommo-



Fig. 246.—Showing the course of incident rays in myopia.

dates for the observed pupil, will see a circle of light with a blurred margin. When the illumination on the retina moves down the circle of light which the observer sees will move up, *i.e.*, in the opposite direction to the movement in the case of the hypermetropic eye (Fig. 246).

Now suppose that the observer's eye is one metre in front of the observed eye, and that the latter has 1 D of myopia. In this case the far point of the observed eye will be at the situation of the observer's eye, say at the level of his pupil (Fig. 247). In this case a very slight movement of the light on the observed fundus will throw the image at the far point

off the observer's eye altogether. In other words, the observed pupil will appear to be completely bright or completely dark.

If, again, the observed eye is emmetropic, its far point will be at infinity. We may regard it as being infinitely far behind the observed eye. Here again there will be scarcely any shadow, though in reality there is a very faint shadow moving in the same direction as for the hypermetropic eye.

The above is a simple explanation of the theory of retinoscopy. The question of the type of mirror is an entirely subsidiary one. It merely has to do with the direction of movement of the immediate source of light, *i.e.*, the point of light in front of the eye which has been considered above. The image of a real light behind the patient's head, formed by a concave mirror, is situated in front of the mirror. If the mirror is tilted up, the image moves up. The image of a real

FIG. 247.—Showing the course of the emergent rays at the point of reversal. So long as λ_1 is in the pupillary area of O_2, the pupil of O_1 appears uniformly illuminated, and there is no shadow. Directly λ_1 passes to λ_2 the whole of the light is cut off, so that the pupil of O_1 becomes completely dark.

light behind the patient's head, formed by a plane mirror, is situated as far behind the mirror as the light is in front of it. When the mirror is tilted up, the image moves down.

Hence under the actual conditions of retinoscopy with a plane mirror, when the mirror is tilted to the right the immediate source of light moves to the left, and—

(*a*) In the hypermetropic eye, the circle of light on the fundus moves to the right, and the shadow seen in the pupil moves to the right ;

(*b*) In the myopic eye (above — 1 D) the circle of light on the fundus moves to the right, and the shadow seen in the pupil moves to the left ;

(*c*) In the myopic eye of — 1 D there is no shadow :

(*d*) In emmetropia and myopia of less than — 1 D there is a very faint shadow moving to the right.

Stated as a mere guide to practice, with the plane mirror the shadow moves in the same direction as the mirror in hypermetropia and in the opposite direction in myopia above

one dioptre: in myopia of one dioptre there is no shadow and in emmetropia and myopia of less than one dioptre there is a very faint shadow moving in the same direction as the mirror.

In actual retinoscopy the whole of the image of the illuminated area of fundus cannot be seen at once: the shadow is part of the circumference. In high degrees of ametropia the shadow has a distinctly curved border, it is very dark, and it moves very slowly (Fig. 218). In low degrees of ametropia the border of the shadow looks straight: it is faint, and it moves very rapidly.

The movement of the shadow, being a purely optical phenomenon, is of course independent of the cause of the ametropia. Consequently in astigmatism, if one axis is hypermetropic and the other myopic (mixed astigmatism) the shadow moves in opposite directions in the two meridians. Often the periphery of the cornea is flatter than the centre; correction of the refraction of the central part, which is the more important, will then differ from that of the peripheral part. These variations produce very puzzling shadows in many cases.

Fig. 248.

Retinoscopy is applied to the estimation of refraction by placing correcting lenses in front of the eye and noticing the effect upon the shadow. When the shadow has completely disappeared we know that the eye has been made myopic 1 D if the surgeon is at one metre from the patient.

Retinoscopy is conducted in a dark room. The light is placed behind and above the patient's head. The surgeon sits at one metre from the patient. The patient wears a trial frame: the eye not under observation is covered by a screen. A mydriatic should be used by all but skilled observers, and is necessary to them in many cases. A plane mirror should be used. The patient looks at the observer's forehead.

The light is reflected into the eye, and the mirror is slowly tilted from one side to the other. The direction in which the shadow moves is noted. The horizontal meridian should be observed first, then the vertical. If the shadow appears to swirl round, not moving in the same meridian as the mirror, the eye is astigmatic, and the mirror is not moving in a direction which corresponds with either axis. A direction of movement can then be found in which the shadow will move either

directly with or against the mirror ; this is one of the principal axes of the astigmatism. The other axis is at right angles (regular astigmatism).

If the shadow moves with the mirror a low convex glass is put in the frame in front of the eye. If the shadow still moves in the same direction a stronger convex glass is used, and so on until no shadow can be seen. A still stronger convex glass is placed in the frame. The shadow now probably moves against the mirror. We now know that the refraction has been over-corrected. The point at which there is absolutely no shadow—the point of reversal—is somewhere between the last two lenses, and we know that at that point the refraction of the eye *plus* the lens is equivalent to one dioptre of myopia. If, for example, the shadow can still be seen to move with the mirror with + 4 D lens in the frame, and moves against it with + 5 D, we shall not be far wrong in considering that the point of reversal is + 4·5 D. A + 4·5 D lens would therefore make the eye one dioptre myopic. The actual refraction is therefore + 3·5 D. If there is no astigmatism the patient ought to be able to read 6/6 with this lens. If he is under atropin a further correction must be made before glasses are ordered. Atropin not only paralyses the ciliary muscle, but also inhibits the physiological tone of the muscle. This is found by experience to be equivalent to about 1 D of accommodation. Hence the glass which should be ordered to correct the distant vision after the mydriatic has passed off is + 2·5 D.



Fig. 249.

Similarly for spherical myopia. Supposing — 4 D eliminates the shadow against the mirror and — 4·5 D gives a distinct shadow with the mirror, we know that — 4·25 D will leave the eye with still — 1 D. Hence the refraction under atropin is — 5·25 D. The correction for atropin gives — 6·25 D as the lens which corrects distant vision without a mydriatic. The tone of the ciliary muscle is often less in myopia than in hypermetropia : since myopia should be under- rather than over-corrected, it is wiser to order very little more than the atropin correction, *e.g.*, 5·5 D in the above example.

In astigmatism each principal meridian is corrected sepa-

rately. When one meridian is approximately corrected the shadow assumes the shape of a band. The edge of the band is parallel to the axis of the corrected meridian. Even if the light is not moved in a direction accurately at right angles to this meridian the shadow still seems to move in the same direction. This is due to an optical illusion. If a straight edge, A B, is placed obliquely behind a circular hole in a card and is then moved horizontally in the direction of the arrow C, it will appear to be moving in the direction of the arrow D at right angles to its own edge. The shadow is most sharply defined if the mirror is moved at right angles to its edge, *i.e.*, at right angles to the corrected meridian.

The results are usually recorded thus (Fig. 250), the directions of the lines indicating the directions of the axes :—

The numbers should represent what the surgeon believes to be the refraction of the eye under the mydriatic, not numbers



Fig. 250.

to which further modifications have to be made, *e.g.*, the actual lenses used. In the examples given (*a*) is a case of regular simple hypermetropic astigmatism according to the rule ; (*b*) regular mixed astigmatism according to the rule ; (*c*) regular compound hypermetropic astigmatism against the rule ; (*d*) regular compound hypermetropic astigmatism with oblique axes. The exact direction of the axis in astigmatism is usually determined by subjective trial. In children it may be found by placing the requisite cylinder in the trial frame and rotating it until no shadow can be observed in any direction. In the examples given the correcting lenses required would be (*a*) + 2 D cylinder, axis vertical ; (*b*) — 1 D sphere combined with + 2·5 D cylinder, axis vertical, or + 1·5 D. sph. ⊂ — 2·5 D cyl., axis horizontal ; (*c*) + 1 D sph. ⊂ + 2 D cyl., axis horizontal, or + 3 D sph. ⊂ — 2 D cyl., axis vertical ; (*d*) + 2 D sph. ⊂ + 1·5 D cyl., axis 60° down and in, or down and out, according as it represents the left or the right eye.

The shadows in regular astigmatism are not always easy to

correct, owing chiefly to differences in curvature of different parts of the cornea. Usually the periphery of the cornea is flatter than the centre. The centre of the pupillary area will then be corrected by a different lens from the periphery, especially with the dilated pupil. From this cause various conflicting shadows may be seen, the commonest being the so-called "scissors" shadows, where two shadows appear to meet each other and cross as the light is moved in a given direction (Fig. 251). In conical cornea a triangular shadow with its apex at the apex of the cone, *i.e.*, usually slightly below the centre of the cornea, appears to swirl round its apex as the mirror is moved.



Fig. 251.

In irregular astigmatism the shadows move in various directions in different parts of the pupillary area; they cannot be accurately corrected by spherical or cylindrical lenses, but some improvement of vision may be obtained.

In conclusion, a word of warning must be given. The correction of a given refraction by retinoscopy may be extremely easy or extremely difficult. A vast number of refractions should have been carefully corrected and confirmed by subjective tests before a surgeon should consider himself justified in ordering glasses without supervision from an expert.

CHAPTER XXIV.

ERRORS OF REFRACTION.

Myopia, or "short sight," is that dioptric condition of the eye in which, with the accommodation at rest, incident parallel rays come to a focus anterior to the light-sensitive layer of the retina. Myopia may be due theoretically to any of the following conditions:—*A*. Abnormal length of the eye—*axial myopia*. *B*. Abnormal curvature of the refracting surfaces—*curvature myopia*: (*a*) too strong curvature of the cornea; (*b*) too strong curvature of one or both surfaces of the lens. *C*. Abnormal refractive index of the media—*index myopia*: (*a*) too high index of the cornea or aqueous; (*b*) too high total index of the lens, due to (α) too high index of the nucleus; (β) too low index of the cortex; (γ) both these causes; (*c*) too low index of the vitreous. *D*. Abnormal position of the lens, *i.e.*, displacement forwards. *E*. A combination of the above abnormalities.

There is no question that increased length of the eye is the most important factor in the high degrees. It is not improbable that the other factors are of more importance than is commonly thought in the lower degrees. This view is supported by the fact that, as shown by actual measurement, the radius of curvature of the normal cornea may vary from 7 to 8·5 mm. Curvature myopia occurs commonly as a factor in astigmatism, but is rare as a cause of spherical myopia, and is then associated with disease of the cornea—conical cornea. Index myopia is seldom seen clinically, but it accounts for myopia as a premonitory symptom of senile cataract, when it is due to increased refractive index of the nucleus of the lens; it also accounts for myopia in some cases of diabetes, with or without cataractous changes in the lens.

The increase in length of the eye affects the posterior pole and the surrounding area; the part of the eye anterior to the equator may be absolutely normal (Fig. 252). In most cases the myopia is of low degree, *i.e.*, up to 5 or 6 D (simple myopia). Less commonly the error reaches a considerable degree in childhood or early youth and increases steadily up to twenty-five or

more, finally amounting to 15 to 25 D or more (progressive myopia). It is impossible, clinically or pathologically, to draw a distinct line of demarcation between the two forms.

In low myopia the only symptom may be indistinct distant vision. In other cases and in high myopia there is often, in addition, discomfort after near work, due largely to disproportion between the efforts of accommodation and convergence (*vide* p. 199). The eyes are unduly sensitive to light. Black spots are seen floating before them, and some-



Fig. 252.—Horizontal sections of emmetropic and myopic eyes from the same patient superposed, showing the identity of the pre-equatorial regions. (Heine.)

times flashes of light are noticed ; the latter may occur irrespective of any tendency to detachment of the retina (*vide* p. 350). In very high myopia the eyes are prominent, the pupils are large, and the anterior chamber appears deeper than normal, probably only owing to the dilatation of the pupil. There may be an apparent convergent squint due to a large negative angle γ (*vide* p. 518). A true divergent strabismus may be found, either concomitant or affecting only one eye. Vision may be very poor, even with correction ; scotomata may be present, both central and peripheral.

Ophthalmoscopically, in low myopia there may be a quite

normal fundus; the optical defect will of course be noticed, especially on examination by the direct method. In the majority of cases of moderate myopia there is a "myopic crescent" (Plate XVIII., Fig. 2). This is a white crescent at the temporal border of the disc; very rarely it is nasal. In higher degrees of myopia it may extend to the upper and lower borders, or a complete ring may be formed round the disc. The crescent is occasionally absent even in cases of high myopia.

The bulging at the posterior pole in high myopia is called a posterior staphyloma. It is distinguishable clinically only



FIG. 253



FIG. 254.

FIG. 253.—Diagrammatic horizontal section of normal disc.
FIG. 254.—Diagrammatic horizontal section of myopic disc. *N*, nasal side; *T*, temporal side; *R. Pigment*, retinal pigment epithelium; *R*, retina; *Ch.*, choroid; *Scl.*, sclerotic. (Modified from Heine.)

by its optical and pathological effects. The term should not be used as a synonym for myopic crescent, as is often done. Optically posterior staphyloma causes the high error of refraction, and the edges may be actually visible by the indirect method owing to the presence of a crescentic shadow two or three disc diameters to the nasal side of the disc and concentric with it and to the change in course of the retinal vessels (staphyloma posticum verum). Pathologically, posterior staphyloma causes degenerative changes in the choroid and overlying retina: these are commonly described as "myopic choroiditis," but this term should be abandoned since the condition is non-inflammatory and should be called myopic choroido-retinal atrophy. The changes are generally

limited to the posterior pole and the surrounding area (Plate XVIII., Fig. 2). Small yellowish, white, or pigmented spots, and not infrequently white branched lines, usually horizontal, are found at and around the macula. The spots coalesce, forming irregular areas which may extend to the disc. Patches of choroidal atrophy are common near the disc; they may fuse with each other and with the myopic crescent so as to form a ring round the disc. Small foci occasionally occur at the periphery. Hæmorrhages in the macular region are generally described in high myopia. I believe them to be rare; the appearance of small hæmorrhages is very nearly simulated, but the spots undergo no change for an indefinite time; they are probably due to bunches of dilated



FIG. 255.—Age incidence of ametropia. (Herrnheiser.) Ordinates, percentages; abscissæ, ages.

capillaries, usually choroidal, rendered visible by rarefaction of the retinal pigment layer. The retinal pigment epithelium often loses much of its pigment in high myopia, so that the fundus is tigroid, and the choroidal vessels are well seen; this condition is not inconsistent with good vision.

Black specks in front of the eyes are often complained of in myopia. Dusty vitreous opacities may be visible with the ophthalmoscope (*vide* p. 104), or in high myopia large floating streamers. The normal muscæ volitantes are seen more plainly by myopic than by other eyes, probably because the entoptic image is generally larger.

A rare, but serious, change in the fundus is a small circular claret-coloured or black spot at the fovea. It may appear quite suddenly, being accompanied by great diminution of central visual acuity: it is probably due to intrachoroidal hæmorrhage or thrombosis.

Detachment of the retina is liable to occur spontaneously in high myopia (*vide* p. 350).

Ætiology.—Myopia is rare in the new-born, the percentage of cases increasing rapidly during the first two decades, remaining constant afterwards at 10 to 15 per cent., as compared with 30 per cent. for emmetropia and 50 to 55 per cent. for hypermetropia (Fig. 255). As regards sex, there is no doubt that the higher grades are commoner in women than in men. High myopia is as common among peasants as among the educated classes who do more near work. It is doubtful if near work is a cause of myopia, a view which has been held since the time of Kepler. There is, however, no doubt that it has a deleterious influence upon the disease. This has been attributed by Donders to (*a*) pressure of the extrinsic muscles upon the globe in strong convergence; (*b*) increased intraocular pressure from vascular congestion, due to the position of the head; (*c*) congestion of the fundus, leading to softening of the tissues. Accommodation has long been indicted as a cause of myopia. Many facts are against this view, *e.g.*, (*a*) accommodation occurs much more forcibly in hypermetropia; (*b*) it does not increase the intraocular pressure; (*c*) it does not affect the choroid farther back than the equator, and does not affect the sclerotic at all.

The mechanism whereby convergence influences the production of myopia is the subject of many theories, *e.g.*, pressure on the vortex veins, increased intraocular tension and so on.

Neither accommodation nor convergence alone suffices to explain the genesis of myopia. There must be some individual predisposition which provides any such accessory causes with advantageous conditions. There is probably a congenital weakness of the sclerotic. Other theories invoke shortness of the optic nerve (certainly false), special conformation of the skull leading to increased interpupillary distance or alteration in the position of the pulley of the superior oblique, &c. The view that moderate and high myopia are essentially distinct diseases, due to different causes, is probably untrue.

The cause of the myopic crescent has given rise to much discussion. It is probably congenital in origin, allied to other congenital crescents (*vide* p. 369), but there is no doubt that it may become altered by the conditions obtaining in the myopic eye. Anatomically there is considerable distortion of the papilla in myopia. It has been attributed to dragging, produced by the development of the posterior staphyloma: whether caused thus or not, it is an influential factor. Some

authors ascribe the crescent to this cause (distraction crescent). In well marked cases the head of the nerve is pulled over to the temporal side. The retina, including the pigment epithelium, is pulled slightly over the nasal edge of the disc (supertraction crescent). On the temporal side the pigment epithelium stops short at a variable distance from the disc and the choroid is atrophic here (Fig. 254). This part appears ophthalmoscopically as the crescent.

The fact that the crescent may be absent in high myopia and is often present in low militates against the view that it is caused entirely by traction. It is not due to accommodation for the same reasons that myopia is not caused by this factor.

As regards *prognosis*, low or moderate degrees of myopia (up to 5 or 6 D), unless occurring in young children, have a good prognosis. They are not likely to progress, and in some of the conditions of civilised life they may even be an advantage to the individual. The same condition in a child, before the age of schooling, is of grave prognosis, because it is almost certain to progress, so that in a few years there may be 10 or 15 D of myopia, accompanied by serious fundus changes and defects of vision. The prognosis in high degrees of myopia is always grave. It must be judged by the acuity of vision after correction and the condition of the fundus. In all cases there is some danger of retinal detachment occurring.

Treatment consists in wearing suitable correcting glasses and attention to the hygiene of the eyes. Each case must be considered on its merits.

As regards the ordering of glasses in myopia every surgeon agrees that *myopia must never be over-corrected*. Opinions differ as to details. In low myopia, up to 5 or 6 D (young children excepted), no harm is done by ordering the full distant correction for constant use, but if it is done the patient must be warned not to hold near work closer than ordinary reading distance. Many surgeons order glasses weaker by 2 or 3 D for near work. This has the effect of making the patients artificially presbyopic, *i.e.*, if they hold the work at reading distance they exert 2 or 3 D less accommodation than an emmetropic person would do, or than they themselves would do if wearing full correction. Many patients are more comfortable for near work with the weaker glasses; others derive no benefit. There is no doubt that the principle is derived from the fallacy that accommodation, *per se*, has a deleterious effect upon myopia. But there are some inherent objections to the weaker glasses. The patients often bring

their work closer than reading distance. As far as accommodation is concerned this matters less with the weak glasses than with the strong, but mere accommodation is of little importance. It is convergence which is the important factor. In order to read at reading distance there must be some convergence. If the artificial presbyopic correction is given no stimulation to converge is supplied by the act of accommodation, so that in order that physiological requirements may be satisfied the visual axes should be parallel. This can only be effected by combining the glasses with prisms, bases in (*vide* p. 500). If the work is held too near still more convergence is required, and the arguments apply still more strongly.

In general, in low myopia, the full correction may be ordered for constant use, with minute instructions as to near work; in the event of any discomfort being experienced, weaker glasses should be ordered for near work, especially if much reading and sewing, &c., is engaged in.

In high myopia it is wise always slightly to under-correct even for distance, and the same or still weaker glasses may be ordered for near work. In the highest grades the patient often sees best with glasses which are decidedly weaker than the full correction; he should be allowed to choose those he prefers. One reason is that the strong minus glasses very markedly diminish the size of the retinal images and make them very bright and clear. The retinal images are diminished because the glasses have to be worn farther from the eye than the anterior focal plane (*vide* p. 37); glasses for high myopia should therefore be made to fit as close to the eyes as possible. The very bright, clear images are uncomfortable because the retina is unduly irritable, probably owing to the fact that it has become accustomed to large indistinct diffusion images. Moreover much artificial astigmatism, and therefore distortion of the image, is produced by looking obliquely through strong glasses: it is found to be most disconcerting to those who begin the use of glasses or have them much strengthened late in life. Very short-sighted people get into the habit of turning the head rather than the eyes to avoid looking obliquely through the glasses. Some high myopes can find their way about much better without any glasses.

In very high myopia the requisite amount of convergence for near work may be impossible. Reading and other near work then becomes purely uniocular. Generally one eye is better than the other, and this eye is always used. The effort to maintain convergence under impossible conditions

is soon given up, which results eventually in the disused eye becoming divergent. There are other factors which tend to cause divergent strabismus in myopia (*vide* p. 529).

As regards hygienic measures in myopia, especially in the young, near work, apart from being held in the proper position, must be restricted. It is particularly important that it should not be done too continuously. More work can be done, with less harm to the eyes, by the interpolation of frequent short intervals of rest. It is best to give precise instructions as to the amount and distribution of near work. The illumination must be good, but not too bright, and it should come from behind and beside the patient's head. If the light is bad there is a strong tendency to bring the book or work closer to the eyes in order to enlarge the retinal image. Reading in bed and stooping over near work must be forbidden.

If the eyes are irritable, or the myopia is progressing, complete rest is imperative. Atropin should be instilled once a day for a prolonged period, tonics, especially iron and arsenic, should be given, and a change of air to the country, with plenty of healthy exercise, is desirable.

Operative treatment for high myopia.—If an eye has axial myopia of 24 D, its length will be about 31 mm. (*vide* p. 484). If the crystalline lens of such an eye is removed, parallel rays will be focussed upon the retina without the intervention of any correcting lens, and the retinal images of distant objects will be larger than those of the emmetropic eye. Hence the extraction of the lens has been strongly advocated in high myopia. In completely successful cases the improvement is enormous. The operation is, however, attended with grave dangers. The eye with high myopia is a diseased eye, which withstands operative measures badly. The vitreous is likely to be fluid and to contain opacities. The retina and choroid are probably diseased, and the tendency to detachment of the retina is increased by operation. No dogmatic rules can yet be given for the operation. I am guided by the following principles. (1) Only young patients should be operated upon; (2) the operation should be discission without subsequent curette evacuation unless it becomes imperative on account of tension; (3) there must be at least 15 D of myopia; (4) the fundus must be fairly healthy; (5) one eye only must be operated upon. The operation might be performed under less favourable circumstances if vision was so bad as to be useless, but such cases are rare.

Hypermetropia (*Syn.—Hyperopia*) or "far sight" is that

dioptric condition of the eye in which, with the accommodation at rest, incident parallel rays come to a focus posterior to the light-sensitive layer of the retina. Hypermetropia may be due theoretically to any of the following conditions: —*A*. Abnormal shortness of the eye—*axial hypermetropia*. *B*. Abnormal curvature of the refracting surfaces—*curvature hypermetropia*: (*a*) too slight curvature of the cornea; (*b*) too slight curvature of one or both surfaces of the lens. *C*. Abnormal refractive index of the media—*index hypermetropia*: (*a*) too low index of the cornea or aqueous; (*b*) too low total index of the lens, due to (α) too low index of the nucleus, (β) too high index of the cortex, (γ) both these causes; (*c*) too high index of the vitreous. *D*. Abnormal position of the lens, *i.e.*, displacement backwards. *E*. Absence of the lens—*aphakia*. *F*. A combination of the above abnormalities.

As in myopia, the chief factor in clinical hypermetropia is abnormality in the length of the eye; in this case the eye is too short. It must be remembered that a small eye, though too short, is not necessarily hypermetropic, since there may be uniform diminution of all the parts. This is, perhaps, most easily understood if a diagram such as Fig. 28 is considered; if such a diagram is uniformly diminished, *e.g.*, by photography, the parallel rays will still come to a focus on the retina. As a matter of fact hypermetropic eyes are almost invariably also smaller than normal, a fact which is of great pathological importance (*vide* p. 268).

Curvature hypermetropia occurs commonly as a factor in astigmatism; it is almost unknown as a cause of spherical hypermetropia. Index hypermetropia accounts for the hypermetropia of old age (*vide* p. 49), and it is to be attributed to increased refractive index of the cortex of the lens.

Hypermetropia rarely exceeds 6—7 D, which is equivalent to a shortening of the optic axis of 2 mm. Individual cases of much higher degrees, without other anomaly, such as coloboma or microphthalmia, have been recorded—up to 24 D.

In the young the condition may cause no symptoms. When symptoms are present or arise, they are chiefly referable to the abnormal amount of accommodation which these eyes are subjected to, and to the lack of consonance between accommodation and convergence (*vide* p. 499). As has been pointed out, the healthy youth has an ample reserve of accommodation, and if he happens to be hypermetropic he accommodates for distant and near objects without being conscious of the act. If he is weakly or does much near work the perpetual overaction

of the ciliary muscle is likely to produce symptoms ; the condition is often called accommodative asthenopia or " eye-strain." The symptoms are noticed chiefly after reading, sewing, &c., especially in the evening by artificial illumination. The eyes ache and burn ; they may feel dry, so that blinking movements are more frequent than usual, or there may be lacrymation. The conjunctiva and edges of the lids become red, and actual blepharitis may be caused. If near work is persisted in, headache, usually frontal, comes on. Typical migraine may occur.

In young children hypermetropia is a predisposing cause of convergent strabismus (*q.v.*). In all cases latent convergence is often found in hypermetropes, though other forms of heterophoria may occur (*vide* p. 530). The presence of heterophoria increases the tendency to headache, &c.

In older patients no symptoms may be caused until the power of accommodation has diminished to the extent that the far point is beyond the range of comfortable reading distance. Near work has to be held farther off than usual in order to be seen clearly. The greater the degree of hypermetropia the sooner will this symptom arise. In other words, apparent presbyopia commences at an earlier age than usual. It must be carefully borne in mind that hypermetropia predisposes to glaucoma (*q.v.*) in elderly people.

Ophthalmoscopically the fundus may exhibit no abnormality. A bright reflex, suggesting the appearance of watered silk, is commoner in hypermetropic than in emmetropic or myopic eyes. The inferior crescent is also more common in these eyes than in others, as also abnormal tortuosity of the retinal vessels. In some cases optic neuritis is nearly simulated—pseudo-papillitis (*vide* p. 359).

Anatomically the eye is shorter than normal in hypermetropia : it is also usually smaller. The changes are not confined to the post-equatorial segment as in myopia. The diameter of the cornea is often reduced, and regular astigmatism is common. The anterior chamber is shallower than normal, owing partly to the normal size of the lens (*vide* p. 267). Little weight should be attached to the old observation that the circular fibres of the ciliary muscle are hypertrophied, the meridional atrophied in hypermetropia. No anatomical abnormalities are found in the retina, choroid or optic nerve.

The new-born are almost invariably hypermetropic (mean 2·5 D). In the first decades of life the hypermetropia curve falls rapidly, remaining at about 50 per cent. after the twentieth year (Fig. 255). Hypermetropia shows no predilection for

either sex. It is a well-known fact that savages are usually hypermetropic. The higher mammals, especially the carnivora, are hypermetropic.

Treatment consists in prescribing the correcting glasses. Unless there are definite symptoms there is no reason for insisting upon the use of glasses in the young or middle-aged. In elderly people the hypermetropia must be corrected for near work: the ordinary presbyopic correction must be added to the hypermetropic correction, but care should be taken that these cases are rather under- than over-corrected (*vide* p. 489).

In young children the requisite correction is estimated under atropin, confirmed if possible by subjective tests. The correction, allowing for the effect of atropin upon the tone of the ciliary muscle (*vide* p. 469), is ordered for constant use or only for near work according to the severity of the symptoms. If the degree of hypermetropia is high the use of the glasses may be commenced while the child is still under the influence of atropin. In older patients with high hypermetropia it is often unwise to order the full correction at once. The ciliary muscle has been overworked so long that complete relaxation does not occur immediately. If the full correction is ordered the eye, with its contracted ciliary muscle, *plus* the glass, is made myopic: the patient cannot see clearly at a distance, and is liable to discard the spectacles. In these cases rather more than the amount of manifest hypermetropia should be ordered. The patient is told to return in three or six months, when stronger glasses are ordered, and so on until the full correction can be borne with comfort.

Astigmatism is that condition of refraction in which a point of light cannot be made to produce a punctate image upon the retina by any spherical correcting lens. The varieties of regular astigmatism have been already enumerated (*vide* p. 43).

Regular astigmatism, the only form which permits of optical correction, invariably produces greater or less defect in visual acuity. It is particularly liable to cause the worst forms of asthenopia or "eye-strain"; the asthenopia in these cases is only in part accommodative. It is often worse in the lower degrees of astigmatism than in the higher. This is probably due to the eye endeavouring so to accommodate as to produce a circle of least diffusion (*vide* p. 42) upon the retina. Aching of the eyes, severe headaches, and typical migraine are complained of; the eyes quickly become fatigued with reading, and the letters are described as "running together."

Regular astigmatism is usually a congenital defect, due in

most part to difference in curvature of the cornea in different meridians. It must be remembered that frequently the cornea is not alone at fault. Corneal astigmatism may be increased or partially corrected by lenticular astigmatism: hence the methods for correcting astigmatism, such as the ophthalmometer, &c., which are wholly dependent upon estimation of the corneal defect, are quite untrustworthy except in aphakia.

Regular astigmatism may be traumatic, following a wound, usually surgical, in the corneo-scleral margin. The contraction of the scar causes flattening of the cornea in the meridian at right angles to the wound. The astigmatism due to this cause continues to alter for many weeks after the injury, so that glasses should not be ordered for at least six weeks.

Treatment.—In all cases in which astigmatism causes asthenopic symptoms the full correction should be ordered for constant use, *i.e.*, both for distant and near vision. If there is a high degree of hypermetropia or myopia, associated with a low degree of astigmatism, the effect of the cylinder upon distant vision should be tested. If it produces no appreciable improvement simple spherical glasses should be tried first. It should be remembered that glasses placed before the eyes only correct the refraction accurately when the visual axis passes through the optical centre of the lens. When the eyes are directed to one side the lenses also act as prisms, and further the lenses are tilted relatively to the eyes so that an astigmatic effect is produced. In the case of high spherical lenses the astigmatic effect is considerable, and may easily counteract or double the effect produced by a weak cylinder combined with the sphere. Hence weak cylinders are seldom of much use when combined with high spheres.

In low astigmatic errors the instruction as to the use of glasses depends upon the amount of asthenopia. The relief of the discomfort experienced may not be worth the trouble of wearing glasses constantly. In these cases they should be ordered for near work only, and if this fails to eliminate the symptoms the advice should be given to wear them constantly.

Aphakia is the condition of the eye when the crystalline lens has been removed. The eye is extremely hypermetropic if it was emmetropic or had only a low grade of ametropia before removal of the lens. The hypermetropia, as estimated by the correcting lens required when worn in the usual position, is about 10 or 11 D if the eye was previously emmetropic.

The optical conditions of the aphakic eye are very simple. It consists of a curved surface, the cornea, separating two media,

air and vitreous, of different refractive indices. Knowing the radius of curvature (8 mm.) and the refractive indices (1 and 1·33) it is easy to calculate the focal distances. The anterior focal distance is 23 mm. and the posterior 31 mm., as compared with 13 mm. and 23 mm. respectively for the normal eye. If the aphakic eye were 31 mm. long parallel rays falling on the cornea would be brought to a focus on the retina and no correcting glass would be required for distance. It is easy to calculate the amount of axial myopia of a phakic eye which is 31 mm. long. It has been already pointed out that in the phakic eye 1 mm. of elongation is equivalent to an axial myopia of 3 D. Therefore an elongation of 31 — 23 mm., *i.e.*, 8 mm., equals 24 D (*vide* p. 479).

The retinal image of the aphakic eye is about one-third as large again as the emmetropic retinal image. Hence vision of 6/6 with a correcting glass after extraction is not quite as good as it seems.

Accommodation is, of course, lost. The anterior chamber is deep, the iris tremulous, and there is often a coloboma of the iris upwards. In cases of doubt as to the absence of the lens the Purkinje-Sanson reflexes from the lens surfaces should be sought.

With the ophthalmoscope opacities will probably be found in the pupillary area, consisting chiefly of remnants of the lens capsule. They should be examined by oblique illumination, by the mirror, and by the direct method. If they are dense, discission is indicated before attempting to correct the refraction ; if they are slight, the advisability of needling depends upon the amount of vision obtained with correction.

In addition to the hypermetropia, there is always some astigmatism in those cases in which a corneal or corneo-scleral section has been made. If the section is in the upper part of the cornea, the astigmatism is against the rule ; *i.e.*, the cornea is flattened in the vertical meridian. The astigmatism usually amounts to 2 or 3 D. It gradually diminishes, fairly quickly at first and very slowly after the first few weeks, as the cicatricial tissue in the scar contracts.

Treatment.—The refractive error is determined by retinoscopy and by subjective tests. The ophthalmometer may afford help in these cases. Great patience is often necessary, for the patients do not readily accommodate themselves to the new optical conditions. A 10 or 11 D convex lens combined with a + 2 or + 3 D cyclinder, axis horizontal, is about the correction usually required for distance. The sphere must be

stronger by 1 D for near work. A small amount of false accommodation can be obtained by slightly altering the distance of the glass from the eye. The correcting glasses should not be ordered earlier than six weeks after the operation, both on account of the necessity of resting the eye and because the astigmatism changes rapidly during the first few weeks.

If one eye only has been operated upon, the other being cataractous, reversible spectacle frames may be ordered. In them the bridge is horizontal, so that when the distant glass is being used the near glass is in front of the eye which has not been operated upon, and *vice versâ*. Reversible frames, however, never fit very satisfactorily.

The aphakic eye is specially liable to erythropsia (*q.v.*) and should therefore not be exposed to very bright light.

Anisometropia is the condition in which the refraction of the two eyes shows a considerable difference. A slight difference is very common. The condition may cause asthenopic symptoms. All varieties and degrees of anisometropia occur. In the lower grades there is usually binocular vision, though it is imperfect. In the higher grades it is impossible without correction. Vision is then uniocular, and there is some danger of the eye which is not used becoming divergent. If one eye is nearly emmetropic and the other myopic, the former may in some cases be used for distant, the latter for near, vision.

Treatment.— Correction of anisometropia offers many difficulties. It has already been mentioned that if correcting glasses are placed at the anterior focal plane of the eye, the retinal images are the same size as the emmetropic retinal image In practice the glasses are farther from the eyes. Consequently with convex glasses the retinal image is enlarged, with concave diminished. In high grades of anisometropia, therefore, there will be a considerable difference in the size of the retinal images. of the two eyes. Patients find it difficult or impossible to fuse these sharp but diverse images. Moreover on looking obliquely through the glasses the prismatic effect and the distortion are different in the two eyes, enhancing the discomfort. No universal rules can be given for the glasses which should be ordered. The following suggestions will generally be found to work well :—

If the difference between the two eyes is less than 4 D, the full correction should be ordered for constant use ; they should be perseveringly worn for at least six weeks. If still they cannot be borne, it will probably be necessary to correct only the less ametropic eye for distance.

In patients of less than twelve years of age the full correction should also be ordered for constant use, even if the difference is greater than 4 D. The more ametropic eye should be exercised alone, as in cases of concomitant strabismus (*vide* p. 527). Very often the treatment will fail, but it should be tried in the interests of binocular vision. It is almost certain to fail in older patients.

When the full correction cannot be worn constantly, both eyes can often be made to work together in comfort for near work by making each eye artificially presbyopic to the extent of half the difference between the two eyes. For example, suppose one eye is emmetropic and the other has 3 D of myopia, the patient will be most comfortable with + 1·5 D in front of the emmetropic, and — 1·5 D in front of the myopic eye for near work.

The Correction of Errors of Refraction.

The correction of errors of refraction has been already briefly sketched. It will be well, however, to outline the method to be adopted in systematically examining for and correcting these errors, and to indicate the requirements which should be satisfied by spectacles.

If the patient is less than fifteen years of age,—

(1) Test the distant and near vision, if the child knows his letters :

(2) Test the pupil reactions ;

(3) Test the muscular balance by the screen test (*vide* p. 531) ;

(4) Examine the fundi with the ophthalmoscope.

Then order ung. atropini, 1 per cent., to be inserted with a glass rod, three times a day for at least three days.

At the next visit—

(1) Determine the error of refraction by retinoscopy ;

(2) Thoroughly examine the fundus with the ophthalmoscope ;

(3) Confirm the retinoscopy by subjective tests, if the child knows his letters ;

(4) Order the correction according to the principles enunciated in the paragraphs devoted to the respective types of refractive error.

If the patient is between fifteen and twenty-five years of age, the same procedure should be adopted, but in many cases the prolonged action of atropin so seriously interferes with the patient's employment that it may be replaced by homatropin.

If the patient is between twenty-five and forty, —

(1) Test the distant vision, the manifest hypermetropia, and the near vision ;

(2) Test the pupil reactions and the range of accommodation ;

(3) Thoroughly examine the eyes with oblique illumination and by the ophthalmoscope.

If it is concluded that the defect is simply due to error of refraction, the further procedure depends upon the results of the subjective testing :—

(1) If the vision is 6/6 and J 1, with a low degree of manifest hypermetropia and few asthenopic symptoms, glasses may be ordered according to the amount of manifest hypermetropia (*vide* p. 482).

(2) If the vision is less than 6/6 no Hm, but J 1 is read fluently when the type is held closer to the eyes than normal, the patient has probably simple myopia. In general homatropin should be instilled, and the glasses ordered according to the retinoscopy and subjective tests under the mydriatic. The expert may feel justified in ordering glasses without using a mydriatic, judging by the subjective tests and his ophthalmoscopic examination, confirmed by retinoscopy without a mydriatic.

(3) If the vision is less than 6/6, and 6/6 cannot be read with any spherical glass, or if some letters only of 6/6 can be read—letters with oblique lines, *e.g.*, Z, being missed in that line and even in the other lines—the patient is probably astigmatic. Homatropin must be instilled, and the refraction corrected by retinoscopy.

If the patient is over forty, the examination will be exactly as for one between twenty-five and forty, except that presbyopia must be taken into account, and greater care must be exercised in instilling a mydriatic. Presbyopia affects the distant vision in hypermetropes in such a manner that although 6/6 may not be read with the unaided eye, it may be possible with a convex lens. Its effect upon near vision is discussed elsewhere. It is of little use to test the near vision of a presbyope without the glass which is necessary to correct the presbyopia, since no useful knowledge is obtained. If the vision cannot be improved up to 6/6 with a spherical lens, the patient is probably astigmatic, or has some disease of the eye, *e.g.*, incipient cataract. If ophthalmoscopic examination indicates merely the presence of astigmatism, homatropin (*never atropin*) should be instilled, but *in every case in which this is done one drop of eserin, 0·5 per cent., is instilled into the*

eye before the patient leaves. After estimating the error of refraction by retinoscopy it is advisable to see the patient again after the effects of homatropin have passed off before ordering glasses for near vision.

Spectacles.—In children spectacles with large round or "round oval" "eyes" should be ordered, otherwise the child may look over them.

In adults with astigmatism spectacles or rigid pince-nez must be ordered, never "folders." The latter are never to be recommended, and are absolutely contraindicated in astigmatism.

It is very important that all glasses fit accurately. In distant glasses the lenses must be centred so that the centres are exactly opposite the centres of the pupils when the visual axes are parallel. Near glasses are decentred slightly inwards, and the lenses are tilted so that the surfaces form an angle of 15° with the plane of the face: they are then approximately at right angles to the visual axes when the eyes are directed downwards in reading.

Various forms of bifocal glasses are sometimes used. In them the upper part contains the distant correction, the lower part the near If recommended, patients should be warned that they may experience some difficulty in going downstairs and so on, since objects on the ground will appear blurred when looking through the glasses.

If tinted glasses are desirable, *e.g.*, in high myopia, albinism, &c., the correcting lenses may be tinted. For use in tropical countries Crookes's glass (*vide* p. 142) may be used.

CHAPTER XXV.

ANOMALIES OF ACCOMMODATION.

Presbyopia has been already explained (*vide* p. 48). It is a physiological condition and therefore not properly included amongst anomalies of accommodation. It is convenient, however, to discuss its clinical treatment here, since it is so nearly allied to the correction of errors of refraction.

It has been shown that no convex lens of greater strength than 4 D should be ordered to correct presbyopia in the absence of hypermetropia; further, that the rule that a presbyope requires + 1 D for every five years after forty errs on the side of being a somewhat liberal allowance. Rather less, and never more, should be ordered. Some people, especially if they have hypermetropia, and therefore still stronger glasses for near work, have discomfort with their proper presbyopic correction. It is usually due to the absence of any stimulus, derived from the accommodative effort, to converge (*vide* p. 499). Theoretically the visual axes should be parallel when the presbyopic correction is used. Convergence, however, is necessary in order that both eyes may see the near object. It may be eliminated by combining prisms, bases in, with the correcting glasses.

In many occupations, *e.g.*, bootmaking, carpentering, and so on, the work is held at a greater distance than ordinary reading distance. The correction for work must then be ordered according to the distance, a weaker glass being required for a distance greater than 22 cm.

Myopes of course may require no glass for near work. Their presbyopic correction is estimated by the algebraic sum of their myopia and presbyopia.

Paralysis of Accommodation, or *cycloplegia,* occurs in disease as well as from the direct action of drugs (cycloplegics) such as atropin and homatropin. Unilateral cycloplegia is generally due to drugs, contusion (*vide* p. 400), or to paralysis of the third nerve. Bilateral paresis, less commonly paralysis, is most frequent after diphtheria, but may occur after debilitating illness, influenza, syphilis, diabetes, tabes, cerebral disease, &c.

Paresis of accommodation occurs as a premonitory symptom of glaucoma.

In complete paralysis the sphincter pupillæ is also generally paralysed, so that the pupil is widely dilated. In paresis the pupil may be scarcely at all affected, especially after diphtheria, but in this disease the reverse of the Argyll Robertson pupil may be met with, viz., loss of reaction to accommodation with retained reaction to light. The symptoms depend upon the condition of the refraction. If the patient is myopic, the defect may pass quite unnoticed; if he is emmetropic, near vision will be alone affected; if he is hypermetropic, both distant and near vision will be affected, but particularly the latter. In paresis it may be possible to diagnose the condition only by carefully measuring the range of accommodation.

In diphtheritic cases the paralysis of accommodation follows the primary attack at an interval of several weeks, and is often associated with paralysis of the palate, loss of knee jerks, &c. The sore throat may have been very slight and its diphtheritic character unrecognised. The lesion is probably nuclear, either toxic or hæmorrhagic. Cycloplegia in middle life should arouse suspicion of diabetes. It also occurs in chronic alcoholism.

The prognosis is good in cases due to drugs or diphtheria. In traumatic cases the condition may be permanent.

Treatment is that of the cause. Post-diphtheritic cases should be treated with tonics, especially strychnine. Whenever the condition is bilateral near work can be carried on by using suitable convex glasses, as in the correction of presbyopia. As a rule, however, the eyes should be kept at rest, so that it is inadvisable to order glasses. Miotics are sometimes used, but they may do harm and seldom do good. The constant current may be tried.

Spasm of Accommodation.—It has already been mentioned that the ciliary muscle has physiological tone, which is abrogated by atropin, and is equivalent to about one dioptre. In some cases it is found that atropin produces a much greater effect. This can only be due to spasm of the ciliary muscle. It is found only in young patients, and, contrary to what might be expected, more often in myopes than in hypermetropes. In any case an actual or relative myopia is produced. Spasm of accommodation is produced artificially by the instillation of miotics.

In spontaneous spasm of accommodation there is nearly always some error of refraction. The eyes have usually

been subjected to too much near work under unfavourable circumstances. The condition should not be diagnosed unless proved to be present by the use of atropin.

Treatment consists in the use of atropin for several weeks. The amount of near work must be limited and carried out under good conditions, the error of refraction being carefully corrected.

SECTION V.

DISORDERS OF MOTILITY OF THE EYE.

CHAPTER XXVI.

ANATOMY AND PHYSIOLOGY OF THE EXTRINSIC OCULAR MUSCLES.

The internal rectus is inserted into the sclerotic about 5·5 mm. to the nasal side of the corneo-scleral margin, the inferior rectus 6·5 mm. below, the external rectus 7 mm. to the temporal side, and the superior rectus 7·5 mm. above (Fig. 256).



Fig. 256.—Lines of insertion of the recti muscles of right eye seen from in front.



Fig. 257.—Lines of insertion of the superior oblique muscle and of the superior, external and internal recti of right eye, seen from above.

The tendons are about 10 mm. broad. The origin of these muscles around the optic foramen is much to the nasal side of the posterior pole of the eye. It has been proved that when the extrinsic muscles act they turn the eye around a spot which is called the centre of rotation (Fig. 258). This spot is situated about 13·5 mm. behind the centre of the cornea. It

lies in the same horizontal plane as the lateral recti. Consequently when the internal or external rectus acts it rotates the eye horizontally inwards or outwards respectively around a vertical axis through the centre of rotation without any rotation about the horizontal axis. When, however, the superior rectus acts, it not only pulls the eye upwards, but also inwards, while there is some rotation of the cornea, so that the vertical meridian assumes a direction from above down and out (torsion). Similarly when the inferior rectus acts the eye is pulled down and in, the vertical meridian of the cornea being deviated so that it lies from above down and in (Fig. 259).

The oblique muscles are inserted into the sclerotic behind the level of the centre of rotation (Fig. 257). Their direction of action is from behind forwards and inwards. Hence the superior oblique pulls the eye downwards and outwards, the inferior oblique upwards and outwards. The mechanism is so arranged that when the superior rectus and inferior oblique act simultaneously the eye moves directly upwards; *i.e.*, the upward movement caused by each muscle is summated, while the inward movement and corneal rotation of the superior rectus are exactly compensated by the outward movement and contrary corneal torsion of the inferior oblique. Similarly when the inferior rectus and superior oblique act simultaneously the eye moves directly downwards.



FIG. 258.—Diagram of the lines of action of the extrinsic muscles of left eye. (After Fick.) *C*, centre of rotation. The arrows show the directions of action of the muscles. *R.e.*, external rectus; *R.i.*, internal rectus; *R.s. & inf.*, superior and inferior recti; *O.s.*, superior oblique; *O.i.*, inferior oblique.

Every movement of the eyeball is a synkinesis (*vide* p. 51). In adversion not only does the internal rectus act, but also the superior and inferior recti, and it has been shown that the antagonistic muscles are not merely relaxed, but are actively inhibited. In abversion the external rectus and both obliques are in action. In elevation the superior rectus acts consonantly with the inferior oblique. In depression the inferior rectus acts with the superior oblique. The movements already

described are all around three primary axes—vertical (movements in and out), coronal (movements up and down), and sagittal (torsion)—which pass through the centre of rotation. Still more complicated are the movements about secondary axes, *i.e.*, axes passing through the centre of rotation in some other direction, such as movements up and in, up and out, down and in, down and out.

Not only is there uniocular synkinesis: under normal circumstances there is always also binocular synkinesis. Adversion of one eye is accompanied by abversion of the other eye—conjugate deviation; elevation or depression of one eye is always accompanied by elevation or depression respectively of the other eye. The only exception to this rule is the bilateral



Fig. 259.—Diagram of the lines of action of the extrinsic muscles upon the corneæ. (Elschnig.) R, right eye; L, left eye; T, temporal sides; N, nasal sides. The dotted lines show the torsional effects.

adversion of the eyes in convergence. Elevation of both eyes is accompanied by slight abversion (divergence), depression by slight adversion (convergence).

The oculomotor, or third cranial nerve, supplies all the extrinsic muscles except the external rectus and superior oblique; it also supplies the sphincter iridis and ciliary muscle. The superior oblique is supplied by the fourth nerve, and the external rectus by the sixth nerve. A thorough knowledge of the arrangement of the nuclei of the cranial nerves in the midbrain and medulla, and of the course and relations of the nerves to their destinations, is requisite for accurate diagnosis of the seat of the lesion in cases in which they are involved.

The third and fourth nuclei form a large continuous mass of nerve cells situated near the middle line in the floor of the

aqueduct of Sylvius beneath the corpora quadrigemina or colliculi (Figs. 260—62). The cells nearest the middle line towards the anterior part of the third nucleus are smaller than the others: they, with the cells of the opposite side, form an unpaired nucleus with two divergent horn-like processes in front (the Edinger-Westphal nuclei) (Fig. 264), which probably supplies fibres to the ciliary muscle (accommodation) and sphincter iridis (constriction of the pupil). It is probable that in the great large-celled lateral nucleus the levator palpebræ is represented most anteriorly, then from before backwards elevation of the eye, adversion, and depression, while abversion is relegated to the sixth nucleus, much farther back in the medulla (Fig. 264). There is little decussation of the fibres from the third nuclei of the two sides in the anterior part, but a considerable amount in the posterior part.



Fig. 260.—Diagram of the positions of the nuclei of the third, fourth, and sixth nerves, seen from above. *S.c.q.*, superior corpus quadrigeminum; *I.c.q.*, inferior corpus quadrigeminum.

The fourth nerve is unique among motor nerves in having a dorsal decussation. Nearly, if not quite, all the fibres decussate in the superior medullary velum and are distributed to the superior oblique muscle of the opposite side.

The sixth nucleus is in the immediate vicinity of the facial (seventh) nucleus (Figs. 260, 263), the fibres from which make a large bend around the sixth nucleus (Fig. 265). Hence vascular and other lesions of the sixth nucleus are very liable to be accompanied by facial paralysis on the same side. All the fibres of the sixth nerve are distributed to the external rectus of the same side.

The peculiarities of distribution of the fibres from the third, fourth and sixth nuclei to muscles partly on one side and partly

on the opposite side of the body show that the nervous mechanism of co-ordination of these muscles is extremely complex.



FIG. 261.—Diagram of transverse section of the mesencephalon at the level of the third nucleus (level of 1, Fig. 266). *Supr. c.q.*, superior corpus quadrigeminum; *R*, red nucleus; *I.g.b.*, internal geniculate body; *E.g.b.*, external geniculate body; *O.Tr.*, optic tract.



FIG. 262.—Diagram of transverse section of the mesencephalon at the level of the fourth nucleus (level of 2, Fig. 266). *Aq.*, aqueduct of Sylvius; *Inf. c.q.*, inferior corpus quadrigeminum; *p.l.b.*, posterior longitudinal bundle; *Supr. ped.*, superior peduncle of the cerebellum; *Pyr. Tr.*, pyramidal tract.

A large and important tract of nerve fibres, derived in part from the anterior columns of the spinal cord, lies below and close to the third, fourth and sixth nuclei. This is the posterior longitudinal bundle (Figs. 261 — 63, 266). Fibres pass between it and the nuclei under consideration; they probably have important functions in the co-ordination of movements and equilibration, which are so intimately related with vision. Among these fibres are also some which link up the sixth nucleus of one side with the third nucleus of the other in some such manner as depicted in Fig. 266, though the exact course of the fibres has not been definitely proved. These fibres are concerned in conjugate deviation of the eyes to one or other side. Hence when one sixth nucleus is destroyed the patient is unable to turn his eyes to the same side, though the power of convergence is unimpaired. Nuclear sixth nerve paralysis therefore causes loss of conjugate deviation of the eyes to the same side, and is very likely to be associated with facial paralysis on the same side, whereas peripheral sixth nerve

paralysis causes only loss of power of movement of the same eye to the same side.

The student should revise his knowledge of the anatomical relations of these and the neighbouring cranial nerves in their course from the nuclei to their respective terminations.



FIG. 263.—Diagram of transverse section of the pons at the level of the sixth nucleus (level of 3, Fig. 266). *p.l.b.*, posterior longitudinal bundle; *D*, Deiter's nucleus; *s.o.*, superior olive; *C.r.*, restiform body; *Pyr. Tr.*, pyramidal tract.

Orientation. — Orientation of objects in space depends upon their relation to the nodal point of the eye, *i.e.*, the position of an object is determined by the line passing through the object and the nodal point, the spot where this line cuts the retina being the position of the retinal image of the object. Conversely an object is said to be *projected* along the line joining the retinal image with the nodal point. Objective orientation determines the



FIG. 264.—Diagram of the probable position of the nuclei of origin of the fibres to the ocular muscles in the third, fourth and sixth nuclei.

relative positions of objects to each other. Subjective orientation, or the exact relation of the situations of objects to ourselves, is much more complex, depending upon an

accurate knowledge of the position of the body and of the eyes in the body, derived largely from the muscular sense.

Corresponding Points.—When a distant object is looked at the visual axes are practically parallel: the object forms an image upon each fovea centralis. An object to one side of the object looked at forms its retinal images upon the temporal side of one retina and upon the nasal side of the other; these are called corresponding points. Points on the two retinæ which are not corresponding points in this sense of the term are called *disparate points*. If an object forms its retinal images upon disparate points it will be seen double (binocular diplopia). If the disparity is slight there is a great tendency to move the eyes so that the images may be fused. It will be noticed that the two foveæ are corresponding points.



FIG. 265.—Diagram of the sixth nucleus and its relation to the nucleus and the emerging fibres of the seventh nerve. The relations cannot be accurately depicted in one plane.

When a near object is looked at the eyes converge the requisite amount to bring the two retinal images of the object upon the two yellow spots.

Binocular Vision.—When the eyes are normal the individual sees clearly with both eyes the object looked at. The retinal images of the two eyes are not, however, identical. This is obvious when it is remembered that there is a considerable distance between the two eyes. If the object is a solid body, *e.g.*, a cube, the right eye sees a little more of the right side of the object, and *vice versâ*. The two images are fused psychologically, and it is this fusion of the slightly diverse images, combined with other facts derived from experience, which enables the person to appreciate the solidity of objects. The estimation of the relative distance of objects in or near the line of vision is still more complex. It probably depends upon the fact that the images of objects farther or nearer than the object fixed are situated at disparate points on the retinæ. A more distant object will produce heteronymous diplopia, a nearer homonymous diplopia, as can be shown by experiment. The diplopia is suppressed in actual vision, but it produces a psychological impression which is translated into appreciation

of distance. It will suffice if it is well understood that accuracy of stereoscopic and topical vision depends upon good sight with both eyes simultaneously.

Convergence and Accommodation.—When a distant object is observed by an emmetropic person the visual axes are parallel and no effort of accommodation is made. If a near object is observed the eyes converge upon it and an effort of accommodation corresponding with the distance of the object is made. Convergence can be tested roughly by making the patient fix a finger or pencil which is gradually brought nearer to the eyes in the middle line. The eyes should be able to maintain convergence when the object is 8 cm. ($3\frac{1}{2}$ inches) from the eyes. If outward deviation of one eye occurs before this point is reached the power of convergence is deficient. There are various methods of recording the amount of convergence. One very convenient method employs the *metre angle* as a unit. Suppose an object to be situated in the median line between the two eyes at a distance of one metre from them. Then the angle which the line joining the object with the centre of rotation of either eye makes with the median line is called one metre angle (Fig. 267). With an interpupillary distance of 60 mm. this angle is about 2°. If the object is two metres away the angle is approximately half as great, or $\frac{1}{2}$ m.a. If the object is 50 cm. away the angle will be 2 m.a. Now, the amount of accommodation which an emmetropic eye exercises in order to see clearly an object 1 m. away is 1 D, 2 m. away 0·5 D, 50 cm. away 2 D, &c. Hence with an emmetropic person the amount of convergence, reckoned in metre angles, is the same as the amount of accommodation reckoned in dioptres.



FIG. 266.—Diagram of the course of the fibres from the sixth nucleus which are concerned in conjugate deviation of the eyes. 1, 2, 3, lines of section of Figs. 261–263.

The amount of convergence can also be measured by prisms. If an object one metre distant is looked at through a prism

with the base directed outwards placed before one eye, *e.g.*, the right, it may still be seen as a single object (Fig. 268). Now in order that the object may form its image upon the fovea of this eye it is necessary that the eye should be turned inwards an amount corresponding with the angle of deviation of the prism (*vide* p. 24). This method, besides affording a method of recording amounts of convergence, also shows that the relationship between convergence and accommodation is somewhat elastic. In the experiment described, although the amount of accommodation exercised remains the same, the amount of convergence is altered. Indeed, if the relationship were quite unalterable a hypermetropic person would invariably have diplopia, for his accommodation is always in excess of the corresponding value of the amount of convergence exerted by an emmetrope. Moreover, the power to converge would gradually be lost *pari passu* with loss of accommodation in advancing age.



FIG. 267.—Diagram of the metre angle. *Cr.*, *Cl.*, centres of rotation of the right and left eyes.

If in the experiment described above the prism is held before the right eye with its base inwards it will still be found possible to see the object single (Fig. 269). Further, a distant object is still seen single under the same conditions if the prism has only a small angle of deviation. This can only be accomplished by active divergence of the eyes to an amount corresponding with the angle of deviation of the prism. The power of divergence, which may therefore be considered to be negative convergence, is much less than the power of convergence. Just as the difference in the amount of accommodation between the far point and the near point is called the amplitude of accommodation, so the difference in convergence between the far point and the near point is called the *amplitude of convergence*. Whereas, however, negative accommodation, *i.e.*, ability to see a point beyond infinity, or in less mathematical terms, so to flatten the lens that a myope could see clearly without glasses, is impossible, negative convergence, as has been seen, is possible within small limits. The amplitude of convergence therefore

consists of a negative portion and a positive portion. The former is measured by the strongest prism, base inwards, which can be borne without producing diplopia in distant vision. The latter is measured by the strongest prism, base



FIGS. 268–269.—Diagrams of the action of adducting and abducting prisms. O, object of fixation ; *f*, *f′*, left and right foveæ centrales.

outwards, which can be borne without producing diplopia in the nearest possible vision.

The convergence synkinesis is so co-ordinated that the energy exerted is accurately divided between the two internal recti. Hence it is found that the effect is the same in the above experiments whether the prism is placed before only one eye, or a prism of half the strength is placed before each eye.

CHAPTER XXVII.

PARALYTIC AND KINETIC STRABISMUS. SYNKINESES. NYSTAGMUS.

Strabismus (στρέφειν, to turn) or *squint* is a generic term applied to all those conditions in which the visual axes assume a position relative to each other different from that required by the physiological conditions. Strabismus may be provisionally divided into two great groups : (1) those due to known cause ; (2) those due to unknown cause. To the first group belong (*a*) those due to paresis or paralysis of one or more of the extrinsic muscles—*paralytic strabismus ;* (*b*) those due to irregular activity or over-activity of individual muscles or groups of muscles—a sub-group which I propose to designate *kinetic strabismus.* To the second group belong (*a*) those cases which are characterised by the fact that the visual axes, though abnormally directed, retain their relative position in all movements of the eyes ; they are therefore termed *concomitant* or *comitant strabismus.* Another sub-group of the second class is (*b*) cases in which there is *latent strabismus* or *heterophoria.*

Paralytic Strabismus.

Signs and Symptoms.—(1) *Limitation of Movement.*—In paralysis of an ocular muscle the ability to turn the eye in the direction of the normal action of the muscle is diminished or lost. In slight paresis the defect in mobility may be so small as to escape observation without special tests. In all positions in which the affected muscle is not brought actively into play the visual axes assume their normal relationship.

Limitation of movement is tested roughly by fixing the patient's head and telling him to follow the movements of the surgeon's finger. The finger should be held vertical in testing horizontal movements, horizontal in testing vertical movements. An accurate record of the movements of each eye can be obtained by taking the field of fixation. The patient is seated at the perimeter as for recording the field of vision. With the head fixed and the other eye screened the patient

looks as far as possible along the arc of the perimeter, test types being moved in from the periphery until he is just able to read them. The normal field of fixation is about 50° downwards and 45° in all other directions.

When the eyes are turned in the direction of the normal action of the paralysed muscle the affected eye remains stationary. It deviates therefore relatively to the other eye; this position is called the *primary deviation* (Fig. 270).



FIG. 270.—Diagram of primary deviation in paresis of the right external rectus. *p*, angle of primary deviation; *n*, *n'*, left and right nodal points.

The angle of deviation is the angle which the line joining the object observed with the nodal point makes with the visual line.

If the sound eye is covered by a screen, and an attempt is made to fix an object so situated that the paralysed muscle is brought into play, it will be found that the eye behind the screen deviates more than the primary deviation of the paralysed eye. For example, if the right external rectus is paralysed and the left eye is covered, then on attempting to fix an object situated to the right with the right eye the left eye will deviate very much to the right, so much in fact that

its line of vision is well to the right of the object fixed. Hence, if the screen is removed suddenly the left eye will spring back to the left so as to take up fixation. This deviation of the sound eye is called the *secondary deviation* (Fig. 271). The reason why the secondary deviation is greater than the primary is that in conjugate deviation of the eyes the nervous energy is equally distributed between the muscles of the two eyes. Now the effort to take up fixation with the paralysed or



FIG. 271.—Diagram of secondary deviation in paresis of the right external rectus. *S*, screen in front of left eye ; *s*, angle of secondary deviation.

paresed eye is much greater than normal. Consequently the sound eye behind the screen moves through a greater distance than normal, *i.e.*, through a distance corresponding with the excessive effort exerted. This feature is of great importance because when well marked it distinguishes paralytic squint from the concomitant type in which the secondary deviation is equal to the primary.

(2) *Diplopia.*— The chief complaint of patients with paralysis of an extrinsic muscle is often that they see double. Diplopia

occurs only over that part of the field of fixation towards which the affected muscle or muscles move the eye. If both eyes are functional and one deviates, *i.e.*, if the visual axes are not parallel in looking at a distant object, or if the amount of convergence is not accurately adapted to the position of the object in near vision, *binocular diplopia* results. When the deviation is due to paralysis of one or more extrinsic muscles, the eye on the sound side fixes the object accurately, while the other eye deviates. Suppose the left eye fixes accurately while the right deviates inwards, a bright, sharply defined foveal image is seen with the left eye. The image formed by the object on the right retina, falling as it does upon the line joining the nodal point with the object, lies to the nasal side of the retina. The patient being unconscious of the malposition of his eye orients the object subjectively as if the eye were straight. He knows from experience that objects which form their images upon the nasal side of the retina are situated to the temporal side. He therefore projects the object with this eye to the right of its actual position. This is called *homonymous diplopia*, because the object as seen by the right eye is to the right of the object as seen by the left eye (Fig. 272).



FIG. 272.—Diagram of homonymous diplopia. *f*, *f'*, left and right foveæ; *n*, *n'*, left and right nodal points. The image of O formed at *a* is projected as if *a* were the fovea, *i.e.*, to O'.

If the right eye deviates outwards, *heteronymous* or *crossed diplopia* results, because the object as seen with the left eye lies apparently to the right of the object as seen by the right eye (Fig. 273).

In binocular diplopia the image seen by the squinting eye (false or apparent image) is less distinct than that seen by the fixing eye (true image), because only in the latter case does the image fall upon the fovea centralis. The angular displacement of the false image is equal to the angle of deviation of the eye.

(3) *False Orientation.*—It will be seen from what has

already been said that false orientation is a necessary accompaniment of binocular diplopia. Suppose that a patient whose right external rectus is paralysed shuts his left eye and attempts to fix an object situated towards the right. Let him now quickly strike at the object with his extended index finger. The finger will pass considerably to the right of the object. This is called *false projection*. It depends upon exactly the same principle as the increase of the secondary deviation. The object is projected according to the amount of nervous energy exerted; as this is greater than that exerted under normal circumstances, the object is projected too far in the direction of action of the paralysed muscle. It is essential that the finger should be directed at the object quickly, otherwise the error is noticed and compensated for. For example, if under the same circumstances the patient is told to walk towards an object situated at some distance to the right, he first steps too far to the right, then recognises his mistake and corrects it. In old paralysis the patient may learn by experience completely to compensate for the deficiency.



Fig. 273.—Diagram of heteronymous (crossed) diplopia.

(4) *Position of the Head.*—The patient holds his head so that his face is turned in the direction of action of the paralysed muscle. For example, in paralysis of the right external rectus the patient keeps his head turned to the right. The object of this manœuvre is to abolish the diplopia and its attendant unpleasant consequences as much as possible. In complex paralysis the position of the head is still such as to relieve the diplopia to the maximum extent, the position being adopted unconsciously.

"*Ocular torticollis*" is a term sometimes applied to tilting of the head to compensate defective vertical movements of one eye. It is distinguished from true torticollis in that there is a simple tilting of the head, the chin not being rotated towards the opposite shoulder; moreover, the sterno-mastoid is not unduly contracted.

It occurs chiefly in cases of congenital origin—probably malinsertion of the muscles—but has been met with after interference with the pulley of the superior oblique in frontal sinus operations. The vertical defect is made manifest by placing the head straight, when diplopia is also elicited.

(5) *Vertigo, &c.*—Vertigo, leading to nausea, and even vomiting, is due partly to diplopia, partly to false projection. It occurs chiefly when the paralysed muscle is called upon to exert itself. When the gaze is turned from the region of correct to that of false localisation, objects appear to move with increasing velocity in the direction in which the eye is moving. The unpleasant symptoms are counteracted partially by altering the position of the head, or completely by shutting or covering the affected eye.

In paralyses of long standing, false orientation gradually ceases (*vide supra*). Diplopia also tends to disappear or become less troublesome; the patient learns to ignore the impressions derived from the affected eye. Contracture of the antagonists of the paralysed muscle gradually sets in, which has the effect of increasing the primary deviation. Since the retinal image is thus thrown farther to the periphery, where the sensitiveness is less (*vide* p. 62), its suppression is facilitated.

Investigation of a Case of Ocular Paralysis.—The patient usually seeks advice on account of diplopia. In some cases the nature of the case is obvious immediately from the strabismus or from the manner in which the head is held. In most cases these features are too slight to decide the diagnosis.

(1) The first procedure should be to cover one eye in order to determine whether the diplopia is uniocular or binocular.

(2) Having decided that the diplopia is binocular the patient should fix the surgeon's finger, and the field of fixation of each eye should be carefully investigated (*vide* p. 506). In cases of complete paralysis of one or more muscles it may be possible to make an accurate diagnosis from the observation of the defective movements combined with investigation of the exact positions of the images of the finger in different areas of the field of binocular fixation. In cases of paresis the differentiation of the images is too obscure to permit of the solution of the problem by this means.

(3) In such cases the diplopia must be investigated by more delicate tests. The patient is taken into a dark room. A red glass is placed before one eye in order to distinguish its

image. A lighted candle is then moved about in the field of binocular fixation at a distance of at least four feet from the patient, the patient's head being kept stationary. The positions of the images are accurately recorded upon a chart with nine squares marked upon it (Fig. 274). The examination may be carried out by the surgeon turning the patient's head in various directions while the candle is kept stationary. The following data are derived from this examination :—

(*a*) The areas of single vision and diplopia ;

(*b*) The distance between the two images in the areas of diplopia ;

(*c*) Whether the images are on the same level or not ;

(*d*) Whether one image is inclined or both are erect ;

(*e*) Whether the diplopia is homonymous or crossed.



Fig. 274.—Diplopia chart for the right external rectus. The oblique line through the chart shows the limit of the fields of single vision and of diplopia. The dotted arrows show the positions of the false image in different parts of the field of diplopia.

These data, if concordant, are sufficient to diagnose the paralysis. The false image is determined by the direction in which the images are most separated from each other. This is the direction of the normal action of the paralysed muscle. The false image can often be recognised by being the fainter of the two or by being tilted ; by covering one eye it can be shown to which eye this image belongs.

It must be remembered that these tests are purely subjective. In many cases the patients are stupid or their intelligence is obscured by intracranial disease ; or contracture of the antagonistic muscles may have set in. Consequently the answers are not infrequently discordant, and accurate diagnosis may be extremely difficult or impossible. There are two not infrequent causes of ambiguity. The paresis may unmask a latent squint (*vide* p. 530), or the patient may fix with the paralysed eye, especially if this eye has the greater acuity of vision.

The nature of the diplopia and the position of the images in each of the nine areas of the field of fixation for paralysis of each individual muscle should be worked out by the student. In performing this exercise he should rely upon his knowledge

of the anatomy of the muscles and their consequent action in each position of the eye. Considerable ingenuity has been used to devise mnemonics for determining the position of the false image. One of the most satisfactory is shown in the accompanying diagram (Fig. 275). It may be pointed out that all the signs, with the exception of the deviation of the eye, viz., defective movement, false projection, increase of diplopia, secondary deviation, and position of the head, are towards the side of the paralysed muscle.

Varieties of Ocular Paralysis.—If one muscle alone is affected



FIG. 275.—Werner's mnemonic for paralyses of elevator and depressor muscles. Note that the right muscles are to the right, the left to the left; recti in centre, obliques outside, superior recti above, superior obliques below. The diagram illustrates the normal actions of the muscles and the type of diplopia caused. For example:—*Actions:* The recti adduct, the obliques abduct; the superior muscles (*R.s.r.*, *L.s.r.*, *R.s.o.*, *L.s.o.*) produce inward torsion, the inferior muscles outward torsion; the movement follows the direction of the arrow, *e.g.*, *R.s.r.* moves the eye inwards and upwards and causes inward torsion. *Diplopia:* The false image (broken arrows) is displaced in the direction of action of the muscle: therefore, for muscles in the upper half (*L.i.o.*, *L.s.r.*, *R.s.r.*, *R.i.o.*) the diplopia occurs on upward movement of the eyes, and the false image is higher than the true. The diagram also shows whether the diplopia is homonymous or crossed. If the patient fixes with the paralysed eye the figure must be rotated so that the false image becomes vertical.

it is generally the external rectus or the superior oblique, since each of these is supplied by an independent nerve.

Affection of several muscles simultaneously is usually due to paralysis of the third nerve. All the extrinsic and intrinsic muscles of one or both eyes may be paralysed—*ophthalmoplegia totalis* (*vide* p. 546). If only the extrinsic muscles are affected the condition is called *ophthalmoplegia externa;* if only the intrinsic (sphincter pupillæ and ciliary muscle) *ophthalmoplegia interna.*

Conjugate paralysis is the term applied to abolition of certain

synkineses. Thus ability to look up, to the right or left, or down may be lost. Inability to converge may also occur. Such defects might be conveniently termed *symparalyses.*

Paralysis of the External Rectus.—There is limitation of movement outwards, and the face is turned towards the paralysed side. Diplopia occurs on looking to the paralysed side. It is homonymous; the images are on the same level, and erect, becoming more separated on looking more towards the paralysed side. The false image is slightly tilted on looking up or down as well as towards the paralysed side.

Paralysis of the Superior Oblique.—There is limitation of movement downwards and towards the paralysed side; the face is turned downwards and towards the *sound* side. Diplopia occurs on looking down (Fig. 276). It is homonymous;



FIG. 276.—Diplopia chart for the right superior oblique.



FIG. 277.—Diplopia chart for the right third nerve. The area enclosed by the curved line is the area of single vision.

the false image is lower and its upper end is tilted towards the true image. The distance between the images and the inclination of the false image increase on looking down and towards the paralysed side. The patient has great difficulty in going downstairs, and vertigo is usually a particularly prominent symptom.

Paralysis of the Third Nerve.—In complete paralysis of the third nerve there is ptosis, which prevents diplopia. On raising the lid with the finger the eye is seen to be deflected outwards and somewhat downwards, owing to the tone of the two unparalysed muscles. The pupil is semi-dilated and immobile, and accommodation is paralysed. There is a slight degree of proptosis, owing to loss of tone of the paralysed muscles. There is limitation of movement upwards and inwards, to a less degree downwards. With the lid raised

there is diplopia, which is crossed, the false image being higher, with its upper end tilted towards the paralysed side (Fig. 277).

Paralysis of the third nerve is often incomplete, and individual muscles may occasionally be affected alone.

Ætiology.—Paralysis of ocular muscles may result from a lesion situated in any part of the nerve tracts from the cerebral cortex to the muscles. The site may therefore be intracranial or intraorbital. Cortical lesions usually cause loss of synergic movements, *e.g.*, conjugate deviations, but simple ptosis may be due to such a lesion. The diagnosis of nuclear and peripheral lesions depends largely upon knowledge of the anatomical relations of the nuclei and nerves. It is beyond the scope of this work to treat the subject exhaustively here. It may be mentioned that paralysis of the external rectus, sometimes bilateral, is common in babies. It may be due to the use of forceps during delivery, the sixth nerve being most exposed to pressure, or to maldevelopment of the nucleus. In the latter case there is loss of conjugate deviation to the same side (*vide* p. 496), and the seventh nerve is normal. In acquired sixth nuclear paralysis in adults the seventh nerve is usually implicated (*vide* p. 496). In congenital paralysis of the external rectus contracture of the antagonists does not occur. Other nuclei may be maldeveloped, or they may be picked out by the lesions of syphilis, tabes, or disseminated sclerosis.

The commonest cause of ocular paralyses is syphilis, which may affect the nerves at their origin or in any part of their course (see Section VI.). Syphilitic paralysis is usually a late manifestation. The third nerve is affected most frequently, but not necessarily equally in all its branches. Tabes is responsible for a large proportion of the cases (*vide* p. 541).

Ocular paralyses of intracranial origin may be due to affections of the blood vessels—hæmorrhage, thrombosis, etc.—or to external pressure—tumours, bloodclots, periostitis, etc.

Other causes are diphtheria, diabetes and other toxic conditions, injury, etc. Paralysis of both extrinsic and intrinsic ocular muscles is a common and early feature in encephalitis lethargica. Paralysis, especially of the external rectus, sometimes follows spinal anæsthesia with stovain, etc. Ophthalmoplegic migraine is a rare cause (*vide* p. 379).

Paralysis of the external recti is common in cases of intracranial tumours with high intracranial pressure, and generally has no localising value. It may be due to traction on the nerves, or more probably to pressure by the anterior inferior

cerebellar and internal auditory arteries, which cross them at right angles and often lie ventral to them—the nerves are strangulated between the vessels and the œdematous and swollen pons (Cushing). This may also account for the spinal anæsthetic cases, and for ophthalmoplegic migraine, the third nerve passing between the superior cerebellar and the posterior cerebral arteries.

Injury to and disease of the orbit may affect the nerves of muscles in this situation by rupture, pressure, inflammation, &c.

The prognosis varies with the cause. Ocular paralyses are so often early signs of grave nervous disease that the prognosis should always be guarded. Those due to peripheral disease, dependent upon syphilis or "rheumatism," may speedily recover. Long-standing cases rarely recover.

Treatment.—Syphilitic cases should be treated by mercury and increasing doses of iodides, and this treatment should be applied to all doubtful cases. Salvarsan should be tried in intractable cases. Some cases improve on salicylates, or colchicum and iodides. Diaphoresis may be used. The constant current is chiefly useful in keeping up the metabolism of the muscles until innervation is re-established. It probably has little therapeutic influence upon the paralysed nerve.

Occasionally symptomatic treatment affords relief to the patient. The diplopia may sometimes be relieved by suitable prisms, but this treatment is rarely of much use owing to the variation in the amount of the deviation in different positions of the eyes. Occasionally good is done by exercising the weak muscle with strong prisms (*vide* p. 534). In old cases an operation may be indicated, usually tenotomy of the antagonist with advancement of the paralysed muscle, thus putting the affected muscle under better mechanical conditions. It is only suitable for paretic, not paralytic, cases, and should never be adopted until all other means have failed. It is therefore seldom indicated.

If diplopia is very troublesome and cannot be relieved by the means suggested, spectacles should be ordered with a ground glass in front of the affected eye.

Kinetic Strabismus.

Aberrant forms of strabismus occur as the result of irritative intracranial lesions, and are due, not to paralysis, but to irregular action or over-action of certain muscles, caused by

unequal stimulation of the nerve centres or nerves. Such squints are common in meningitis and lesions of the mid-brain or cerebellum, such as tumours (glioma, tubercle, gumma, &c.). The occurrence of the squint only during epileptiform fits or its irregularity of type may render the diagnosis from paralytic squint easy, especially when there are other prominent symptoms of cerebral irritation. In other cases, especially in the early stages of the disease, the diagnosis from paralytic or concomitant squint may be extremely difficult.

Synkineses.

The extrinsic muscles take part in many normal and pathological synkineses. When the eyes look up the levatores palpebrarum raise the lids and in extreme upward movements the frontales also contract. In congenital ptosis (*q.v.*) upward movement of the eyes is often defective. On looking down the lid follows the globe. In exophthalmic goître the lid follows tardily or not at all (von Graefe's sign); in total facial paralysis the lid follows the globe on looking down, though the eye cannot be closed voluntarily. On closing the eyes, as in sleep, the eyes turn upwards and outwards. The same movement of the eyes occurs on attempted closure in total facial paralysis. On the other hand the eye sometimes closes in total seventh nerve paralysis on synergic activity of other facial muscles, as in laughing. The so-called "jaw-winking" synkinesis is particularly striking. In these rare cases one levator palpebræ is thrown into activity during eating, and sometimes on reading aloud. The lid movement is usually specially associated with lateral movements of the jaw, due to action of the pterygoid muscles, which are innervated by the fifth nerve. In most cases, but not all, there is slight ptosis of the affected lid, and in cases with congenital ptosis the synkinesis occurs on sucking. Allied to the jaw-winking cases are others in which spasmodic lid movements occur on lateral deviation of the eyes. The convergence pupillary synkinesis has already been mentioned: to it may be added the contraction of the pupil on forced closure of the lids. In rare cases spontaneous rhythmical variations in the size of the pupil are accompanied by ocular or lid movements. They are usually associated with congenital or early infantile paresis of the third nerve. The pupil contracts rapidly to about 2 mm. diameter, then after 5—10 seconds dilates slowly to 6—7 mm., contracting again after 15—20 seconds. Contraction is hastened by action of the internal rectus, dilatation by action of the external rectus. The movements are accompanied by spasm and relaxation of the ciliary muscle.

Nystagmus.

Nystagmus (*νυστάζειν*, to nod) is the term applied to rapid oscillatory movements of the eyes, independent of the normal movements, which are not affected. The oscillations are involuntary, though in rare cases normal persons can imitate them. They are usually lateral, but vertical, rotatory, and mixed rotatory and lateral or vertical nystagmus are not uncommon. The condition is almost always bilateral, though the movements may be much more marked in one eye than the other. In such cases it may be necessary to examine the eye very carefully with the ophthalmoscope (corneal reflex, retinal vessels, &c.) before the presence of nystagmus can be demonstrated. Unilateral nystagmus does occur, but it is probable that many of the cases described are really bilateral.

Nystagmoid jerks, *i.e.*, larger rhythmic jerking movements, most pronounced at the extreme limits of the normal movements of the eyes, should be distinguished from true nystagmus. They are not uncommon in normal people under certain conditions—fatigue, railway travelling, &c. The fundamental cause is probably quite different from that of true nystagmus, though both may occur together.

Nystagmus may be congenital or early infantile, or it may be acquired. These two groups of cases should also be carefully distinguished on account of their different pathological foundation. Congenital and early infantile nystagmus, *i.e.*, nystagmus dating from birth or within a few weeks of birth, occurs in congenitally malformed eyes, in albinism, and in eyes with congenital or early developed opacities of the media, *e.g.*, leucoma or anterior polar cataract due to ophthalmia neonatorum (*q.v.*), macular changes, &c. The cause in these cases is inability to develop normal fixation. Fixation is developed during the first few weeks of life, the eyes being moved aimlessly and independently before it is acquired. Any cause seriously diminishing the acuity of macular vision occurring at this period is liable to give rise to nystagmus ; if the eye is blind, nystagmus is not developed. Nystagmus is present in total colour blindness (*q.v.*), in which vision is carried out by the rods alone, and there is therefore a central scotoma. In some congenital cases it is impossible to discover such a cause. In a few such cases ancestors or relations have been albinos.

Nystagmus may be acquired in infancy after the period at which fixation is developed. This form occurs in *spasmus*

nutans, in which it is associated with nodding movements of the head. It occurs in the first year of life. The nodding of the head may be antero-posterior (affirmation), lateral (negation), or rotatory. It develops some weeks before the nystagmus, ceases during sleep, and disappears before the nystagmus. The whole symptom-complex disappears in time—one of the few cases in which nystagmus disappears spontaneously. The nystagmus may disappear in one eye before the other; such cases may be mistaken for true unilateral nystagmus. In rare cases head nodding with nystagmus is congenital and hereditary, and in these cases persists throughout life (Hancock).

Nystagmus in adults occurs in disseminated sclerosis, disease of the cerebellum and vestibular tracts, and of the semicircular canals (*e.g.*, occasionally on syringing the ears), Friedreich's ataxia, &c. In disseminated sclerosis the movements are generally horizontal and are elicited in the early stages only in extreme lateral positions of the eyes. Cerebellar irritative lesions cause coarse nystagmus towards the side of the lesion and fine nystagmus to the opposite side. Some of these cases show analogy with hippus (*vide* p. 57), and like it are probably dependent upon the rhythmic activity of nerve centres. Nystagmus may also occur in adults as an "occupation neurosis," the commonest form being coal-miner's nystagmus (*vide infra*).

In congenital and early infantile nystagmus the patient is wholly unconscious of the movements, since objects do not appear to move. Vision is usually defective in spite of correction of errors of refraction which generally accompany the defect. In some cases of acquired nystagmus in adults objects appear to move.

The prognosis is good in spasmus nutans and in miners' nystagmus if the occupation is changed, though recovery is slow. In all other cases it is bad, though it tends to diminish with advancing years. Treatment is therefore palliative, consisting in correction of refraction, wearing smoked glasses in albinism, and treating any disease which may be present.

Labyrinthine Nystagmus occurs in disease of the internal ear in which the semicircular canals are involved, and can be produced in normal subjects by rotation in a specially designed chair or by passing a galvanic current through the head. The nystagmus is rhythmic, with a rapid and a slow component, is bilateral, and horizontal or rotatory, but varies according to the semicircular canal stimulated. Either pair of semicircular canals can be

stimulated by rotation with the head in a suitable position. Destruction of one labyrinth causes rhythmic nystagmus towards the opposite side, which ceases if the other labyrinth is destroyed.

Miners' Nystagmus occurs chiefly in those who have worked long at the coal face. The patient complains of defective vision which is worse at night, headache, giddiness, photophobia, dancing of lights and movements of objects. The nystagmus is essentially rotatory and very rapid; in latent cases it is elicited by fixing the head and making the patient look up. In severe cases the lids are nearly closed and the head is held backwards; there is tremor of the head and eyebrows. The disease is six times as common in pits which use safety-lamps as compared with those that use naked lights, and varies inversely with the illumination (Llewellyn). Continual looking upward at work is only a contributory cause, other such causes being ill-health, accidents, errors of refraction, subnormal pigmentation, &c. Scotopia, or vision in a dull light, is carried out almost entirely by the rods. Under these circumstances visual acuity is greatest 10°— 15° outside the fovea, and there is a physiological central scotoma. There is great difficulty in keeping up fixation, and the evidence is strongly in favour of the view that this is the essential ætiological factor in miners' nystagmus. Improvement in miners' lamps and in the lighting of mines would probably eliminate the disease, which is a cause of enormous economic loss in compensations, &c.

CHAPTER XXVIII.

CONCOMITANT STRABISMUS. HETEROPHORIA. CONGENITAL DEFECTS.

Concomitant Strabismus.

In concomitant strabismus the visual axes, though abnormally directed, retain their abnormal relation to each other in all movements of the eyes. It differs therefore in this respect from paralytic strabismus, in which the relationship of the visual axes to each other changes with every movement of the eyes in the direction of action of the paralysed muscle or in the resultant direction of action of the paralysed muscles when more than one is affected. The secondary deviation in concomitant squint is equal to the primary deviation, a fact which demonstrates the absence of paralysis (Figs. 278—9). The deviation of the visual axes may be convergent or divergent, the former being the more common.



Fig. 278.—Diagram of primary deviation in concomitant convergent strabismus.

In every case in which the direction of the visual axes is aberrant paralytic strabismus must first be eliminated by testing the movements of the eyes in all directions with the finger. If they are found to be normal and there is no complaint of diplopia, it may be concluded that there is no paralysis. It does not follow that there is concomitant squint. The convergence or divergence of the axes may be only apparent. A marked appearance of convergent squint is sometimes seen in myopic eyes, of divergent squint in hyper-

metropic eyes. It will be observed later that true concomitant convergent squint is most commonly associated with hypermetropia, divergent with myopia, *i.e.*, the opposite of apparent strabismus.

Apparent strabismus is due to the fact that the visual axis of the eye is very rarely coincident with the optic axis (Fig. 280). The optic axis, *i.e.*, the axis upon which the cornea and lens are centred, passes through the centre of rotation of the eye and approximately through the centre of the pupil. The visual axis passes through the nodal point and the fovea centralis, thus crossing the optic axis and making a small angle with it. This angle is very nearly equal to an angle which is called the angle gamma γ; it is commonly spoken of clinically as the angle γ. In the emmetropic eye the angle γ is said to be positive, *i.e.*, the optic axis cuts the retina internal to the fovea centralis. In hypermetropic eyes the angle γ is also positive but greater than in emmetropia. In myopia the angle γ is absent or negative, *i.e.*, the visual axis and the optic axis coincide or the latter cuts the retina external to the fovea centralis.



FIG. 279.—Diagram of secondary deviation in concomitant convergent strabismus.

Now, neither of these lines can be seen, and the direction of the line of vision is judged by the position of the pupil. Hence the greater the size of a positive angle γ the more the eye will appear to look outwards. If the angle γ is negative the eye will appear to look inwards. Therefore in high hypermetropia there will be an apparent divergent squint, in high myopia an apparent convergent squint. The latter is the more striking because the emmetropic eye usually has a positive angle γ of 5°, thus producing an apparent divergence of 10°, which, however, we are accustomed to regard as the normal position of the eyes.

Having decided that the case is not one of paralytic strabismus, it is necessary next to show that it is real, not merely

apparent. This is easily done as follows: The patient is told to fix the surgeon's index finger, which is held up at least two feet from the eyes. If it is held closer, as is too often done, normal convergence will vitiate the result. The surgeon's left hand or a screen is held in front of the patient's right eye; the left eye will now be accurately fixing the finger. The screen is then moved so as to cover the left eye, fixation being now taken up by the right eye. If the right eye moves inwards or outwards at the moment when it takes up fixation there is a true squint; if it remains absolutely motionless the squint is apparent only.

Having now eliminated both paralytic and apparent strabismus, it is almost certain that the case is one of true concomitant squint (see, however, p. 529). In concomitant squint one eye maintains fixation while the other is adverted or abverted. If in the test just described the fixing eye is covered by the screen, the deviating eye generally moves outwards or inwards through an angle equivalent to the angle of the deviation in order to take up fixation. At the same time the eye behind the screen moves inwards or outwards through exactly the same angle (secondary deviation). If now the screen is removed it will generally be found that fixation immediately reverts to the first eye. Such a case is usually termed one of *unilateral strabismus*. In some cases fixation is still retained by the second eye. If it is so retained for a considerable period, *e.g.*, while the patient blinks several times, the squint is said to be *alternating*. Inquiry will then generally elicit the fact that the patient sometimes squints with one eye, sometimes with the other: indeed, this may be noticed while watching the patient.



FIG. 280.—Apparent strabismus. *a b*, optic axis upon which the refractive surfaces are centred; *f*, fovea centralis; *n*, nodal point; *c*, centre of rotation; O, point of fixation; O *c*, line of fixation; O *n f*, line of vision; O *c b*, angle γ. It is practically equal to O *n b*, which can be measured. In actual practice the guide to *a b* is taken from the centre of the pupil; *a b* does not usually pass accurately through the centre of the pupil, so that the result is always only approximate. The angle γ is to the nasal side in hypermetropia and emmetropia.

Occasionally patients with alternating strabismus can fix with either eye voluntarily, but usually they are unconscious which eye is fixing. Concomitant squint may be *constant*, or occur only at intervals—*periodic*.

It has been mentioned that when the fixing eye is covered with the screen the deviating eye usually moves so as to take up fixation. In unilateral squints of long-standing this eye may remain motionless and can only be moved into the primary position by moving the finger, a condition which is called *eccentric fixation* (*vide* p. 521). Since it occurs only with marked deviation of long standing there is generally no difficulty in distinguishing it from apparent squint.

In performing the preliminary test to eliminate paralytic strabismus it will often be found that in true concomitant squint with considerable deviation the eyes do not move as much as usual in the direction opposite to that of the deviation. Thus, in convergent squint it may be very difficult to get the eyes to move outwards as much as normal, so that the margin of the cornea lies under the external canthus. Similarly in divergent squint it may be very difficult to get the eyes to move inwards as much as normal, so that the margin of the cornea is well covered by the internal canthus. This defective movement is commonly attributed to "insufficiency" of the external or internal recti respectively. In convergent strabismus it is probably due, not to any defect in the external rectus or its innervation, but to the fact that, fixation being dependent upon one eye, there is little stimulus to outward movement as soon as the point of fixation has passed beyond the field of fixation of this eye, *i.e.*, as soon as the nose cuts off vision of the finger. In many such cases the eye will move out completely if the finger is moved rapidly. In divergent strabismus defective inward movement is sometimes due to mechanical causes, viz., the size of the myopic eye (*vide* p. 529).

It has already been mentioned that in concomitant strabismus there is no diplopia. It may be present in the earliest stages, but is invariably absent in the later. This is due to psychological suppression of the image of the squinting eye. In most cases suppression is aided by actual defect, usually ametropia, in this eye, but it is not the complete explanation, since suppression of the image of the squinting eye is also the rule in alternating squint, in which both eyes are frequently quite normal or have the same degree of ametropia. Suppression is doubtless aided in all cases by the peripheral situation of the image in the squinting eye, but there is no doubt that the

seat of suppression is really in the brain : that is, in the interpretation of the stimuli reaching the brain from the eyes those derived from the squinting eye are deliberately neglected. The ability to exclude the impulses derived from the squinting eye is an important point in attempting to arrive at the rationale of concomitant strabismus. It follows from this fact that people with convergent squint have only uniocular vision, or, at most, very imperfect binocular vision.

Except in alternating strabismus the vision of the squinting eye is nearly always very defective, which is partly due in most cases to errors of refraction. In convergent strabismus the eyes are nearly always hypermetropic, with or without astigmatism, and often the squinting eye has greater ametropia. There can be little doubt that some such inherent defect may determine which eye will deviate, though it is probably never the fundamental cause of the squint.

In some cases the vision in the squinting eye is defective beyond any explanation derived from objective defects — ametropia, &c. In some such cases there is reason to believe that the defective vision dates from birth—*congenital amblyopia*. In all unilateral squints of long standing, very defective vision in the squinting eye is the rule. It is commonly attributed to the prolonged suppression of the images derived from this eye, and is hence called *amblyopia ex anopsiâ*. This explanation is not altogether satisfactory, since cases are well known in which vision has long been excluded by congenital cataract, yet is quite good after successful operation. The vision in the amblyopic eye is often reduced to 6/60, and may be reduced to counting fingers. Cases of recovery of sight after loss of the fixing eye have been recorded, but unfortunately this result certainly does not invariably follow. The visual acuity may be greater in the false position than when the retinal image falls upon the fovea (eccentric fixation with " false macula "). All power of fixation may be lost by the amblyopic eye.

It has already been mentioned that the gross movements of the eyes are nearly or quite perfect in concomitant strabismus. This applies equally to dynamic convergence and accommodation. The eyes start in an abnormal position, and normal movements are superposed. Thus, in fixing a new object, the normal amount of dynamic convergence is superposed upon the abnormal static convergence or divergence.

Concomitant strabismus always commences in childhood, generally in infancy. It may become manifest after a fright, an attack of whooping cough, measles or other debilitating

illness, and is often popularly attributed to some such cause. It must be carefully distinguished from the squinting which normally occurs during the first few weeks of life, before fixation is developed (*vide* p. 514); this is not concomitant. Many important factors in the ætiology of concomitant strabismus are known, and a proper appreciation of them is essential to rational treatment. No theory of the fundamental causation which has yet been advanced satisfactorily explains the condition.

Attention was early drawn by Donders to the common association of convergent strabismus with hypermetropia. He explained the relationship by the fact of the normal association of convergence and accommodation (*vide* p. 499). Hypermetropes have to exercise an effort of accommodation to see distant objects, still more to see near objects. The effort of accommodation is associated in the normal person with a corresponding effort of convergence. If this rule be supposed to hold good for hypermetropes, the strong accommodation may be regarded as inciting an effort of convergence which is excessive for the actual point of fixation. The hypermetrope is therefore in a dilemma. He must either converge accurately for the object, in which case he will not accommodate sufficiently to see it clearly: or he must accommodate accurately for it, in which case he will converge too much. This will cause homonymous diplopia, unless he is able to suppress the image of one eye, which is exactly what the patient with convergent strabismus does.

Regarding divergence as negative convergence, the association of divergent strabismus with myopia is explained by the same theory.

There is no question that this factor is one of great importance, as is shown by the cure of some cases of squint by suitable correcting glasses. It is not, however, the fundamental cause of squint, for if it were (1) all uncorrected hypermetropes would have to squint; (2) there would be ametropia in all cases of concomitant strabismus. The latter corollary is found to be false, for convergent strabismus, other than apparent strabismus, is occasionally found to be associated with myopia, and in alternating strabismus there is often little or no ametropia.

In spite of these facts the great importance of the association between accommodation and fixation must be strongly insisted upon. Convergent strabismus most frequently develops between the ages of two and six, *i.e.*, just at the period when

the fixation of near objects throws a strain upon accommodation. It is often periodic at this stage, and noticed only when near objects are looked at. Moreover, there is an undoubted tendency for the deviation in all cases of convergent strabismus to diminish with age, *i.e.*, with the diminution of accommodation. The relative infrequency of convergent squint in adults, compared with its frequency in children, cannot be explained solely by the fact that many cases undergo successful treatment. It may be remarked that the amblyopia persists in the formerly squinting eye; hence in every case in which satisfactory objective evidence of the cause of defective vision in one eye cannot be discovered, the patient should be asked if he ever squinted.

As already mentioned, greater ametropia in one eye, opacities in the refracting media, introcular disease, and so on, are never the fundamental cause of strabismus, though they may determine the particular eye which loses fixation. This is especially the case when concomitant squint is preceded by latent squint (*q.v.*). It is easy to understand that when there is disturbance of muscular equilibrium which can only be overcome by special effort, any slight defect in one eye may determine the development of a manifest squint.

The application of Donders' theory to divergent strabismus associated with myopia has been mentioned. Here, since near objects are seen with little or no accommodation, the impulse to convergence is too weak. Since infants are rarely myopic, this form of divergent squint does not develop in early childhood. There are other factors besides deficient accommodation which tend to produce divergence in myopia. One is the mechanical conditions of the myopic eyeball, which, being abnormally large and long, adapts itself to the axis of the orbit. Further, the internal recti act under mechanical disadvantage from the same cause. Moreover, in very high myopia the far point of the eye is so close to it that it is impossible for convergence to be effectual—it becomes impossible to see the object with both eyes at the same time. The better eye is then used and the other is allowed to take up the position of rest, which is usually one of divergence. Such a strabismus may remain periodic for near work only for many years; in other cases it becomes constant.

Spontaneous cure rarely if ever occurs in divergent strabismus, which tends to increase with age.

Concomitant strabismus has proved a fertile field for conjecture. There are several points bearing upon the subject

which are obscure, such as the development of fixation and of binocular vision, the occurrence of congenital amblyopia, and of amblyopia ex anopsiâ, and so on. Many theories relating to these points have been stated and restated so frequently that they are accepted as facts. In reality, many are not susceptible of demonstration, and none have been proved. The theorist on strabismus accepts the "facts" which fit his theory, and rejects the remainder.

The prevailing theory at the present time is that strabismus is due to defect of the fusion faculty, or the capacity of combining psychologically the impressions derived from the two eyes. Some authors go so far as to hypothecate a "fusion centre" in the brain. No one will deny that the fusion of the images derived from the two eyes in binocular vision is an act in consciousness, and that it has a physical basis. This physical basis is a set of accurately co-ordinated nerve impulses. Binocular vision, then, depends upon the accuracy of co-ordination of these impulses, and this is a function of the nerve-complex as a whole, not of any particular "centre." It is indeed possible that the strabismus is caused by inco-ordination of the afferent impulses upon which binocular vision depends, though this advances the true ætiology of the disease but little. It is just as probable that the inco-ordination or deficiency of the afferent impulses is a result of the strabismus, which is itself due to some other cause; or both may be due to a common cause.



FIG. 281.—Diagram of the position of the corneal reflex as a guide to the angle of the squint.

In every case of concomitant strabismus the angle of the deviation should be measured, so that the effects of treatment may be accurately observed. A rough indication of the angle of the squint can be obtained from the position of the corneal reflex when light is thrown into the eye with the ophthalmoscopic mirror (Fig. 281). The light is thrown in from a distance of about two feet, and the patient is told to look at the mirror; an infant does this reflexly. In the fixing eye the corneal reflex will be in the centre of the pupil, or slightly to the inner side if there is a large angle γ, to the outer side if there is a negative angle γ. The light is then turned on to the squinting eye. If the reflex is about half-way between the centre of the pupil and the corneal margin, there

is a deviation of about 20° ; if it is at the corneal margin, about 45°. This test is only a rough one.

The best test in children is with Priestley Smith's tape (Fig. 282). It consists of a tape 1 metre or 60 cm. long, with a ring at each end. To one ring a second tape is attached, having a tangent scale upon it. The graduations are, of course, different, according to the length of the first tape. At the other end of the tangent tape is a small weight. The measurement is carried out in the dark room. The free ring is held by the patient or an assistant on the cheek immediately below the fixing eye. The surgeon passes one finger of the hand, which also holds his ophthalmoscope, through the other ring, and keeps the tape taut. With his disengaged hand he holds the tangent tape at right angles to the distance tape, at the same time holding up the index finger as an object of fixation. The light from the ophthalmoscope is thrown into the squinting eye, and the patient is told to look at the fixation finger. This is moved along the tangent tape until the corneal reflex is in the centre of the pupil. The angle of the squint is then read off on the tape. The direction in which the tangent tape is held, whether to the surgeon's right or left, depends, of course, upon the side of the squinting eye and the nature of the squint, whether convergent or divergent. This test is by far the best of any yet devised for small children.



FIG. 282.—Diagram of Priestley Smith's tape for measuring the angle of squint. R, right eye with internal squint; L, left eye; O, ophthalmoscope; O F, graduated tape; F, fixation point (observer's finger); W, weight. The angle measured is O L F, which is equal to O R R′, the angle of the squint.

The angle of deviation can also be measured with the perimeter (Fig. 283). The squinting eye is placed at the centre

of the arc, and the patient fixes an object *six metres away*, situated slightly over the position of the ordinary fixation spot of the perimeter. The arc of the perimeter is turned to the horizontal position on the side towards which the squinting eye is directed. The surgeon passes a candle flame along the arc until the corneal reflex is in the centre of the cornea of the squinting eye. He then reads off the angle of the squint on the arc. If extreme accuracy is desired, the angle γ should be measured and allowed for. It is done by covering the sound eye and making the patient fix the fixation spot of the perimeter with the squinting eye. The candle flame is again carried along the arc until the corneal reflex is again in the centre of the cornea. The angle γ is read off on the arc. The method is not suitable for children, and can only be employed with intelligent adults. One objection to it is the tendency to fix the ordinary fixation spot of the perimeter instead of a distant spot in the same line of vision. If this mistake is made errors arise, more particularly because the amount of dynamic convergence exerted in these cases is not always the same as with normal eyes.



FIG. 283.—Measurement of the angle of squint by the perimeter.

The angle of deviation can also be conveniently measured on a tangent scale set against the wall, the corneal reflex of a candle flame being again used as a guide.

Treatment.—The routine treatment of a case of concomitant convergent strabismus in a child is as follows :—

(1) Record the distant vision of each eye if the child is not too young, the angle of the deviation, &c. Order ung. atropini, 1 per cent., three times a day for at least four days. At the end of this period estimate the error of refraction by retinoscopy and confirm the result subjectively if possible ; reliance should be placed on the retinoscopy rather than on subjective tests. Again measure the angle of the squint, which

is likely to be less under atropin than without a mydriatic. Order the full correction for constant use. A smaller correction for the effect of atropin should be made than in hypermetropia without squint. If the error is considerable I usually subtract only 0·5 D for atropin instead of 1 D; if the error is small I order the full atropin correction to start with. Great care must be taken to correct all astigmatism, especially in the squinting eye. The patient is told to return in a month.

If the child is less than two years old I do not order glasses except in rare cases. Some surgeons order them in all cases. I prefer to eliminate accommodation by keeping both eyes under the influence of atropin; the 1 per cent. ointment need be applied only once a day. The child should be examined at regular intervals until it is considered advisable to order glasses.

(2) After the glasses have been used constantly for a month the child is again examined. The vision is tested and the angle of the squint again measured. The treatment now depends upon the condition of the vision in the squinting eye. If, as is usually the case, this eye is amblyopic, an effort should be made to improve the vision in it by continual exercise. In order that this eye may be used the other must be prevented from seeing, or at any rate from seeing clearly. It may be done by placing a pad of cotton wool under the glass in front of the fixing eye for a certain period, *e.g.*, ½—1 hour, three or four times a day. This method is liable to be neglected or carried out inefficiently. A better method is to order the instillation of atropin *into the fixing eye only*, once a day. This eye will then generally be used only for distant vision, the squinting eye being used for seeing near objects. The child is examined at intervals of a month or two, and any improvement in vision in the squinting eye, as well as any change in the angle of deviation, carefully recorded. The treatment requires watching, as in some cases the deviation becomes transferred to the atropised eye, and the vision in this eye may deteriorate.

(3) The further treatment depends upon the size of the angle of deviation, the condition of vision in the squinting eye, and a variety of other factors which differ in each case. If there is any evidence of some degree of binocular vision it may be advisable to attempt to cultivate it by *orthoptic treatment*. This consists essentially in specially devised stereoscopic exercises. It is excessively tedious and seldom successful. It has, however, one overwhelming argument in

its favour, viz., that when successful it cures the squint. The cure is complete, *i.e.*, the patient is placed in the same condition as a normal person; his eyes are straight and he has binocular vision. No other treatment can be said to cure the disorder. The eyes can be put straight, but this cures only the deviation; the other elements of the disorder remain unaffected. Unfortunately orthoptic treatment succeeds only in very favourable cases, and then only after prolonged and very persevering efforts. In the majority of cases it is useless to attempt it, and in all cases it is useless unless carried out systematically and thoroughly. For the details of the treatment monographs on the subject must be consulted.

Orthoptic treatment may demand operative procedure at this stage or even earlier. If orthoptic treatment is considered hopeless or has been abandoned the further treatment consists in continuance of that described under (1) and (2), but the question of the advisability of operative interference has now to be considered,

(4) *Operation.*—Surgeons differ so much in the principles which influence them in advising operation and in the nature of the operation employed that it will be best to describe the treatment which I adopt myself.

If the deviation is 10° or less tenotomy of the internal rectus of the squinting eye should be performed, under a general anæsthetic if necessary. This will cure the deviation or reduce it to a negligible quantity.

If the deviation is more than 10°, advancement of the external rectus of the squinting eye, usually with tenotomy of the internal rectus of the same eye, will be necessary. This operation should be performed under a general anæsthetic only in very rare cases—indeed only in cases in which orthoptic treatment is indicated. The operation should therefore be postponed until the child is old enough to be operated upon under cocain. The age varies with the individual; some quite young children, *e.g.*, under ten, submit well if judiciously managed. The operation is quite painless, except when tension is put upon the muscle. It is almost impossible to avoid slight tension during certain stages, but it need only be momentary. The objection to operating under a general anæsthetic is that the position of the eyes varies so much in different stages of anæsthesia that it gives no criterion of the final position after the anæsthetic has passed off. Very good results can, however, be obtained by an experienced operator if he keeps firmly in mind the amount of the deviation

and ignores the position actually present under the anæsthetic.

The treatment of alternating concomitant convergent squint without appreciable error of refraction is purely cosmetic. These patients have no binocular vision, and it is useless to attempt to develop it unless the case is seen when the patient is very young, or immediately after the squint has been first noticed. Usually there is considerable deviation, so that an advancement operation is required. It should be postponed until a perfect result can be guaranteed, *i.e.*, until the operation can be performed with local anæsthesia.

The treatment of concomitant divergent strabismus is similar to that of the convergent type. The refraction must be first carefully corrected, and it is advisable to order a full correction for constant use unless the myopia is very high. Tenotomy of the external rectus is seldom indicated in these cases because the benefit derived is too slight ; an external tenotomy will not correct much more than 5° deviation. Advancement of the internal rectus should be left in most cases until it can be performed under cocain. No operation is advisable in very high grades of myopia, since the size of the eye may render success mechanically impossible (*vide infra*).

In divergent strabismus over-correction is indicated, for these eyes show a great tendency to revert to their former position. In convergent strabismus the deviation should be fully corrected only in adults, on account of the tendency of the deviation to diminish pari passu with the loss of accommodation (*vide* p. 523). Moreover, slight convergence is less unsightly than divergence.

Divergent Strabismus other than Concomitant and Paralytic.—There are cases of divergent squint which are not paralytic, nor are they strictly concomitant. Some are very nearly allied to the true myopic divergent squint and may arise from it, *e.g.*, the divergence of the worse eye after binocular fixation has been completely abandoned. Similarly the mechanical divergence of extreme myopia is neither concomitant nor paralytic.

Another form of divergence, allied to the unilateral divergence in unequal myopia, is met with occasionally in any form of anisometropia in which the difference in refraction between the two eyes is great. In these cases, which are seldom susceptible of full correction (*vide* p. 486), one eye only is used for fixation. It is not surprising, therefore, that the unused eye assumes a position of rest, which is usually one of divergence. It is more surprising that relatively few such cases of aniso-

metropia develop a manifest squint. When a squint is developed the patient not infrequently complains of diplopia, which is the more trying the less the divergence. These cases are difficult to treat. Prolonged use of the fullest possible correction, aided by stereoscopic exercises, should be tried first. If it fails tenotomy of the external rectus of the diverging eye may effect a cure in slight deviations. In other cases an advancement is indicated ; in such cases extreme delicacy of manipulation is required. Operation undertaken for cosmetic reasons may sometimes induce or increase diplopia by approximating the true and false images ; it is usually permanent in these cases and exceedingly distressing.

An eye may diverge as the result of over-effect from an internal tenotomy. Such cases were at one time not uncommon, owing to an attempt being made to obtain a greater effect than can be reasonably expected from the operation. This is only attainable by liberating the muscle very freely from Tenon's capsule ; it is then prone to slip back so that it fails to become re-attached to the globe. The squint therefore resembles completely a paralytic squint. The muscle will usually be found imbedded in the tissues lying under the caruncle. When the attempt is made to move the eye inwards the caruncle is seen to be actively retracted. An advancement operation should be performed, but perfect movement is seldom regained.

Finally, a blind eye diverges sooner or later.

Latent Strabismus or Heterophoria.

It is found in some apparently normal persons that in the screen test (*vide* p. 503), when the screen is removed from before one eye, that eye moves slightly inwards or outwards to regain binocular fixation ; if the screen is placed in front of the other eye and then removed, this eye also moves slightly inwards or outwards respectively to regain fixation. When both eyes are fixing there is no deviation. Such a squint is called a latent squint or, in opposition to the normal condition of orthophoria, heterophoria. If the latent squint is one of convergence the condition is called esophoria, of divergence exophoria. Sometimes one eye is higher than the other ; this condition is usually called hyperphoria : as a matter of fact it is impossible in these cases to be sure whether there is absolute hyperphoria of one eye or hypophoria of the other, the condition being relative only.

It must be concluded that when the eyes are screened they take up a position of rest, the extrinsic muscles exercising merely their normal tone. In cases of latent squint the position of rest is not orthophoria, with the visual axes parallel, but heterophoria, with some deviation of the axes. During normal vision the requirements of binocular vision demand a suitable readjustment of the visual axes, which can be brought about only by tonic contraction of certain muscles—in esophoria of both external recti, in exophoria of both internal recti. This involves a perpetual strain, which often manifests itself as asthenopia. As might be expected the deviation is liable to become manifest in conditions of bodily fatigue. Some periodic squints are due to this cause, and the periodicity may be very rhythmic. Thus a child may squint in the evening when he is tired ; after a good night's rest the squint has disappeared, and may not return until the second or third day, the sequence being accurately repeated. Often latent squints give no trouble until school time arrives or adult life is reached. Here the demands of near vision increase the strain. No symptoms arise perhaps until after reading or writing for an hour or two. Then " the letters seem to run together." This is due to relaxation of the overstrained muscles ; the eyes momentarily assume the position of rest, and diplopia, which is not realised as actual double vision, causes blurring of the print. With an effort the blurring is overcome, but eventually this becomes impossible, headache supervenes, and the work has to be abandoned.

Analysis of the cases shows that slight eso- and exo-phoria are quite common and give rise to little or no trouble, which is not difficult to understand when it is remembered that over-action of both internal recti is physiological in ordinary convergence on near objects. These muscles are therefore accustomed to act together and little strain is felt. The same is true in less degree of the external recti. Only when the deviation is great —5°—10° or more—is asthenopia frequently present. Very slight degrees of *hyperphoria*, however, almost invariably cause extreme discomfort, for in these cases over-action of muscles which are not accustomed to work together is necessary in order to keep the visual axes in the same plane. For instance, in the primary position of the eyes there must be over-action of one superior rectus and inferior oblique, combined with over-action of the other inferior rectus and superior oblique, and the readjustment in other positions of the eyes must be very complex.

It may be impossible to discover slight degrees of heterophoria by the screen test. More delicate tests have therefore been devised. All depend upon disassociating the two eyes. The simplest method is that of so altering the appearance of the retinal image in one eye that it affords no stimulus to fusion with the image of the other eye.

The patient is placed six metres from a candle or bright spot of light in a dark room. A Maddox rod (Fig. 284), which consists of four or five cylinders of red glass side by side in a brass disc, is placed in the trial frame before one eye. The spot of light seen through the red cylinders appears as a long red line. If the cylinders are placed with their axes horizontal the red line will be vertical. If there is orthophoria the bright spot will appear to be in the centre of the vertical red line; if there is eso- or exo-phoria the red line will be to one side of the spot. The angle of the deviation is measured by the strength of the prism which it is necessary to place in front of the Maddox rod in order to bring the red line and the spot together. The nature of the deviation is indicated by the position of the base of the prism, whether out or in.



Fig. 284.—Maddox rod.

The Maddox rod is then turned round so that the cylinders are vertical; the red line will now be horizontal. If there is no hyperphoria the line will pass through the bright spot. If there is hyperphoria the red line will be below or above the spot according as the relative hyperphoria is in the eye with the rod in front of it or in the other. The amount of deviation is measured by the strength of the prism required to correct it.

The deviation in latent squint is often different in near vision from that in distant, so that both must be tested. The deviation in near vision is tested by means of a special card (Plate XXI.). A strong prism—about 12°—is placed base down or up before one eye. In orthophoria the arrows are exactly in the same vertical line. In eso- or exo-phoria the lower arrow points to a number in the upper scale. This number gives the angle of deviation, which may be confirmed by counteracting the deviation with a prism, base in or out, before one eye.

Goldfinches may often be seen in a flock, their bright yellow wings flashing in the sunshine, as they dart from prickly plant to plant, doing good service.

Maddox's scale to measure latent insufficiency or excess of convergence at the distance of 25 cm. with a prism of 12° (*i.e.*, 6° deviation or 6° d) base upwards before the R. eye. The figure to which the lowest arrow points indicates in degrees the amount of latent convergence (black letters) or divergence (red letters) which may be present. The figures under the waved line are—teens (ten, twelve, thirteen, &c.) and the capital letters, A, B, C, D, represent metre angles for an interocular distance of 64 mm.; if the lower arrow point to black A there is 1 metre angle (1 m.a. or 1 M) of convergence, if to black B, 2 m.a., &c. The printed matter under the numbers is introduced to ensure accurate accommodation for the distance (25 cm.).

[*To face p.* 532.

Besides the actual measurement of the deviation in latent strabismus the strength of the muscles involved should also be tested by forcing them to a maximum effort against prisms. With the patient seated six metres from a candle the highest prism, base down before the right eye, still permitting of single vision, gives the range of superduction of that eye. Subduction and abduction can be measured in the same manner. Adduction gives less concordant results. The normal limits of super- and sub-duction are 1·5° to 2·5°; of abduction 4° to 5°.

Allied to these defects, though not strictly speaking a latent squint, is insufficiency or weakness of convergence. It will be revealed by the ordinary tests of motor balance. If there is more exophoria or less esophoria in near vision than in distant there is insufficiency of convergence. The defect causes asthenopia in near work.

Treatment.—The lower degrees of esophoria, and to a less extent of exophoria, cause no symptoms and need no special treatment. Slight exophoria often causes symptoms in young adults much engaged in near work. It is relieved by suitable prisms, bases in; if the general health improves, or the amount of near work is diminished, the prisms can be dispensed with later. Hyperphoria is most likely to cause asthenopic symptoms. It is corrected by ordering suitable prisms to be combined with the glasses which correct any refractive error. If the spherical error is sufficiently great the prismatic effect may be obtained by decentring the lenses. The total prismatic error should be divided equally between the two eyes in ordering the correction. Thus, if there is hyperphoria of 3° as measured by a 3° prism, base down before the left eye, a prism of 1½° is ordered before each eye, base up for the right, base down for the left. When this treatment does not succeed, and the deviation is considerable, tenotomy of the superior rectus may be necessary, but such cases are rare, and operative interference should not be lightly undertaken, for it is apt to be disappointing.

The rational treatment of large degrees of eso- or exo-phoria consists in exercising the weak muscles against prisms. This is usually only temporarily beneficial, but the muscles can be kept in good order by repeating the exercises at intervals. The asthenopia can be relieved by ordering prisms to correct the defect, *i.e.*, prisms with their bases directed in the opposite sense to those used for exercise. This should be avoided except in such cases as those already mentioned, since it generally tends to increase the defect, so that stronger prisms

have to be ordered from time to time. Stereoscopic exercises are beneficial. In severe cases operative interference—tenotomy—may be indicated.

Insufficiency of convergence may be treated by prism exercises. The following simple exercise is often sufficient without having recourse to prisms. Any error of refraction is corrected with glasses which are ordered to be used constantly. While reading the patient gradually brings the book nearer and nearer, until the print becomes blurred. He then slowly moves the book back to ordinary reading distance. The process is repeated. At about every tenth line the patient looks into the distance, so as to relax his accommodation and convergence. Two or three pages should be read in this manner three or four times a day for several weeks. The course is repeated as often as necessary.

If convergence training fails prisms, base in, must be ordered with the reading glasses. Care must be taken not to overcorrect presbyopia (*vide* p. 489).

Operations on the Extrinsic Muscles.

Tenotomy.—Instruments required: speculum, fixation forceps, strabismus hook, tenotomy scissors. I prefer a flat tenotomy hook (Moorfields' pattern) to the ordinary hook with a bulbous end. Local anæsthesia suffices in all but very young patients; a few drops of adrenalin (1 in 1,000) may be instilled with advantage. I usually stand above the patient's head in all operations on the extrinsic muscles.

The conjunctival sac having been douched, and the speculum inserted, the conjunctiva is seized over the muscle and a vertical incision, 8–9 mm. long, is made with scissors. It must be remembered that the internal rectus is inserted nearest to, the external farther from, the corneal margin (*vide* p. 492). The tissue underlying the conjunctiva at the lower border of the muscle is then seized with the forceps and incised, care being taken to hold the scissors so that they are tangential to the globe. If this is properly done Tenon's capsule is opened. The hook is now taken in the right hand, which still retains the scissors, and the point is introduced into the opening in Tenon's capsule. It is passed backwards, then upwards between the muscle and the sclerotic. In this manœuvre the point of the hook must be kept firmly against the globe. The point of the hook appears at the upper border of the muscle. It is freed from any conjunctival or subconjunctival tissue

which may cover it. The hook is drawn forwards until it lies close under the insertion of the muscle, but undue traction must not be exerted, as it causes severe pain. The hook is transferred to the left hand. The point of one blade of the scissors is introduced along the hook below the tendon, and the muscle is divided between the hook and the sclerotic. The hook should be introduced again and moved about in such a manner that any attachments which remain may be caught up and divided.

A pad and bandage are worn for two days, the eye being freely irrigated with boric acid lotion.

With the exception of a small puncture in the conjunctiva the whole operation can be performed subconjunctivally by an expert. This method has the advantage of preventing so much retraction of the caruncle in tenotomising the internal rectus as usually follows tenotomy by the open method. It is less easy and slightly more dangerous. The retraction of the caruncle in the open method can be counteracted by inserting a conjunctival suture (silk, No. 0), which however has the disadvantage that it must be removed in a few days.

No attempt should ever be made to obtain a greater effect from tenotomy of the internal rectus than 10°, of the external rectus 5°. If a greater effect is desired an advancement of the opponent must be performed.



Fig. 285.—Prince's forceps.

Advancement. — Instruments required: speculum, fixation forceps, tenotomy scissors, strabismus hook, Prince's advancement forceps (Fig. 285), needle holder (Fig. 286), curved needles (No. 2) (Fig. 287). Prince's advancement forceps have one smooth blade and one armed with sharp spikes which pass through holes in the smooth blade when the blades are locked. These forceps are usually made in right and left pairs. There is no necessity for this: it need simply be remembered that the smooth blade must be passed below the muscle. The best needle holder is the one figured; it has narrow jaws, so that the sharply-curved needles are not broken by their pressure.

Apart from exceptional circumstances (*vide* p. 528), advancement should always be performed under local anæsthesia. A few drops of adrenalin (1 in 1,000) should be instilled just before commencing the operation.



Fig. 286. — Silcock's needle holder.

The conjunctival sac having been douched, a mask of sterile gauze is placed over the face (Fig. 288). A hole is cut in it, opposite the eye to be operated upon. The speculum is then inserted. The patient is told to look in the required direction, *e.g.*, in operating upon the right external rectus towards the left. A vertical incision is made in the conjunctiva over the insertion of the muscle, as in tenotomy. A rectangular flap of conjunctiva is then dissected up, parallel horizontal incisions being made from the ends of the vertical incision. Tenon's capsule is then incised at the lower border of the muscle and the hook introduced as in tenotomy. The point of the hook is freed from subconjunctival tissue in the same manner. The smooth blade of Prince's forceps is passed along the hook, which is then removed. The forceps are closed, and locked by the spring catch. They are handed over to the care of the assistant. The patient is then told to look in the opposite direction, the eye being kept steady by the assistant with the advancement forceps. The internal rectus is then tenotomised.



Fig. 287. — Curved needles for advancement. Size No. 2 should be used.

The external rectus is now tenotomised by cutting with the scissors between the Prince's forceps and the sclerotic. The muscle is thoroughly freed from the sclerotic and held vertical by the assistant, without undue tension.

The next stage consists in inserting the stitches (silk, No. 1). If the right external rectus is being advanced the stitches are passed from the muscle towards the cornea ; if the left, in the reverse direction. We will suppose for the sake of description that the left external rectus is being operated upon. The needle is grasped by the holder about half-way between the eye and

the middle of the bend. It is passed into the episcleral tissue about 3 mm. from the corneo-scleral margin. It must be directed almost perpendicularly to the globe at first, so that a firm scleral grip is obtained. This is the most important and most difficult step in the operation. The needle is brought out about 1 mm. from its point of entry. In doing so pressure must be exerted in the direction of the curve of the needle ; if the pressure is exerted in the direction of a chord of the arc of the circle of curvature of the needle the latter will inevitably break. The middle stitch is inserted first, *i.e.*, the needle is introduced in the prolongation of the horizontal meridian of the cornea. Having obtained a firm anterior hold the needle is again grasped by the holder, this time nearer its eye. The



Fig. 288.—Diagram of advancement operation.

muscle is held vertical and placed somewhat on the stretch. The needle is inserted on the posterior surface, as far back as possible. It is passed through the muscle and conjunctiva and the thread is drawn through.

The other two stitches are inserted in the same manner, one above, the other below the first. Instead of bringing them through the muscle and conjunctiva as before, they should be passed through the muscle first, then through the anterior corners of the conjunctival flap. The sutures are kept carefully arranged in order over the gauze mask, which prevents them coming in contact with the face.

The sutures having been inserted the advancement forceps are removed by cutting through the muscle close behind them ; care must be taken not to cut the sutures. The middle stitch

should be tied first. A double turn is taken in the first stage of the knotting. Absolute parallelism in the visual axes is obtained as follows. A light is placed on the ceiling immediately above the patient's face. The gauze mask is slit through so that both eyes can fix the light. When the axes are parallel the corneal reflex of the light should be in the centre of each pupil. I usually tie the sutures with an ordinary surgical knot, but if desired they may be tied in bows, so that further adjustment may be made during the first few days after the operation. The ends of the stitches must be cut short so that they do not touch the cornea.

Both eyes should be bandaged for three days or longer. The stitches are removed under cocain in a week or ten days. A slight astringent lotion may be used until the conjunctival injection has disappeared. Occasionally a knob of granulation tissue appears in the wound ; this should be snipped off. The correcting spectacles must be used as soon as both eyes are uncovered.

The chief difficulty in performing advancement, as already pointed out, is getting a firm anterior hold for the sutures. It is a common opinion that the stitches tear out through the muscle. Consequently a great variety of fancy stitches have been devised for obtaining a firm posterior hold. I am convinced that this rarely happens, for the muscle is surrounded by quite dense connective tissue which prevents tearing out. When an advancement fails it is because a firm grip on the sclerotic has not been obtained. It is useless to pass the needle through conjunctiva only ; it must penetrate well into the episcleral, and even scleral, tissue. There is of course some danger of perforating the globe, and disastrous results have been known to arise from this cause ; they are fortunately very rare.

Since tenotomy produces some slight proptosis and impairment of movement of the eye it is best in moderate deviations to do an advancement only. In such cases the muscle may be tucked or reefed instead of being divided and advanced. The operation is very similar to advancement. Partial tenotomies and devices for lengthening the tendon of an extrinsic ocular muscle are in my opinion of little value.

Congenital Defects.

One or more of the extrinsic muscles may be absent as a congenital defect, or may be abnormally inserted into the sclerotic. In some cases the condition has been proved to be due to the

absence of the motor nervous mechanism. The position of the eyes and their movements may be very varied, but sometimes resemble those of an ordinary internal squint. In the latter case peculiarities of movement, *e.g.*, in and up instead of inwards, can usually be elicited. When each eye is made to fix successively the movements of the other eye are often quite different from each other. There is never double vision and muscular contracture does not occur. There is often congenital ptosis (*q.v.*) not infrequently of the hereditary type, and sometimes nystagmus,

SECTION VI.

SYMPTOMATIC DISEASES OF THE EYE.

CHAPTER XXIX.

OCULAR MANIFESTATIONS OF DISEASES OF THE NERVOUS SYSTEM.

Many diseases which primarily attack other parts of the body give rise to ocular symptoms, and not infrequently first



Fig. 289.—Relations of the cranial nerves to the arteries at the base of the brain.

come under the observation of the ophthalmic surgeon. At the risk of some repetition I propose briefly to review the most important ocular manifestions of such diseases.

The ocular signs of nervous disease often appear super-

ficially to be complicated and confusing. In most cases they are readily explained by the anatomy of the part of the nervous system involved.

Tabes Dorsalis.—*Primary Optic Atrophy* (*vide* p. 365) occurs in about 10–20 per cent. of cases of locomotor ataxia. It is about twice as common in men as in women, most frequent between thirty and fifty years of age, and may precede the appearance of typical tabetic symptoms by some years. It is commonest in the pre-ataxic stage, but it is not true that optic atrophy deters the development of ataxy or exercises any beneficial influence. The onset is gradual, leading to total blindness in two to three years or more. Pallor of the disc may precede the failure of vision by a considerable period, never the reverse. The affection of one eye usually precedes that of the other by a few months, rarely longer. The disease is a primary neuronic degeneration, starting in the ganglion cells of the retina; this accounts for the bad prognosis (cf. Disseminated Sclerosis, p. 543).

The *fields* show progressive contraction, pari passu with the failure in central vision. It is rare for the failure of sight to commence with a central scotoma, thus differing from the onset in disseminated sclerosis, though Fuchs has shown that it occurs. Two types of field are met with: (1) General concentric shrinkage, the colour fields for red and green being very early lost, and central vision much impaired; (2) Irregular sectorial defects, which are sharply defined but gradually spread, though central vision may be quite good. Defective dark adaptation and coloured vision have been described as early symptoms of tabes.

The characteristic *pupillary signs* include the so-called spinal miosis, the Argyll Robertson pupil reaction, inequality of the pupils, and distortion of the pupillary aperture. These signs are found in other diseases, and are to be regarded as signs of syphilis of the central nervous system rather than as pathognomonic of tabes. Their combination is of great diagnostic significance. Argyll Robertson pupils are found in 70 per cent. of tabetics and are almost invariably bilateral. Unequal pupils are found in 30 per cent. of tabetics, but are still more frequently met with in general paralysis of the insane. Ophthalmoplegia interna, *i.e.*, paralysis of the sphincter iridis and of the ciliary muscle, occurs in about 5 per cent. of tabetics and is generally unilateral. It is due to a lesion in the nucleus of the IIIrd nerve. Cycloplegia without mydriasis, or vice versâ, is rare.

Paralyses of the Extrinsic Ocular Muscles are common in tabes, occurring in about 20 per cent. of the cases. The order of frequency of the nerves affected is IIIrd (20 per cent.), VIth (13 per cent.), IVth (3 per cent.), external or total ophthalmoplegia (2 per cent.). It is characteristic of tabetic paralyses that they are partial, *i.e.*, not involving the whole nerve, incomplete, *i.e.*, pareses rather than paralyses, variable and transitory. The affection of the IIIrd nerve is so common that it is always suggestive of a tabetic or syphilitic lesion. Total IIIrd nerve paralysis is rare in tabes, and isolated ptosis is very common (4 per cent. of cases). The pareses of the ocular muscles nearly always occur in the pre-ataxic stage: when they occur at a later stage they are more likely to be permanent. They generally clear up rapidly, but show a marked tendency to recur. They are due to nuclear lesions. Considering that tabes is essentially a disease of afferent tracts it is curious that the Vth nerve almost always escapes. Nystagmus is rare, but the paresed muscles often give rise to jerky movements of the eyes which may be mistaken for nystagmoid jerks. Paralyses of associated movements, *e.g.*, conjugate deviations, are very rare.

In **Combined Sclerosis**, in which both the posterior and the lateral columns of the cord are affected, all the ocular manifestations characteristic of tabes may occur; these are probably atypical cases of tabes. In subacute combined sclerosis, such as is specially associated with anæmia and cachexia, ocular symptoms are rare.

In **Hereditary Ataxy** (*Syn.—Friedreich's Disease*) optic atrophy and paralyses of the ocular muscles are very rare. Nystagmoid jerkings of the eyes, very similar to those occurring in disseminated sclerosis, are very common, but the visual symptoms characteristic of the latter disease are absent. The movements are probably due to the same lack of co-ordination which causes the other ataxic signs of the disease; they occur on voluntary movement, and are not usually present in passive fixation.

Congenital Spastic Paralysis (*Syn.—Little's Disease*) is probably due to subdural hæmorrhage resulting from difficult labour. Of these cases 30—40 per cent. have concomitant convergent squint.

Myelitis.—A considerable number of cases of optic neuritis associated with myelitis have been described. The visual defect usually precedes the signs of myelitis. Its onset is sudden, but one eye may be affected a day or so before the

other. Complete amaurosis generally supervenes rapidly. In some cases there is a central scotoma in the early stages, and there may be pain on moving the eyes, pointing to a retrobulbar neuritis. There is usually only slight neuritis, but considerable swelling of the disc has been seen. In cases which recover the blindness passes off and good vision is restored. Since the site of the myelitis may be lumbar or dorsal, and as anatomical continuity of the inflammatory process from the cord to the optic nerves cannot be demonstrated, it is almost certain that the ocular affection is due to toxins circulating in the blood. There are no signs of general meningitis, and other cranial nerves escape.

During the amaurotic stage the pupils are dilated and immobile. In cases of cervical and upper dorsal myelitis without optic neuritis the pupils are often unequal, owing to implication of the dilatator tracts. In these cases the pupils still react to light.

Disseminated Sclerosis (*Syn.—Multiple or Insular Sclerosis*). —Lesions in disseminated sclerosis often occur in the visual paths (50 per cent. of cases). Unlike the lesions of tabes, the medullary sheaths of the nerve fibres are specially attacked, the axis cylinders remaining relatively little affected. Hence, during the acute stage, defects in conductivity are specially prominent; considerable variations succeed each other, and high degrees of functional restoration are possible. The optic nerves are most frequently attacked, with all the clinical signs of a typical retrobulbar neuritis, but patches of degeneration in the chiasma, optic tracts, or optic radiations may cause characteristic hemianopic changes in the fields. In ordinary cases there is central scotoma with a full field. The scotomata are generally only relative: they are therefore easily overlooked, and can be demonstrated only by the use of small coloured objects. Concentric contraction of the field and irregular peripheral defects, sometimes only for colours, also occur, and these show variations from time to time. Hemianopic fields are rarer than might be expected.

The ophthalmoscopic changes show very little direct relationship to the visual defects. There may be complete blindness with normal fundi, and signs of optic atrophy may coincide with good vision. Owing to the relative escape of the axis cylinders much less peripheral degeneration occurs than might be anticipated. Owing to the recovery of conductivity in the fibres, vision generally improves materially, but repeated relapses are not uncommon. Permanent com-

plete blindness scarcely ever occurs. Uhthoff found marked optic atrophy in 3 per cent., incomplete atrophy in 19 per cent., temporal pallor of the disc in 18 per cent. and optic neuritis in 5 per cent. of cases. The optic nerves are affected, therefore, much more commonly than in tabes, more often indeed than in any other disease of the nervous system except cerebral tumour.

The visual symptoms of disseminated sclerosis may be mistaken for those of toxic amblyopia, retrobulbar neuritis, tabes, or hysteria. The diagnosis from the two former is the most difficult. In ordinary retrobulbar neuritis the central scotoma is usually absolute; in disseminated sclerosis relative. In toxic amblyopia the scotoma is practically always bilateral; in disseminated sclerosis it is unilateral in about half the cases. In retrobulbar neuritis it is nearly always unilateral, and to these cases the early stages of disease of the pituitary gland must be added. Some cases can only be definitely diagnosed by the history and by the development of other pathognomonic signs. With regard to tabes central scotoma is rare, and on the other hand symmetrical concentric contraction of the field is rare in disseminated sclerosis. Moreover, the failure of vision is steadily progressive in tabes, and is bilateral; in disseminated sclerosis it is unilateral and very variable. The diagnosis from hysteria may be difficult, but the regular concentric contraction of the field so often found in this condition scarcely ever occurs in disseminated sclerosis.

Nystagmus occurs in multiple sclerosis (12 per cent. of cases), but nystagmoid jerks are much commoner (50 per cent. of cases). True nystagmus is a very important diagnostic sign, as it is rare in other acquired diseases of the central nervous system: nystagmoid jerks are of much less diagnostic significance. Both are probably due to central changes, and the latter show some analogy to the intention tremor so characteristic of disseminated sclerosis.

Miosis is fairly common in this disease, and to a less degree inequality of pupils. Other abnormal pupil reactions are rare.

Paralyses of extrinsic ocular muscles are much less common than in tabes, and although resembling these in their partial and transitory nature differ from them in that paralyses of associated movements are not uncommon. Thus paresis of convergence, with retained action of the recti in lateral movements, frequently occurs. Paralysis of lateral conjugate

movements is commoner than that of upward and downward movements. These are obviously due to nuclear lesions. Of individual nerves the VIth is more often affected than the IIIrd, and total IIIrd nerve paralysis is never seen (cf. Tabes). Partial ophthalmoplegia externa, with intact intrinsic muscles, also occurs, whereas opthalmoplegia interna is unknown.

Syringomyelia is due to dilatation of the central canal of the cord and excavation of the central grey matter. Since the dorsal and lower cervical regions are often affected inequality of the pupils is the most characteristic ocular sign. It is due to implication of the dilatator tract; the pupil on the affected side is small and reacts to light, but does not dilate after instillation of cocain (*vide* p. 58). Other signs of paralysis of the cervical sympathetic may be present, such as slight ptosis, retraction of the globe, &c. Paralysis of the Vth nerve is also not uncommon, and the VIth nerve may be affected, but very rarely the IVth or IIIrd. Syringomyelia, however, is sometimes complicated with tabes, and in these cases all the ocular manifestations of tabes may occur. The patients are often hysterical, a fact which accounts for the frequency of concentric contraction of the fields of vision.

Myasthenia Gravis. –This disease shows some resemblance to chronic progressive bulbar paralysis, but differs from it, *inter alia*, in the fact that the ocular muscles are almost invariably affected. Most of the patients are young, and have difficulty in articulation, swallowing, and mastication. There is nearly always ptosis and paresis of the orbicularis palpebrarum. The muscles of the extremities and trunk become affected, or the disease may start in them. Dyspnœa is common, especially on exertion. A striking feature is the absence of muscular atrophy. The muscles do not give the reaction of degeneration, but show the "myasthenic reaction," *i.e.*, they respond worse and worse to repeated faradic stimuli. The most characteristic feature is the rapid fatigue of the muscles. The symptoms are least marked in the morning, *e.g.*, the ptosis is much worse in the evening. Reading is only possible for a few minutes owing to failure of convergence and lateral movements of the eyes. Only a few mouthfuls of food can be masticated owing to fatigue of the muscles. The same applies to other voluntary muscles, which are similarly rapidly tired out by electrical stimulation. The muscles recover rapidly in the early stages of the disease after a short rest. The symptoms fluctuate from day to day, and may remain in abeyance for considerable periods. Sensory and

cerebral symptoms are absent, and the reflexes are normal. Many cases die of failure of respiration, though the course of the disease is usually long. No pathological changes are found in the nervous system, but groups of lymphocytes ("lymphorrhages") have been found in the muscles, and the thymus is sometimes enlarged.

The ptosis is nearly always bilateral and is increased by prolonged fixation or attempts to look upwards. Effective compensation by overaction of the frontales is impossible. Ophthalmoplegia externa, partial or complete, occurs in 50 per cent. of the cases. The intrinsic muscles are not affected. Nystagmoid jerks are not uncommon.

Myotonia Atrophica is a familial, hereditary disease, characterised by weakness of muscles (facial, vasti, &c.), and slow relaxation after contraction. The patients frequently develop cataract at an early age—20 to 40—and the cataracts may be the first manifestation of the disease.

Ophthalmoplegia.—Ophthalmoplegia is a somewhat indefinite term applied to widespread paralysis of the muscles of the eye which is thought to be due to nuclear lesions and forms the most prominent feature of the cases. Nuclear paralyses often cause defects of convergence, conjugate deviation of the eyes, and so on; when these are isolated signs the term ophthalmoplegia should not be applied to them. In typical ophthalmoplegia both eyes are affected, though all the muscles need not necessarily be simultaneously paralysed. In ophthalmoplegia totalis all the muscles, extrinsic and intrinsic, are affected: in ophthalmoplegia externa, many or all of the extrinsic muscles; in ophthalmoplegia interna, the intrinsic muscles. Ophthalmoplegia may be acute or subacute and chronic.

Acute or subacute ophthalmoplegia is usually due to poisons or infection, and is relatively rare. The chief poisons are alcohol, lead, and ptomaines; the chief infections diphtheria and influenza. In acute alcoholism the onset is sudden and accompanied by cerebral symptoms—headache, delirium, coma, &c. Bilateral ophthalmoplegia externa comes on suddenly or rapidly, with or without ptosis, and is often followed by facial and bulbar paralysis, with difficulty in speech and swallowing. The intrinsic muscles usually escape. The condition is probably due to acute hæmorrhagic superior polioencephalitis (Wernicke). In lead poisoning the onset is less acute and the intrinsic muscles are more often involved.

In ptomaine poisoning, due to bad food, mussels, &c., the essential feature is bilateral ophthalmoplegia interna, with or without ptosis, but total ophthalmoplegia also occurs. In diphtheria isolated ocular palsies are common, but ophthalmoplegia externa is rare. The pupil often escapes, the ciliary muscle never. In influenza the ophthalmoplegia resembles that of diphtheria—extrinsic muscles and ciliary muscle, the pupil escaping, but the pupil has been known to be affected without the ciliary muscle. The prognosis in alcoholism is bad: other cases usually recover.

Chronic ophthalmoplegia is usually progressive. It commences with ptosis or diplopia. In the course of months or years the paralysis spreads to all the ocular muscles of both sides, except that the intrinsic muscles often escape, and not infrequently the levatores palpebrarum also. These cases of isolated chronic ophthalmoplegia are rare, but the condition is often a precursor or symptom of tabes or general paralysis of the insane, rarely of disseminated sclerosis, &c. It is a very early sign of tabes, and may become associated later with bulbar symptoms. The Argyll Robertson pupil or ophthalmoplegia interna is often present.

Ophthalmoplegia also occurs as a congenital disease or may be acquired early in life as an hereditary familial disease. In these cases there is usually only partial ophthalmoplegia externa, and the condition is not complicated by other nervous disease such as tabes or bulbar paralysis, thus differing from the adult acquired form.

Diseases of the Pons.—The ocular symptoms are of great localising value in diseases of the pons. Of these, tumours are by far the most common, hæmorrhages, thromboses, softening, and abscess being relatively rare.

Tumours.—By far the commonest tumours of the pons are tubercle and glioma, the former being about twice as common as the latter. Both occur most frequently in childhood. Papillœdema or papillitis occurs in about half the cases, and is accompanied by the usual visual symptoms. The most characteristic signs of pontine tumours are due to implication of the motor nuclei and pyramidal tracts (*vide* Figs. 260—266). The VIth nucleus is usually implicated and causes loss of conjugate movement of the eyes to the same side (*vide* p. 496), with retention of convergence: at the onset the external rectus only may be paralysed. Owing to the immediate vicinity of the pyramidal tract it is also generally involved, and as the fibres are affected before they decussate in the medulla

oblongata there is contra-lateral hemiplegia. The intimate relationship of the VIIth nucleus and its afferent fibres to the VIth nucleus has already been mentioned. Hence, facial paralysis combined with loss of conjugate deviation of the eyes to the same side suggests a pontine lesion. Similarly, facial paralysis with contra-lateral hemiplegia (Millard-Gubler's syndrome) has the same significance. If the lesion is situated high up in the pons the pyramidal tract is caught before the fibres to the facial nucleus have crossed. Hence in these cases there is facial paralysis combined with hemiplegia on the same side. The facial paralysis is then usually of the cerebral type, in which the orbicularis palpebrarum, which is said to be innervated from the IIIrd nucleus, escapes. Not infrequently the Vth nerve is partially paralysed, causing for example paralysis of the Vth, nuclear VIth, VIIth, and sometimes VIIIth nerves, with crossed hemiplegia. It is astonishing how large pontine tumours can become without causing death, the nervous structures being pushed aside, especially in the relatively slow development of tuberculous masses. Extension of the disease may lead to IIIrd nerve paralysis, practically never IVth, the fibres of which are protected by the dorsal position of their decussation. If ptosis is the only sign of involvement of the IIIrd nerve its localising value is slight it may be a mere pressure symptom or a cerebral ptosis. Occasionally the opposite pyramidal tract is involved, with bilateral hemiplegia. Owing to the combination of facial paralysis the cases in which the trigeminal is involved are more likely to cause neuroparalytic keratitis (*q.v.*) than in other lesions of the Vth nuclei or intra-medullary fibres. Miosis is not uncommon in tumours of the pons, but the pupillary signs are of little diagnostic value ; nystagmus is a sign of involvement of the cerebellum.

In *hæmorrhages* and *thromboses* in the pons the same motor signs are manifest, and are usually of rapid or sudden onset. There are no ophthalmoscopic changes. The pupils are usually very small in the early stages of pontine hæmorrhage, a point of considerable diagnostic significance in an unconscious patient.

Tumours of the Auditory Nerve (*Syn.—Extra-cerebellar Tumours*).—The peculiar slow-growing neuro-fibromatous or endotheliomatous tumours of the recessus acustico-cerebellaris, usually attached to the VIIIth nerve, give rise to a fairly characteristic syndrome with ocular signs. Early deafness on one side is associated with cerebellar symptoms, among

which nystagmus is common. The VIth nerve is usually involved, generally with paralysis of the external rectus only, rarely with paralysis of conjugate deviation. As might be expected there is very often facial paralysis of the peripheral type, *i.e.*, total including the orbicularis palpebrarum. The Vth nerve is implicated in about a quarter of the cases, but neuroparalytic keratitis is uncommon. In nearly all the cases there is pronounced papillœdema.

Diseases of the Cerebral Peduncle (Figs. 54, 255).—The most characteristic sign of disease of the cerebral peduncle is a combination of paralysis of the IIIrd nerve with contra-lateral hemiplegia, the latter including the face and tongue (Weber's syndrome). The facial paralysis is naturally of the cerebral type, in which the orbicularis palpebrarum escapes, since it is due to a pyramidal tract lesion. If the red nucleus (Fig. 255) is involved tremor and jerky movements occur in the contra-lateral side of the body : this condition, combined with ipso-lateral IIIrd nerve paralysis, forms Benedikt's syndrome. Motor and sensory hemiplegia, contra-lateral to the lesion, without IIIrd nerve paralysis, is less common than Weber's syndrome, and IIIrd nerve paralysis alone is rare. In the usual syndrome the whole IIIrd nerve is involved, the intrinsic muscles rarely escaping : when it occurs it is due to an intra-peduncular fascicular lesion. Both oculomotor nerves are sometimes affected. As might be expected (Figs. 54, 255) implication of the external geniculate body or optic tract may occur, with development of homonymous hemianopia. Since the commonest lesion in this region is solitary tubercle papillœdema occurs in about 10 per cent. of the cases. The most frequent other causes, omitting basal gummatous meningitis, which may affect the peduncle secondarily, are softening and hæmorrhage.

Diseases of the Corpora Quadrigemina and Pineal Gland.—Though there can be no doubt that visual functions are located in the optic lobes of lower animals these functions are submerged in their later representatives, the anterior colliculi. There is no good evidence that lesions of the corpora quadrigemina cause any direct impairment of vision in man. The anatomical relations of the posterior colliculi point to association with hearing, and lesions of the corpora quadrigemina are frequently accompanied by impairment of hearing, which may, however, be due to pressure on the auditory paths. The commonest lesion of these bodies is solitary tubercle, which acts like an intracranial tumour, and glioma. Tumours of

the pineal gland are generally gliomata ; they press upon the colliculi and cause similar though less pronounced symptoms.

As might be expected tumours in this region very frequently cause papillœdema from pressure on the aqueduct of Sylvius, and therewith deterioration of vision. The sign of greatest localising value is loss of upward and downward movement of both eyes. Sometimes only upward movement is lost, never downward alone. The other movements of the eyes are relatively good. It is noteworthy, in opposition to experimental data, that impairment of conjugate lateral movements of the eyes is almost unknown in these lesions. In more than half the cases there is paresis of both IIIrd nerves. Less often only one oculomotor nerve is affected. Extension of the pressure effects may lead to bilateral IVth nerve paralysis or ophthalmoplegia externa, but the VIth nerve, as might be expected, is seldom directly affected. Pupillary changes are common, owing to implication of the IIIrd nerves or papillœdema. Experimental and clinical evidence alike tend to show that, in spite of the intimate relationship of the afferent pupillary paths with the superior colliculi, lesions of these bodies cause no direct permanent changes in the pupillary reactions. The facial nerve is paralysed in about a quarter of the cases : the paralysis is of the cerebral type. It is occasionally accompanied by ipsolateral hemiplegia. These are distant signs, due to pressure. Nystagmus occurs more frequently than with other cerebral tumours, but is usually associated with defects of co-ordination and other signs of implication of the cerebellum. These cases are always difficult to diagnose from cerebellar lesions, and the order of onset of the symptoms is important. If the ocular movements are affected first, and especially if upward and downward movements are lost, the lesion is probably quadrigeminal ; if the cerebellar ataxy precedes the impairment of ocular movements the lesion is probably in the cerebellum (Bruns).

General Paralysis of the Insane. (*Syns. Progressive Paralysis, Paralytic Dementia*).—Like tabes, this is a parasyphilitic disease. It is often accompanied by tabetic signs and symptoms which are due to lesions of the posterior tracts of the cord, identical with those in tabes (tabo-paralysis). The ocular symptoms are most common and unequivocal in these cases, and are to be attributed to the same causes.

The *pupillary* changes are most characteristic. In the early stages inequality of the pupils is most common. It

should be quite definite to be of diagnostic value, for slight inequality is not very infrequent in normal people. It is often accompanied by slight deformation in the shape of the pupil and irregularity of the pupillary margin. The same remark applies to these changes. The pathological nature of the pupillary changes is put beyond doubt when there is the typical Argyll Robertson reaction. It occurs in nearly half the cases, and is therefore an important sign, but less constant than in tabes. In about 5 per cent. of the cases the reactions both to light and convergence are lost, a condition which is rare in tabes and especially frequent in the juvenile form of general paralysis. The sensory reaction, *i.e.*, dilatation of the pupil on painful stimulation of the skin, is very often lost with the light reaction. The Argyll Robertson pupil is rare in cases in which the knee jerks are retained. Spinal miosis is commoner in tabes, unequal pupils in general paralysis. Ophthalmoplegia interna is rarer in general paralysis.

Primary optic atrophy occurs in about 8 per cent. of cases (Uhthoff). It shows exactly the same type and course as in tabes, but is more frequent in the latter disease. Like the pupillary signs, it may precede the onset of the typical cerebral symptoms by a considerable period, especially in those cases which commence with tabetic symptoms.

Paralyses of the extrinsic ocular muscles occur about half as frequently as in tabes, and have exactly the same characteristics, the IIIrd nerve being most frequently involved.

Cerebral Syphilis is the term usually applied to relatively early, direct syphilitic disease of the brain and meninges. Its manifestations differ very materially from those of the parasyphilitic diseases, and the ocular symptoms are of special diagnostic importance. Cerebral syphilis is due essentially to gummatous inflammation of the meninges and the walls of the cerebral blood vessels.

The chief form of brain syphilis is basal gummatous meningitis. It usually arises from the subarachnoid tissue in the region of the chiasma and spreads thence over the base of the brain. The optic nerves, chiasma, and tracts are generally involved. Papillitis, papillœdema, or post-neuritic atrophy are frequently found (about 13 per cent. each), and are usually bilateral. Visual defects are very common, and consist of amblyopia, not infrequently amaurosis, and defects in the fields of vision. Of the latter many cases show homonymous hemianopia from affection of one tract, fewer cases temporal hemianopia. Central scotoma and other signs of retrobulbar

neuritis also occur. The IIIrd nerve is paralysed in a third of the cases, less commonly the Vth and VIth, and least frequently the IVth. The IIIrd and VIth are often affected on both sides. The trigeminal paralysis is always unilateral and often causes neuroparalytic keratitis. Pupillary changes occur, dependent upon the IIIrd nerve lesions. In many cases the process is limited to a small area, oculomotor paralysis or an affection of the visual path being the only signs except headache. A very characteristic feature of basal gummatous meningitis is the inconstancy and variability of the symptoms, temporary and recurrent visual and oculomotor disturbances being very common.

Isolated gummata may give rise to the signs of cerebral tumour, complicated by the fact that they are often multiple. Syphilitic disease of the cerebral vessels is responsible for a large proportion of cases of thrombosis, hæmorrhage, softening, &c.

Intracranial Tumours.—The commonest ocular manifestation of intracranial tumours is papillœdema or papillitis. The latter term is used in this connection for the slight degrees of swelling of the disc, and does not necessarily imply a true inflammatory process.

Analysis of 200 cases of intracranial tumour treated at the National Hospital, Queen's Square, shows the following results (Paton) :—

(1) Precentral tumours are nearly always associated with neuritis fairly severe in character. (2) Postcentral tumours are nearly always associated with neuritis, as a rule moderate, and often of very short duration. (3) Temporo-sphenoidal tumours are always associated with neuritis of about the same degree of severity as in frontal tumours. (4) Of subcortical tumours about one-half develop neuritis—as a rule, moderate in degree—and, as in the case of parietal tumours, frequently of short duration. (5) Optic thalamus and mid-brain tumours are almost invariably associated with optic neuritis of very great severity. (6) Cerebellar tumours are constantly accompanied by neuritis of a grave character. (7) Extra-cerebellar tumours, as a rule, develop neuritis of a grave character. (8) Of pontine tumours, about one-half only develop optic neuritis, and then only when neighbouring parts of the brain, especially the cerebellum, have become involved. The neuritis when it does develop is usually very severe. (9) Ventricular tumours develop a moderate neuritis.

Homonymous hemianopia is due in about half the cases in

which this symptom is present to tumours of the occipital lobes. About 20 per cent. are due to involvement of one tract, either direct or as a pressure symptom. Relatively few are due to involvement of the internal capsule or external geniculate body. Heteronymous hemianopia is much rarer and is due to pressure on the chiasma and tracts by tumours of the pituitary body or distension of the third ventricle.

Paralysis of ocular muscles is relatively rare and nearly always a distant pressure symptom. One or both VIth nerves are often affected, the IIIrd nerve rarely, the IVth practically never. Conjugate lateral deviation of the eyes, which is common in cerebral hæmorrhage and to a less extent in cerebral softening, is rare with tumours of the cerebrum. It is more frequent with cerebellar tumours. Paralyses as true localising symptoms may of course occur with tumours of the crus, pons, &c. To this category belongs trigeminal paralysis, with or without neuroparalytic keratitis: it is rare with cerebral, commoner with cerebellar tumours.

Intracranial Abscess.—Cerebral abscess occurs about three times as often as cerebellar. The majority of cerebral abscesses are due to middle ear disease and affect the temporal lobes. Others are due to traumatism and generally affect the parietal lobes. Rarer causes are metastatic infection, usually derived from the lungs, frontal sinus empyema, and orbital cellulitis. A still greater proportion of cerebellar abscesses is due to otitis media.

Nearly half the cases have either papillœdema or papillitis: it is not infrequently on the side of the abscess only, and in bilateral cases the swelling is generally greater on this side. This sign has therefore greater localising value in intracranial abscess than in tumour. Optic neuritis persists longer after operation for abscess than for tumour, or may even only then commence. As might be expected, optic atrophy is rare during the acute stage: its presence militates against the diagnosis of abscess. Ophthalmoscopic changes are rarer with extradural abscesses.

Homonymous hemianopia indicates a lesion of the occipital lobe, which is rarely due to otitis.

Partial unilateral IIIrd nerve paralysis is fairly common, and the combination of unilateral ptosis and mydriasis has almost pathognomonic significance of ipso-lateral cerebral or cerebellar abscess. Partial IIIrd nerve paralysis with contralateral hemiplegia points to abscess of the temporal lobe with pressure on the IIIrd nerve and internal capsule, or more

rarely to implication of the cerebral peduncle. Paralysis of the VIth nerve is not common, but is found rather oftener in cerebellar than cerebral abscess: it is generally ipso-lateral, but has little localising value. Paralysis of the Vth nerve is rare. Nystagmus is very common with cerebellar abscess, but rare with cerebral. In otitic cases it may be due to disease of the labyrinth.

Oxycephaly and Scaphocephaly (*Syn. Tower Skull*) are due to precocious union of certain cranial sutures, especially the coronal and sagittal. The great wing of the sphenoid is displaced so that the orbit becomes shallow, causing more or less proptosis. In the early stages there is papillœdema, but more commonly only the later stage of post-neuritic optic atrophy is seen. The amount of atrophy varies in degree. The papillœdema is probably due to increased intracranial pressure, owing to continued growth of the brain in a restricted space. Divergent strabismus and horizontal nystagmus are common. Most of the patients are males.

Encephalitis.—Ocular palsies usually usher in an attack of *encephalitis lethargica*. Ptosis is the commonest feature, and other branches of the IIIrd nerve are specially involved. The muscles are usually only partially paralysed, and generally recover. Diplopia is an early symptom, and nystagmus may be present. Papillitis is rare and the pupils are usually normal. The general symptoms are lethargy, with great muscular debility, and other signs of an acute general infection.

Acute *polioencephalitis* accounts for not infrequent cases of paralytic squint following a febrile attack in young children. The VIth nerve is most often involved.

Meningitis.—In *tuberculous meningitis* a moderate degree of papillitis is common (about 25 per cent.) and is generally bilateral. Papillœdema occasionally occurs and indicates the combination of solitary with miliary tubercle. Tubercle in the choroid is frequent and of great diagnostic importance. A review of the literature tends to show that it is less common in tuberculous meningitis than in generalised miliary tubercle, but my own observations lead me to think that it is much commoner than is generally thought. It is often found only a day or two before death. There are often partial ocular pareses, usually of the IIIrd nerve, especially in the form of ptosis. Bilateral IIIrd paralysis is almost unknown, a point of distinction from syphilitic basal meningitis. Unilateral partial VIth nerve paralysis also occurs. Not infrequently

there is a kinetic (not paralytic) conjugate deviation of the eyes and head to one side.

In *epidemic cerebro-spinal meningitis* papillitis is frequently present, never papillœdema ; it is due to a descending infective neuritis. In the early stages there is often kinetic strabismus or conjugate lateral deviation of the eyes. A characteristic sign is the widely open palpebral aperture, often associated with very infrequent blinking. Paralysis of the VIth nerve, usually unilateral, is commoner than that of the IIIrd, though divergent strabismus due to the latter cause has been frequently noted. Total IIIrd nerve paralysis is rare (cf. Gummatous basal meningitis). The pupils vary much, usually showing miosis in the early stages, mydriasis when coma sets in : loss of reaction to light is relatively rare. Conjunctivitis and keratitis sometimes occur, and many cases of metastatic endophthalmitis (*q.v.*) in children are due to the Weichselbaum meningococcus, though it is a relatively rare complication of the disease.

Still and others have shown that the sporadic acute basal meningitis of children is due to the meningococcus. A peculiarity of this disease which I have frequently seen at Great Ormond Street Children's Hospital is complete amaurosis with normal fundi and normal pupil reactions, pointing to the action of toxins on the higher visual centres. The blindness may persist for many weeks after subsidence of other symptoms, and sight may be completely restored. Chronic basal meningitis sometimes shows the same feature, but in these cases optic neuritis and post-neuritic atrophy may occur from secondary hydrocephalus and pressure of the distended third ventricle upon the chiasma and tracts.

Purulent meningitis occurs occasionally in typhoid, and more rarely in pneumonia, influenza, scarlet fever, measles, and septicæmia. In typhoid the diagnosis is difficult, but the presence of papillitis and ocular paralyses points in this direction. Metastatic purulent meningitis, with papillitis or retrobulbar neuritis, occurs in children from obscure causes. Middle ear disease is a not uncommon cause of purulent meningitis. In this condition papillitis or papillœdema is usually due to complications, such as sinus thrombosis or cerebral abscess. When ocular paralysis occurs, the VIth nerve is usually affected, rarely the IIIrd (cf. Intracranial abscess). The facial nerve is most frequently involved, the paralysis often causing lagophthalmia. Conjugate deviation of the eyes is not uncommon. Metastatic endophthalmitis is rarer

than in epidemic cerebro-spinal meningitis. The diagnosis of otogenous meningitis from tuberculous may be difficult or impossible.

Hydrocephalus.—In the congenital and the early acquired hydrocephalus of infancy optic atrophy is not infrequently found. Papillœdema occurs only rarely in spite of the increased intracranial pressure. This fact is doubtless due to the relief of pressure by the enlargement of the skull and the resiliency of the fontanelles and gaping sutures, as well as to the very gradual development. The eyeballs usually deviate downwards, and upward movements are much restricted. This is sometimes due to bulging of the thin orbital plate of the frontal bone, which may be even absorbed. Not infrequently there is considerable proptosis.

The acquired hydrocephalus of later life, after the fontanelles and sutures have closed, is always rather a doubtful inference than a precise diagnosis. The cardinal signs of increased intracranial pressure—headache, vomiting and papillœdema—are present, and to these is often added ataxia of the cerebellar type. The cases are often diagnosed as intracranial tumours, in which localising signs are not infrequently absent or masked. Bitemporal hemianopia may give a clue to the true ætiology, being due to pressure on the chiasma and tracts by the bulging floor of the third ventricle. In some cases there is evidence of previous meningitis, but the most characteristic feature is often the variability of the symptoms. Remissions and intermissions of long duration occur, and recovery or arrest of the condition is not uncommon, often, however, with defective vision due to post-neuritic atrophy.

Fractures of the Base of the Skull.—Unilateral facial paralysis is the commonest cranial nerve lesion in fractures of the base of the skull (22 per cent. of cases): the VIth (4 per cent.), IIIrd (2 per cent.), Vth (1·6 per cent.), and IVth (1 per cent.) follow in order of frequency. Fractures of the base from falls upon the head, &c., very often pass through the optic foramen and involve the roof of the orbit: occasionally both optic foramina are broken. Owing to the intimate union between the dura mater and periosteum the optic nerve is frequently injured. It may be torn through, lacerated by splinters of bone, or compressed by hæmorrhage into the sheath. If the nerve is completely severed there is blindness with a normal fundus in the early stages. In two to four weeks signs of primary optic atrophy appear and progress to total atrophy. Optic neuritis indicates hæmorrhage into the

nerve sheath, and may occur from basal hæmorrhage without fracture of the optic foramen. Hæmorrhage into the sheath or nerve may cause concentric contraction of the field of vision, or quadrant and other sectorial defects: central scotoma appears to be rare. Most cases with rapidly developing papillitis die. Pigmentation in and around the disc may follow hæmorrhage into the sheath. The pupil reactions vary and are not pathognomonic, but there is usually mydriasis on the side of the lesion.

Statistics.—The following statistics (in percentages), derived from Uhthoff's very extensive investigations of the literature and of cases in the Breslau clinic, give some idea of the frequency of important physical signs in certain diseases of the nervous system already discussed. Too much reliance must not be placed upon the percentages, culled as the cases are from very various sources.

	Papilloedema.	Papillitis.	Optic Atrophy.	Homonymous Hemianopia.	Bitemporal Hemianopia.	III Paralysis.	IV Paralysis.	VI Paralysis.	V Paralysis.	Conjugate Deviation of the Eyes.	Nystagmus.
Cerebral Tumour	53	18	8	17	1	14	0·6	11	6	3	4
Cerebellar Tumour	53	24	11	—	—	5	2	18	12	1·5	25
Cerebral Abscess	23	21	0·3	9	—	19	1·6	10	4	6	4
Cerebellar Abscess	23	22	—	—	—	14	—	12	4	6	42
Cerebral Syphilis	14	12	14	11	6	34	5	16	14	1	8
Cerebral Hæmorrhage	11	6·5	1	29	—	9	—	8·4	1	28	10
Cerebral Softening	1·4	2·2	0·8	40	—	2·4	—	0·3	2	12	1·6
Tuberculous Meningitis	5	29	1	—	—	18	1	12	5	8	10
Internal Hydrocephalus	23	20	19	—	—	6	—	13	—	1	13
Pituitary Body Tumours	9	8·5	21	3	32	17	2·5	6	—	—	4
Fractures of the Base of the Skull	9		4	—	—	2	1	4	1·6	2·5	3

CHAPTER XXX.

OCULAR MANIFESTATIONS OF OTHER DISEASES.

The most important ocular manifestations of other diseases than those of the central nervous system have already been discussed incidentally, and it will suffice here merely to enumerate them.

Infectious Diseases.—Mucopurulent conjunctivitis and corneal ulcers are the chief ocular complications of *measles*. They are rare in *scarlet fever*, as is also albuminuric retinitis. Corneal ulcers are common in *small pox*. *Vaccinia* of the eyelids is not uncommon, and may affect the cornea, usually secondarily, sometimes causing disciform keratitis. *Diphtheria* most commonly causes cycloplegia (*vide* p. 490), and rarely paralysis of the external rectus. *Erysipelas* may cause abscesses and gangrene of the lids, orbital cellulitis, and thrombosis of the orbital veins and cavernous sinus. Optic neuritis occurs in *typhoid fever*. Conjunctivitis and the herpetic types of keratitis are common in *influenza* which also causes iritis and optic neuritis.

Diseases of the Respiratory Tract.—Conjunctival hæmorrhages are common in *whooping cough*, and retinal hæmorrhages may also occur. Herpes corneæ occurs in *pneumonia*, but it is remarkable that hypopyon ulcer is rare. Apical *phthisis* may cause irritation of the sympathetic fibres, leading to dilatation of the ipso-lateral pupil.

Diseases of the Circulatory System.—Pulsation of the retinal vessels, embolism and thrombosis of the central artery, arterio-sclerosis, and thrombosis of the central vein of the retina have already been sufficiently discussed. Aneurysm at the root of the neck may cause dilatation of the pupil on the same side.

Diseases of the Blood.—*Chlorosis* has often been held responsible for papillitis, but it is extremely doubtful if the diagnosis is accurate. Retinal hæmorrhages occur in the *secondary anæmias* of carcinoma, ankylostomiasis, &c., and are a prominent sign in *pernicious anæmia*, which really belongs to this group. In this disease they are often of characteristic

colour (*vide* p. 344). The ophthalmoscopic signs of *lukæmia* have already been described. Hæmorrhages in and about the eyes are common in scurvy, purpura, and hæmophilia. Great *loss of blood* often leads to amblyopia or amaurosis, and may be followed by bilateral optic atrophy. Severe ocular symptoms very rarely follow traumatic hæmorrhage, as in war injuries, but most commonly result from intestinal or uterine hæmorrhage. Though both eyes are usually affected there is often an interval of days between them. The discs are hazy at first, later becoming atrophic, with constricted vessels. In some cases slight improvement of sight eventually occurs.

Diseases of the Organs of Digestion.—*Oral sepsis*, especially pyorrhœa alveolaris, is an undoubted cause of iridocyclitis, and probably causes choroiditis and other forms of subacute or acute endophthalmitis. Infective lesions in the mouth may spread by continuity, especially along the veins of the pterygoid plexus, setting up orbital cellulitis or thrombosis of the cavernous sinus. Lamellar cataract is associated with hyoplasia of the enamel of certain teeth. The lacrymal gland is not infrequently affected in *parotitis*, and symmetrical inflammation of the lacrymal and salivary glands is characteristic of *Mikulicz' disease*. Absorption of bacteral toxins from the intestinal canal is almost certainly a cause of iridocyclitis and other obscure inflammations of the uveal tract. Night blindness is associated with some *diseases of the liver*, *e.g.*, cirrhosis, and jaundice causes yellow discoloration of the conjunctiva, but yellow vision (xanthopsia) is much less common than has been thought.

Diseases of the Kidneys.—Albuminuric retinitis and uræmic amaurosis have already been discussed.

Metabolic Diseases.—Ocular complications are common in *diabetes mellitus*, the most frequent being diabetic cataract, retinitis, intraocular hæmorrhages, and retrobulbar neuritis. Œdema of the pigment epithelium on the back of the iris is often seen in microscopical specimens, but iritis is seldom met with. Remarkable changes in the refraction of the eye, both in the direction of hypermetropia and myopia, have been described in this disease, and paralyses of both extrinsic and intrinsic ocular muscles also occur. *Gout* has been held responsible for deposits in the conjunctiva (concretions), conjunctivitis, marginal ulcers of the cornea, episcleritis, scleritis, iritis, and other conditions : it is indirectly the cause of ocular lesions through the kidneys and vascular system. *Rheumatism* is an indefinite entity. Acute rheumatism is doubtless

an infective disease : it practically never gives rise to iritis, but may cause embolism of the central artery of the retina indirectly by its effects on the cardiac valves. It rarely causes optic neuritis. Chronic rheumatism is also probably due to organisms or bacterial toxins, and is thus responsible for iritis, cyclitis, episcleritis, and retrobulbar neuritis. Interstitial keratitis has been described in *myxœdema :* over-feeding with thyroid causes exophthalmos and amblyopia.

Diseases of the Generative Organs.—It can scarcely be doubted that the profound changes which the generative organs, especially in the female, undergo at puberty, in menstruation, parturition, and the climacteric, are often associated with disorders of metabolism and other pathological conditions. Of these the albuminuric retinitis of pregnancy is the most impeccable example among ocular complications. Many others have been described, *e.g.*, conjunctivitis associated with the menses, but their relationship to diseases of the generative organs is obscure. The diseases due to gonorrhœa and syphilis have been described elsewhere.

SECTION VII.

DISEASES OF THE ADNEXA OF THE EYE.

CHAPTER XXXI.

DISEASES OF THE LIDS.

Anatomy.—The lids are covered anteriorly by skin and posteriorly by mucous membrane—conjunctiva tarsi; they end in a free edge about 3 mm. broad—margo intermarginalis. The substance of the lids consists of muscle, glands, blood



FIG. 290.—Diagram of sagittal section of upper lid.

vessels, and nerves, all bound together by connective tissue, which is particularly dense at the posterior part, where it forms a stiff plate—the tarsus (Fig. 290).

The skin of the lids differs from that of the rest of the body merely in its thinness, its loose attachment, and the absence of fat in its corium. It is covered with fine downy hairs, which are provided with small sebaceous glands, and there are also small sweat glands. At the anterior border the hairs are specially differentiated to form a protection to the eyeball.

The cilia or eyelashes are strong, short, curved hairs, arranged in two or more closely set rows. Their sebaceous follicles, like the cilia themselves, are specially differentiated, and are called *Zeis's glands*. Apart from being larger, they are identical with other sebaceous glands. The sweat glands near the edge are also unusually large and are known as *Moll's glands*. They are situated immediately behind the hair follicles, and their ducts open into the ducts of Zeis's glands or into the hair follicles, not direct on to the surface of the skin as elsewhere.

The margin or free edge of the lid is the part between the anterior and posterior borders—the intermarginal strip or margo intermarginalis (Fig. 291). It is covered with stratified epithelium, which forms a transition between the skin and the conjunctiva proper. The anterior border is rounded; the posterior, which lies in contact with the globe, is sharp. The



Fig. 291.—Diagram of intermarginal strip.

capillarity induced by this sharp angle of contact is of importance in the proper moistening of the surface of the eye. Immediately anterior to the posterior border is a single row of minute orifices, just visible to the naked eye. These are the orifices of the ducts of the Meibomian glands. Between this row of puncta and the anterior border is a fine grey line, which is important in operations in which the lid is split, as it indicates the position of the loose fibrous tissue between the orbicularis palpebrarum and the tarsus.

The tarsus consists of dense fibrous tissue; it contains no cartilage cells, so that the term tarsal cartilage is only justified in so far as it defines the consistence of the plate. Imbedded in the tarsus are some enormously developed sebaceous glands, the *Meibomian glands*. They consist of nearly straight tubes, directed vertically, each opening by a single duct on the margin of the lid. The tubes are closed at the upper end, and have numerous small cæcal appendages projecting from the sides, filled with fatty glandular epithelium. The glands

number from twenty to thirty, being rather fewer in the lower than in the upper lid.

The large bundles of the orbicularis palpebrarum occupy the space between the tarsus and the skin. The main central band of the levator palpebræ superioris is inserted into the upper border of the tarsus; an anterior slip passes between the bundles of the orbicularis to be inserted into the skin of the middle of the lid; a posterior slip is inserted into the conjunctiva at the fornix. The inferior rectus and oblique muscles send fibrous strands forwards into the lower lid to be attached to the tarsus and palpebral ligament.

Besides these striped muscles there is a layer of unstriped muscle in each lid. These constitute the superior and inferior tarsal muscles of Müller. The fibres of the former arise among the striped fibres of the levator, pass down behind it, and are inserted into the upper border of the tarsus. The inferior lies below the inferior rectus and is inserted into the lower tarsus.

The arteries of the upper lid form two main arches, superior and inferior, the former lying between the upper border of the tarsus and the orbicularis, the latter in a similar position just above the hair follicles. In the lower lid there is usually only one arch near the free edge. There are two venous plexuses in each lid: a post-tarsal passing into the ophthalmic veins, and a pre-tarsal opening into subcutaneous veins.

The sensory nerve supply is derived from the trigeminal. The third nerve supplies the levator palpebræ, the seventh the orbicularis, and the sympathetic Müller's muscles.

Inflammation of the Lids.

Almost any of the inflammatory conditions which affect the *skin* in general may attack the lids. Erysipelas is dangerous in that it may spread to the orbit, leading to cellulitis and atrophy of the optic nerve, thrombosis of the cavernous sinus, or meningitis. Herpes ophthalmicus is often mistaken for erysipelas: its unilaterality, strict localisation to the course of branches of the ophthalmic nerve, and the characteristic formation of vesicles should prevent this mistake; permanent scarring remains after the attack. Eczema of the lids is common, especially associated with phlyctenular conjunctivitis (*q.v.*) in children, and with atropin irritation (*q.v.*). Abscesses, boils, anthrax pustule, and ulcers of various kinds may affect the skin of the lids. Œdema of the lids may be inflammatory or passive. It is often associated with chemosis of the con-

junctiva in severe conjunctivitis. Great œdema is often caused by bites of parasites, gnats, &c., and by styes, abscesses, and chancre of the lid. In unilateral œdema the condition of the lacrymal sac and nasal duct should be investigated; it is often due to lacrymal abscess. Situated above the internal palpebral ligament it suggests empyema of the frontal sinus; in the lower lid empyema of the antrum. Localised œdema may be due to periostitis of the orbital margin. In all cases of œdema the condition of the eyeball must be determined, with the assistance of Desmarres' retractors if necessary. Œdema of the lids may be caused by serious purulent inflammation of the globe (panophthalmitis), of Tenon's capsule, by phlegmon of the orbit or thrombosis of the cavernous sinus. Passive œdema may be due to nephritis, heart disease, &c., or it may be angioneurotic. Chronic thickening of the lids, resembling œdema, but harder in consistency, may follow recurrent attacks of erysipelas—so-called *solid œdema*.

Blepharitis is a chronic inflammation of the margins of the lids. It may manifest itself as a simple hyperæmia, differing from that caused by weeping, exposure to tobacco smoke, and so on, in being more persistent. The causes and treatment are the same as for the more severe forms of blepharitis. True blepharitis occurs in two forms. In *squamous blepharitis* small white scales, like dandruff, accumulate among the lashes; the latter fall out readily, but are replaced without distortion. If the scales are removed the underlying surface is found to be hyperæmic, but not ulcerated. The condition is probably a seborrhœa.

In *ulcerative blepharitis* yellow crusts glue the lashes together; on removing them small ulcers, which bleed easily, are seen around the bases of the lashes. The lashes fall out or are easily pulled out, and often are not replaced, or grow in a distorted form, owing to injury to the follicles. Blepharitis causes redness of the edges of the lids, itching, soreness, lacrymation, and "photophobia."

The sequelæ of the ulcerative form are serious. If not treated energetically and with perseverance the disease is extremely chronic, causing or being accompanied by chronic conjunctivitis. Care must be taken to distinguish true blepharitis from matting together of the lids by conjunctival discharge; in the latter case removal of the crusts reveals quite normal lid margins. The ulceration is liable to extend deeply, so that the hair follicles are destroyed. Only a few small, scattered, distorted cilia are then found (*madarosis*).

When the ulcers heal the cicatricial tissue contracts. Neighbouring hair follicles are drawn out of place, and a false direction is given to the remaining cilia, so that they may rub against the cornea (*trichiasis*). Or the development of cicatricial tissue may be extreme, so that the edge of the lid becomes hypertrophied and droops in consequence of its weight (*tylosis*).

The lower lid is particularly liable to be displaced by prolonged ulcerative blepharitis. The contraction of the scar tissue drags the conjunctiva over the margin; the posterior lip of the intermarginal strip, instead of being acute-angled, becomes rounded, so that its capillarity is impaired. Tears then tend to run over (*epiphora*), a condition which is accentuated if the punctum becomes everted, so that it ceases to lie in accurate contact with the bulbar conjunctiva (*vide* p. 591). The continual wetting of the skin with tears leads to eczema, which is followed by contraction. The condition is made worse by perpetually wiping the eyes, so that eventually *ectropion* is developed. This causes still more epiphora, a vicious circle being set up.

The causes of blepharitis are multitudinous. The patients are usually children debilitated from living under poor hygienic conditions, or from disease, *e.g.*, anæmia, tubercle, syphilis, measles, &c. The condition may follow chronic conjunctivitis, or be induced by the same causes, especially smoky atmosphere, heat (stokers, cooks), late hours, &c. It may result from a neglected diplobacillary blepharo-conjunctivitis. It is undoubtedly often associated with uncorrected errors of refraction, especially hypermetropia and astigmatism, which probably act by inducing reflex hyperæmia. Occasionally parasites cause blepharitis, *e.g.*, blepharitis acarica, due to demodex folliculorum, and phthiriasis palpebrarum, due to the pediculus pubis, very rarely to pediculus capitis. In the latter condition the cilia are covered with black nits, an appearance being produced which is easily recognised when once seen.

Treatment.—The local treatment of blepharitis must be energetic in the ulcerative form. The crusts must first be removed. This is effected most easily by soap and water, followed by thorough bathing with hot borax or bicarbonate of soda lotion, 3 per cent. The application softens the deposits, so that they can be picked or rubbed off with a pledget of cotton wool. When the crusts have been entirely removed the surface is covered with an ointment of yellow oxide of mercury,

ammoniated mercury, or ichthyol (10 per cent.), which is well rubbed in for at least five minutes. These procedures should be repeated three times a day. In most cases, if the treatment is carried out properly, there is a speedy cure. Unfortunately the treatment is seldom satisfactorily carried out. It is useless merely to smear ointment on the surface of the crusts. It must be applied to the inflamed tissues and rubbed well into the hair follicles.

In more severe cases, or when the above treatment is improperly done, protargol, 15 to 20 per cent., should be rubbed into the margins of the lids with a stumpy camel's hair brush until a lather is formed; this usually takes five minutes. Or the surface may be thoroughly cleaned with bicarbonate lotion, and silver nitrate, 2 per cent., painted on. All loose lashes should be pulled out with epilation forceps. If diplobacilli are present a zinc lotion should be used.

Attention must be directed to the hygienic surroundings, and to the general health. Errors of refraction must be corrected.

Syphilis. A primary sore is occasionally found on the lid margins, commencing in the conjunctiva. It may be caused by a kiss or by removing a foreign body with the tongue. There is generally a small ulcer, covered with scanty greyish secretion and much indurated about the base. If situated near the outer canthus the pre-auricular gland is enlarged, if near the inner canthus the submaxillary, in accordance with the distribution of the lymphatic vessels. The swelling of the glands is always suggestive of syphilis or tubercle, but in all doubtful cases scrapings should be examined for spirochætes and the blood examined by the Wassermann test. If recognised before the glands are affected the sore should be excised or destroyed by radium. At a later stage calomel ointment should be rubbed in. An energetic course of constitutional treatment with salvarsan and mercury should be immediately instituted.

Gummata occur in the lids sometimes, and occasionally may cause enormous thickening of the tarsus (*syphilitic tarsitis*). Isolated gumma may be mistaken for a chalazion. In syphilitic tarsitis the lid may be so swollen and hard that it is impossible to evert it. The pre-auricular gland is swollen. If the onset is slow there is little pain; sometimes the swelling is rapid and very painful. Gummata usually respond rapidly to appropriate treatment with mercury and iodides.

Vaccinia.—The margin of the lid is occasionally inoculated

from the recently vaccinated arm of a baby. Often the inoculated margin in turn inoculates the opposing margin of the other lid. Usually the pustule is at the outer canthus, and the pre-auricular gland is swollen and painful. The history generally serves to elucidate the case. Sometimes the cornea becomes affected, a keratitis resembling disciform keratitis resulting, *i.e.*, a grey disc, denser at the margin.

Inflammation of the Glands of the Lids.

Hordeolum or **stye** is a suppurative inflammation of one of Zeis's glands (Fig. 292). In the early stages the gland becomes swollen, hard and painful, and usually the whole edge of the

Fig. 292.—Hordeolum.

lid is œdematous. An abscess forms which generally points near the base of one of the cilia.

The pain is considerable until the pus is evacuated. Styes often occur in crops, or may alternate with boils on the neck, carbuncles, or acne. Like these conditions, the disease shows deficient resistance of the body to the invasion of staphylococci. It is commonest in young adults, but may occur at all ages, especially in debilitated persons. Not infrequently it will be found that faulty drains account for the defective health.

Treatment.—Hot compresses should be used in the early stages. When the abscess points it may often be evacuated by pulling out the corresponding cilium; but this is usually effected more satisfactorily by an incision with a small knife. It should be remembered that such an incision is exceptionally

painful. The pus should be thoroughly squeezed out and a hot compress applied.

If crops of styes occur the general health must receive attention. When associated with boils, &c., the urine should be tested for sugar, especially in adults. Inquiry should be made as to the condition of the drains. In obstinate cases a staphylococcic vaccine, preferably autogenous, should be used.

Constipation must be counteracted, and tonics are useful, especially iron in some form.

Hordeolum internum is comparatively rare. It is a suppurative inflammation of a Meibomian gland of exactly the same type as the hordeolum externum or stye. It is often called a suppurating chalazion, and some may be due to secondary



Fig. 293.—Chalazion.

infection of a chalazion. The inflammatory symptoms are more violent than in external stye, for the gland is larger and is imbedded in dense fibrous tissue. The pus appears as a yellow spot shining through the conjunctiva when the lid is everted. It may burst through the duct or through the conjunctiva, very rarely through the skin.

Treatment is the same as for external type, except that the incision should be made exactly as for a chalazion (*vide infra*).

Chalazion (*Syns.—Tarsal Cyst, Meibomian Cyst*) is a chronic inflammatory affection of a Meibomian gland. The gland tissue becomes replaced by granulation tissue containing giant cells ; the disease is not caused by the tubercle bacillus, but is probably caused by the chronic irritation of an organism of low virulence. The gland becomes swollen, increasing in size very gradually and without inflammatory symptoms.

Patients usually seek advice on account of the disfigurement (Fig. 293). The smaller chalazia are difficult to see, but are readily appreciated by passing the finger over the skin. If the lid is everted the conjunctiva is red or purple over the nodule, in later stages often grey, or rarely, if infection has occurred (*vide* Hordeolum internum), yellow. The grey appearance is due to alteration in the granulation tissue. This is not very vascular at any stage, but in the later stages the vessels retrogress, the nourishment of the tissue fails and it becomes converted into a jelly-like mass. Only under such conditions is the term "cyst" really applicable. Complete spontaneous resolution very rarely occurs. The contents may be extruded through the conjunctiva, and in these cases a fungating mass of granulation tissue often sprouts through the opening, keeping up conjunctival discharge and irritation. Sometimes the granulation tissue is formed only in the duct of the gland, from which it projects as a reddish-grey, somewhat translucent nodule on the intermarginal strip (*marginal chalazion*).

Chalazia are often multiple or occur in crops. They are commoner in adults than in children.

Treatment.—Quite small chalazia may be left alone: it is very difficult to evacuate them satisfactorily by the ordinary method. Larger chalazia must be incised and thoroughly scraped. The conjunctival sac is well cocained with 2 per cent. solution and a drop of adrenalin instilled. The lid is then everted and the site of the chalazion carefully examined. At the point of greatest discoloration a few crystals (not many) of solid cocain are placed upon the surface and allowed to dissolve, or a little novocain or alypin solution may be injected with a hypodermic syringe. A vertical incision is

FIG. 294. — Beer's knife (devised for cataract extraction; now used only for lid operations).

then made with a sharp scalpel or Beer's knife (Fig. 294). Any semi-fluid contents which may be present escape. A small sharp spoon (Fig. 295) is then inserted into the orifice and the walls of the cavity are thoroughly scraped. The bleeding soon stops, and no dressing is usually necessary. A simple boric acid lotion is ordered for a few days.

The patient should be warned that the swelling will remain for awhile. This is due to the resistant walls, formed by the fibrous tissue of the tarsus; the cavity is kept dilated and becomes filled with blood. Sometimes, especially if the scraping has not been sufficient, granulation tissue sprouts from the wound. This must be snipped off with scissors, curved on the flat, after application of cocain; the cavity should be again scraped out.

Fig. 295.—Sharp spoon.

Very hard chalazia are occasionally met with, particularly near the canthi; it is possible that some of these are true adenomata of the glands. They may require excision, since it may be impossible to scrape them out efficiently.

If a marginal chalazion is not treated the granulation tissue protrudes from the mouth of the gland and may organise into a greyish, somewhat translucent lump of fibrous tissue on the lid margin. It is disfiguring and rather difficult to remove without leaving an irregularity in the line of the lid.

Anomalies of Position of the Lids.

Trichiasis (*θρίξ, τρίχος*. a hair) is the condition of distortion of the cilia, so that they are directed backwards and rub against the cornea (*vide* p. 571). A few only of the lashes may be affected, or the condition may be due to entropion, involving the whole margin of the lid. It may also be caused by congenital distichiasis (*vide* p. 588).

The symptoms are those of a foreign body continually present in the eye—irritation, pain, conjunctival congestion, reflex blepharospasm, lacrymation. Superficial opacities and vascularisation of the cornea are produced; recurrent ulcers of the cornea are not infrequently due to this cause.

Any condition causing entropion (*q.v.*) will cause trichiasis, trachoma and spastic entropion being among the most common.

Other causes are blepharitis, and the scars resulting from injuries, burns, operations, diphtheria, &c.

Treatment.—Isolated misdirected cilia may be removed by epilation, which must, however, be repeated every few weeks. A better mode of treatment is to destroy the hair follicle by electrolysis. The flat positive pole is applied to the temple; the negative, a fine steel needle, is introduced into the hair follicle: a current of two milliampères is used. The negative pole is determined by placing the terminals in saline, when bubbles of hydrogen are given off by it. The strength of current can be gauged by the rate of evolution of gas. It should be remembered that electrolysis is extremely painful and tedious; the pain may be avoided by injecting novocain into the margin of the lid. If the current is of the proper strength, the bubbles evolved at the site of puncture cause the formation of a slight foam.



Fig. 296.—Griffith's ring forceps; useful for holding the lid everted in incising chalazia. The ring is placed on the conjunctival surface and surrounds the chalazion.

If many cilia are displaced, operative procedures must be resorted to. Since they are nearly allied to those performed for entropion, which is generally present, they will be described later (*vide* p. 574).

Entropion (ἐν, in, τρέπειν, to turn), rolling in of the lid, occurs in two forms, spastic and cicatricial. The symptoms are those of the trichiasis (*q.v.*) which is induced.

Spastic entropion is due to spasm of the orbicularis. Strong contraction of the circularly arranged fibres tends not only to approximate the lid margins, but also to turn them inwards or outwards, according to the mechanical support afforded by the globe and orbital contents. If the support is insufficient, entropion is produced. This is well seen when the eyeball has

been removed, but it also occurs when the globe is deeply set owing to absence of orbital fat, &c., especially if the skin of the lids is also redundant. These conditions are found *par excellence* in old people, who are therefore very liable to spastic entropion. It is also caused by tight bandaging, and is favoured by narrowness of the palpebral aperture (blepharophimosis). Spastic entropion is almost invariably restricted to the lower lid.

Cicatricial entropion is caused by cicatricial contraction of the palpebral conjunctiva; in the worst forms, found in trachoma, the tarsal plate is also bent and distorted, sometimes by atrophic, sometimes by hyperplastic, changes. It is an



Fig. 297.—Diagram of skin and muscle operation.

exaggeration of the effect produced by the various causes of trichiasis (*q.v.*).

Treatment of Spastic Entropion.—If due to bandaging, the condition is often cured by simply leaving off the bandage. Wearing an artificial eye relieves the symptoms when the eyeball has been removed. In the spastic entropion of old people temporary relief may be obtained by placing a roll of lint or plaster horizontally just above the margin of the orbit, and bandaging it firmly in position; or the lid may be slightly everted by painting collodion on the skin or by pulling it out with a strip of adhesive plaster.

Permanent relief can be obtained only by operation. The simplest method is the removal of a strip of skin and muscle. Cocain, eucain or novocain is first injected subcutaneously. An oval area of skin, with the long axis horizontal and varying in width according to the amount of entropion and of super-

fluous skin, is marked out with a scalpel or Beer's knife just below the site of greatest displacement. The upper incision must be close to the margin of the lid (Fig. 297). The piece of skin is dissected off. The underlying fibres of the orbicularis are then dissected off with forceps and knife, until the tarsus is exposed. It is usually unnecessary to insert sutures. The operation may be done with one snip of scissors and without cocain if a fold of skin is taken up with T-shaped forceps (Fig. 298). This method is not so satisfactory.



Fig. 298.—T-shaped forceps.

In cases of spastic entropion with much blepharospasm, canthoplasty is sometimes indicated. It consists in widening the palpebral aperture by dividing the outer canthus. The lids are separated with the fingers in such a manner as to put the canthus on the stretch. One blade of strong blunt-pointed scissors is introduced as far as possible into the conjunctival sac behind the commissure. The entire thickness, including skin and conjunctiva, is divided horizontally by a single cut. If only a temporary effect is required, no sutures are inserted. If it is desired permanently to enlarge the palpebral aperture, the conjunctiva is sutured to the skin. Temporary canthoplasty is sometimes indicated in other conditions than spastic entropion, *e.g.*, in simple severe blepharospasm, such as occurs in phlyctenular conjunctivitis, in acute purulent conjunctivitis with much swelling of the lids, and in removal of an enlarged eyeball or an orbital tumour.

Treatment of Cicatricial Entropion.—A very large number of plastic operations have been devised for the relief of cicatricial entropion: only the more simple will be described here. The principles governing the various operations are: (1) altering the direction of the lashes, (2) transplanting the lashes, (3) straightening the distorted tarsus. Subcutaneous injection of cocain or a general anæsthetic is indicated: the former method does not obviate all pain, especially if the tarsus is cut.



FIG. 299.—Lid spatula, which should be made of metal.

The simplest procedure is some modification of Burow's operation. The lid is everted over the end of a metal lid spatula (Fig. 299). A horizontal incision through the conjunctiva and passing completely through the tarsal plate, but not through the skin, is made along the whole length of the lid in the sulcus subtarsalis, *i.e.*, about 2—3 mm. above the posterior border of the intermarginal strip (Fig. 300). Care must be taken not to wound the punctum or canaliculus. The outer end of the strip may then be divided by a vertical incision through the free edge of the lid, including the whole thickness. In this manner the edge of the lid is left attached only by skin, and when cicatrisation has occurred the edge is turned slightly outwards, so that the lashes are directed away from the eye. Relapses are not uncommon, however, and the operation may have to be repeated. It is probably better to keep the edge of the lid everted during the process of healing by means of a spindle-shaped pad of oiled silk (Fig. 301). The pad is kept in position by sutures suitably applied.

In the Jaesche-Arlt operation the zone of hair follicles is transplanted to a slightly higher position. The lid is split from the outer canthus to just outside the punctum along the grey line (*vide* p. 562) between the lashes and the orifices of the Meibomian glands. During this procedure the globe is protected by the spatula inserted between it and the lid, or held by a lid clamp (Figs. 302, 303). The incision extends between the tarsus and the orbicularis for a depth of 3—4 mm., so that the zone containing the hair follicles is thoroughly loosened (Fig. 304). An elliptical piece of skin is then removed from

the lid. The lower incision extends through the skin down to the tarsus at a distance of 3—4 mm. from the edge of the lid and parallel with it for its whole length. The middle part of the upper incision is 6—8 mm. from the edge of the lid. The elliptical piece of skin thus marked out is removed, without taking any orbicularis. The two skin incisions are then sutured. In this manner the zone of lashes is transplanted to a higher level. The gaping wound in the intermarginal strip may be filled in with skin from that excised, or by a Tiersch graft or a graft of mucous membrane ; this tends to prevent the follicles from being drawn down again when the wound



FIG. 300.—Diagram of modified Burow's operation for entropion.



FIG. 301. — Diagram of modified Burow's operation for entropion.

cicatrises. Care should be taken not to produce ectropion by removing too much skin.

Ectropion, rolling out of the lid, occurs in several forms, the chief being spastic, cicatricial, senile, and paralytic. The symptoms are due to the epiphora induced and to the chronic conjunctivitis caused by exposure. In severe cases the cornea may suffer from imperfect closure of the lids.

Spastic ectropion results from blepharospasm when the lids are well supported by the globe and when they are short, firm, and without redundant skin. It is therefore seen in children and young patients, and is readily induced by phlyctenular conjunctivitis (*vide* p. 159). Mechanical ectropion is caused by extreme proptosis or thickening of the conjunctiva, such as occurs after purulent conjunctivitis and trachoma. In the latter disease the tarsus is often distorted. Upper and lower lids are frequently affected simultaneously.

Cicatricial ectropion results from destruction of the skin by

injury, burns, ulcers, gangrene, operations, &c. Caries of the orbital bones is a common cause in children. Chronic conjunctivitis and blepharitis also cause cicatricial ectropion, which is



Fig. 302.—Desmarres' entropion forceps for right eye.



Fig. 303.—Wilde's entropion forceps.

increased by the wetting of the skin with tears and the eczema thereby induced.

Senile ectropion is found only in the lower lid, and is due to relaxation of the tissues and degeneration of the orbicular muscle fibres. The condition is increased by the conjunctivitis and epiphora which are set up.

Paralytic ectropion results from the laxity of the lids induced

by paralysis of the orbicularis. Only the lower lid is affected, the upper being kept in contact with the globe by its own weight.

Treatment.—Non-operative treatment is chiefly serviceable

Fig. 304.—Diagram of Jaesche-Arlt operation for entropion.

in spastic ectropion. Here a well-fitting bandage, unless contraindicated by other factors, will often cure the displacement. A bandage should also be applied in ectropion paralyticum, but the condition is cured only by restoration of the innervation. The slighter degrees of senile ectropion are also amenable to



Fig. 305.—Diagram of Snellen's sutures for ectropion.

non-operative treatment, though it may be advisable to slit the canaliculus in order to cure epiphora. The patient should be instructed not to pull the lid down when wiping the eye.

A large variety of operations has been devised for ectropion : only the simpler procedures will be described.

Snellen's sutures are indicated in some cases of spastic and senile ectropion. In this operation two loops of thread, inserted at the junction of the middle with the inner and outer thirds of the lid, through the ectropionised conjunctiva, are made to hold the fornix in its proper position (Fig. 305). A silk thread is armed with a needle at each end. One needle is passed in at one of the two spots indicated at the level of the top of the tarsus, *i.e.*, in the position where the fornix ought to be. The needle is passed vertically downwards under the skin and brought out just below the level of the orbital margin. The other needle is inserted similarly 2 or 3 mm. to one side of the first, and is carried down parallel with it. The second suture is inserted in exactly the same manner at the junction of



Fig. 306.—Dimmer's modification of Kuhnt's operation for ectropion.

the middle with the other third of the lid. The sutures are then tied over pieces of rubber tubing laid vertically, the ectropionised fornix being thus drawn down into its normal position. The sutures may be tightened from day to day so as to cause the formation of cicatricial bands along their tracks : this was more effectual in the days when antiseptic precautions were not observed. The effect is seldom permanent.

In paralytic ectropion lateral tarsorrhaphy may be indicated. In this operation the palpebral aperture is shortened by uniting the lids at the outer canthus. The edges of the upper and lower lids are freshened for the requisite distance, the lashes being excised. The lids are then sutured together as in central tarsorrhaphy (*vide* p. 216).

In many cases of ectropion, especially senile, the lower lid is

stretched and elongated. The ectropion may then be cured by shortening the lid. A wedge-shaped piece of the lid, including the whole thickness, is excised at the outer canthus. The edges of the skin wound are then sutured together, and similarly the edges of the conjunctival wound.

A more elegant method of shortening the lid is Dimmer's modification of Kuhnt's operation (Fig. 306). A triangular piece of conjunctiva and tarsus is excised, the apex of the triangle being towards the fornix. The lid is then split along the grey line from the triangle to the outer canthus. A triangular area of skin is removed at the outer canthus and the skin is slid outwards so that the gap in the tarsal plate is closed, the



Fig. 307.—Diagram of V—Y operation for ectropion.

requisite length of the margin of the lid at the outer canthus being denuded of cilia.

In the slighter cases of cicatricial ectropion the V–Y operation of Wharton Jones is indicated (Fig. 307). A V-shaped incision, with the apex away from the lid margin, is made through the skin, the limbs of the V enclosing the cicatrix. The skin is freed from the underlying tissues and is also well undermined at the edges. The margins of the incisions are sutured in such a manner that a Y-shaped cicatrix results: the edge of the lid is thus raised to its normal position.

More extensive cicatricial displacement requires some form of blepharoplastic operation, flaps of skin being taken from the cheek or temporal region, or skin grafts employed. Each such case must be treated on its own merits and will often exercise the ingenuity of the surgeon.

Symblepharon (*σύν*, with, together, *βλέφαρον*, eyelid) is the

condition of adhesion of the lid to the globe (Fig. 308). Any cause which produces raw surfaces upon two opposed spots of the palpebral and bulbar conjunctiva will lead to adhesion if the spots are allowed to remain in contact during the process of healing. Such causes are burns from heat or caustics, ulcers, diphtheria, operations, &c. Bands of fibrous tissue are thus formed, stretching between the lid and the globe, involving the cornea if this has also been injured. The bands may be narrow, but are more frequently broad, and may extend into the fornix so that the lid is completely adherent to the eyeball over a considerable area (symblepharon posterius). Bands limited to the anterior parts and not involving the fornix are called symblepharon anterius. Total symblepharon, in which the lids are completely adherent to the globe, is rare.



Fig. 308.—Symblepharon.

Pronounced adhesions cause impairment of mobility of the eye, so that diplopia may be complained of. The adhesion may be so intimate that it is impossible to close the lids efficiently, lagophthalmia, with its baneful consequences, resulting. There is often much disfigurement.

Treatment.—The prevention of symblepharon is of the utmost importance (*vide* p. 397). When it is already established operation is necessary. Symblepharon anterius is usually easily remedied by dividing the bands and preventing re-formation of adhesions in the manner already described. When the bands are broad, and especially if there is symblepharon posterius, the separation of the lid from the globe is difficult. There is no guide to the limitations of sclerotic and tarsus, and great care has to be exercised lest the globe be punctured. The prevention of re-formation of adhesions is much more difficult, and is successful only if the raw surfaces are covered with epithelial grafts. These may be either conjunctival or skin flaps, Tiersch grafts, or grafts of rabbit's or frog's mucous membrane.

Ankyloblepharon (*ἀγκύλη*, a thong, *βλέφαρον*, eyelid) is

adhesion of the margins of the two lids. It may be either a congenital condition or due to burns, &c. It may be partial or complete, and is often combined with symblepharon. The treatment depends upon the amount of symblepharon. If it is very extensive operation may be contraindicated. In other cases the lids are separated and kept apart during the healing process. If the adhesion extends to the angle of the lids, the latter must be covered with an epithelial graft, otherwise the condition will recur.

Blepharophimosis (*βλέφαρον*, eyelid, *φιμός*, a muzzle) is the condition in which the palpebral fissure appears to be contracted at the outer canthus. It is really normal, but the outer angle is obscured by a vertical fold of skin. The latter is due to eczematous contraction of the skin following prolonged epiphora and blepharospasm. Mere narrowing of the palpebral aperture is often called blepharophimosis, and may be a congenital condition: it is really a form of ankyloblepharon.

The condition may require no treatment, disappearing spontaneously after the inflammation has subsided. In other cases canthoplasty is indicated.

Lagophthalmia (*λαγώς*, a hare) is the condition of incomplete closure of the palpebral aperture when the eyes are shut. It may be due to narrowing of the lids from cicatrisation or congenital deformity, ectropion, paralysis of the orbicularis, proptosis due to exophthalmic goître, orbital tumour, &c., or to laxity of the tissues and absence of reflex blinking in people who are extremely ill or moribund. Owing to exposure the cornea becomes epidermoid (xerosis corneæ) or keratitis sets in. The treatment is that of keratitis e lagophthalmo (*q.v.*).

Ptosis (*πίπτειν*, to fall) is the term given to drooping of the upper lid, due to paralysis or defective development of the levator palpebræ superioris. Ptosis may also be caused by thickening and increased weight of the lid (*vide* p. 565). The condition may be unilateral or bilateral, partial or complete. In the higher degrees the lid hangs down, covering the pupil more or less completely, interfering with vision. An attempt is made to counteract the effect by overaction of the frontalis and by throwing back the head, the eyes being pulled downwards by the inferior recti. A very characteristic attitude is thus adopted. Forced contraction of the frontalis causes the eyebrows to be raised and throws the skin of the forehead into wrinkles. Partial ptosis may be masked by this means, but becomes manifest if the patient is asked to look up while the

eyebrows are fixed by firm pressure with the fingers against the bone.

Ptosis may be congenital or acquired. The congenital form is usually, but not invariably, bilateral, and is due in most cases to defective development of the muscles. Some cases have been proved to be caused by maldevelopment of the third nucleus. The condition is not infrequently hereditary. There is nearly always defect in the upward movement of the eyes, due partly to absence of the posterior insertion of the levator into the fornix (*vide* p. 563), partly to coincident maldevelopment or defective innervation of the superior rectus. It may be pointed out here that defective upward movement of the eyes is the commonest congenital defect of bilaterally associated extrinsic muscles.

Acquired ptosis is usually unilateral. It may be part of the symptom-complex of paresis or paralysis of the whole of the third nerve, or may be due to paresis or paralysis of the branch supplying the levator. Isolated ptosis without other signs of oculomotor paralysis may result from disease of upper level centres (cerebral ptosis). Acquired ptosis may also be due to direct injury of the muscle or its nerve supply, as by wounds, fractures, &c. Mechanical ptosis is due to deformity and increased weight of the lid brought about by trachoma, tumours, &c. ; it also occurs from lack of support in phthisis bulbi, anophthalmia, &c. Bilateral ptosis may occur in the acquired form, notably as part of the syndrome of myasthenia gravis.

The amount of ptosis sometimes alters with the position of the globe, attaining its highest pitch in abduction of the eye, its least in adduction or attempted adduction. Occasionally in both the congenital and acquired forms the lid rises when the jaw is moved, as in mastication, though it remains immobile when an attempt is made to look upwards (*vide* p. 513). This is an example of synkinesis or associated movement.

Treatment.—In cases of paralysis of the third nerve treatment must be directed to removal of the cause. The fact that this nerve is so frequently affected in syphilis must be borne in mind ; these cases respond to treatment better than others. In cases of incurable paralysis and in congenital and mechanical ptosis the deformity can be removed only by operation. In complete paralysis of the third nerve operation is usually contraindicated on account of the abduction of the eye. If the lid is raised in these cases the diplopia becomes manifest ; simultaneous advancement of the internal rectus may diminish

the diplopia and the deformity, but is unsatisfactory and unlikely to give a permanent result.

Operations for ptosis ameliorate the condition but seldom give permanent results. In slight cases excision of an elliptical area of skin, with or without excision of the underlying fibres of the orbicularis, improves the appearance temporarily.

Of the many operations which have been devised for the more severe cases Hess's operation is one of the best (Fig. 309). It may be performed under local anæsthesia by subcutaneous injections of novocain. The eyebrow is shaved. An incision is made in the line of the eyebrow for 2·5 cm. The skin of the lid is then



Fig. 309.—Diagram of Hess's operation for ptosis.

undermined through this incision so that it is completely separated from the orbicularis and tarsus over its whole area. Three silk sutures are then inserted. Each suture is armed with two half-curved needles. The first needle is inserted a little to one side of the middle line of the lid at a point 5 or 6 mm. from the margin of the lid. It is passed upwards through the orbicularis, anterior to the tarsus, and is brought out through the frontalis about 4 or 5 mm. *above* the incision in the eyebrow. The other needle on the same suture is inserted a little to the other side of the middle line of the lid at the same distance from the lid margin and is carried up parallel to the first and brought out in the same manner. The other two sutures are inserted in exactly the same manner on each side of the first. On pulling the sutures upwards the skin of the lid is thrown into a fold which corresponds with the normal fold at the upper margin of the orbit. In this manner the

lid is raised the requisite amount: care must be taken that it is not raised too much, otherwise lagophthalmia will result. The effect produced at the time of operation should be slightly in excess of the desired permanent effect. The incision in the eyebrow is sutured, and then the lid sutures are tied round small pieces of rubber tubing, applied vertically to the forehead. The sutures may be tightened or loosened, if necessary, during the first few days. They are removed after ten days or a fortnight.

Another good operation for congenital ptosis is *Motais'*. In this a narrow tongue from the mid-line of the superior rectus tendon is insinuated into a pocket between the skin and tarsus and sutured to the skin near the ciliary margin. After this operation the lid is carried upwards on upward movement of the globe.

Injuries of the Lids.

Injuries of the most various kinds—contusions, wounds, burns, &c.—are very common. They must be treated upon general principles, but special attention must be directed to three points—(1) wounds of the skin of the lids, (2) injury of the bones of the orbit, (3) injury of the eyeball.

Wounds in the direction of the fibres of the orbicularis gape little and heal without conspicuous scarring; hence surgical wounds should be made in this direction as far as possible. Vertical wounds gape, cause disfiguring cicatrices, and often lead to ectropion, or other distortion, especially if there is adhesion to the subjacent bone. The worst wounds are such as sever the lid vertically in its whole thickness. If they do not unite by first intention a notch (traumatic coloboma) is left in the lid margin, and disfigurement, lagophthalmia, and epiphora result. Vertical wounds severing the canaliculus require special care (*vide infra*).

Injury to the bones of the orbit may affect the orbital margin or deeper parts. Fractures involving the margin may be diagnosed by careful palpation –unevenness, crepitation, &c. Fractures of the walls of the orbit often manifest themselves by *emphysema*. It is due to communication of the subcutaneous tissues with the nasal air sinuses, air being forced into the tissues on blowing the nose, sneezing, straining, or coughing. There is great swelling, with a peculiar soft crepitation on palpation. Fracture of the orbital bones may be followed by retraction of the globe (traumatic enophthalmos), or may be part of a more serious fracture of the base of the skull. In the latter event the optic foramen is often involved, causing laceration or compression of the optic nerve (*vide* p. 368).

Injuries involving the globe require special care both in diagnosis and treatment. In every case of injury of the lids the eyeball must be very carefully examined. Palpation will usually demonstrate considerable reduction of intraocular pressure if the eye is injured ; it indicates rupture of the globe. Inspection may be difficult on account of excessive swelling and ecchymosis. In such cases the eye must be examined at all costs, the lids being separated by retractors, under an anæsthetic if necessary.

Contusions are often more alarming in appearance than in reality. There is great swelling and ecchymosis both in the lids and conjunctiva. In all cases a guarded prognosis should be given, for it may be impossible to determine the full extent of the injury to the orbit (*vide* p. 609), or the eye (*vide* p. 397).

Treatment.—Simple contusions with ecchymosis require only cold compresses : a simple boric lotion is ordered for cleansing the conjunctival sac, and boric ointment to prevent the lids from sticking together.

Emphysema should be treated with a pressure bandage, and all straining, blowing of the nose, and so on, must be avoided.

Wounds must be thoroughly cleansed with an antiseptic lotion and brought together by sutures. In wounds involving the canaliculus the inner cut end must be searched for, and the canaliculus slit up (*vide* p. 592). If this is not done before cicatrisation has occurred epiphora will follow, and it will be extremely difficult to obtain an entry into the canaliculus in order to slit it up. Lacerated wounds are likely to leave ugly scars and deformity of the lids : these must be treated by plastic operation (*vide* p. 572, &c.). If suppuration occurs the abscess must be opened and treated on general surgical principles.

Tumours of the Lids.

Benign tumours include xanthelasma, molluscum, warts, nævus, angioma, and other tumours common to the skin and cutaneous glands.

Small clear *cysts* frequently occur among the lashes in old people, due to retention of secretion of Moll's glands. They disappear if the anterior wall is snipped off.

Xanthelasma (*ξανθός*, yellow, *ἔλασμα*, a plate) or xanthoma is a slightly raised yellow plaque, most commonly found in the upper and lower lids near the inner canthus, and often symmetrical in the two lids and in the two eyes. The plaques are most common in elderly women. They grow slowly, and only

require treatment on account of the disfigurement produced. They may be removed after subcutaneous injection of novocain, or destroyed by electrolysis or radium.

Molluscum contagiosum is a small white umbilicated tumour, generally multiple. A substance resembling sebum can be squeezed out of it. Each tumour should be squeezed out after incision and the interior touched with solid silver nitrate stick.

Nævus or mole, usually pigmented, may occur on the lids, generally affecting the margin and involving both skin and conjunctiva. Two are often symmetrically situated on the lids of the same eye, indicating their origin at a time when the lids were still united. The microscopical appearance is characteristic, consisting of "nævus cells," often arranged in an alveolar manner. The growths may take on malignant proliferation.

Hæmangioma, often also called nævus, occurs in two forms —telangiectasis and cavernous hæmangioma. The former are bright red or port-wine coloured spots composed of dilated capillaries. The latter consist of dilated and anastomosing venous spaces lying in the subcutaneous tissue having all the characteristics of erectile tissue; they are not infrequently strictly localised as if partially encapsuled. They appear bluish when seen through the skin and form a swelling which increases in size on crying, lowering the head, &c. Cavernous hæmangiomata are rarely seen in adults, partly due to the fact that they are generally treated in early life, but possibly due to spontaneous atrophy of the growth and thickening of the skin.

Telangiectases may be excised if small. If larger they may be cauterised with the galvano-cautery; only a small area should be attacked at a sitting, the operation being repeated. Electrolysis or carbonic acid snow may also be used. Cavernous hæmangiomata may be excised, preferably from the conjunctival surface, if small. If larger they may be treated by electrolysis. It is a good plan to use electrolysis for a time until the tumour is consolidated with fibrous tissue, and then to excise the mass.

Lymphangioma occurs rarely in the lids.

Symmetrical soft swellings above the inner canthus are sometimes seen in elderly people. They are due to fatty deposits which are continuous with the orbital fat through an aperture in the fascial septum.

Malignant Tumours include carcinomata and sarcomata, the

former being much the more common. Epitheliomata show a preference for spots where the character of the epithelium changes: they therefore commence generally at the edges of the lids. The patients are elderly; the preauricular gland may be enlarged, or if the growth is near the inner canthus, the submaxillary lymphatic glands. Any of the glands of the lid may in rare instances undergo carcinomatous proliferation.

The commonest malignant epithelial growth is the so-called rodent ulcer, which shows a predilection for the inner canthus. It commences as a small pimple which ulcerates. If the scab is removed it is found that the edges are raised and indurated. The ulcer spreads very slowly, the epithelial growth extending under the skin in all directions and penetrating deeply. The surrounding structures are gradually destroyed; lids, orbit, and bones are invaded. The growth is only locally malignant and probably originates in the accessory epithelial structures of the skin—hair follicles and glands. The lymphatic glands are not affected. Rodent ulcer rarely occurs before forty years of age, and the rate of growth is of the order of years.

Sarcoma is rare; it may be round or spindle-celled, pigmented or non-pigmented. Round-celled growths, variously described as lymphoma, lymphosarcoma, pseudo-leukæmic tumours, &c., sometimes affect both orbits and all four lids causing symmetrical proptosis. Occasionally the patients show blood changes as in leukæmia, but these are usually absent. The growth is slow but continuous, and the eyes are lost from lagophthalmia. The malignant growths springing from nævi are usually called sarcomata.

Treatment.—Epithelioma and sarcoma must be thoroughly extirpated at all costs, even if it involves excision of the globe or exenteration of the orbit. Rodent ulcer, if small, should be excised, or destroyed by carbon dioxide snow. If larger it is wise to treat it with radium or X-rays for a time. Considerable improvement, and even cure, has been reported as the result of this treatment, but it is wise always to excise the remnants. In the later stages extensive plastic operations may have to be performed to protect the eyeball. When this becomes impossible the eye must be excised and the morbid tissues freely removed.

Congenital Abnormalities of the Lids.

Symblepharon, ankyloblepharon, ectropion, entropion and trichiasis occur occasionally as congenital malformations. Ptosis is a fairly common congenital defect.

Distichiasis (δίς, double, στίχος, a row) is a rare condition in which there are two complete rows of cilia, often in all four lids. The posterior row replaces the Meibomian glands: it causes trouble by rubbing against the cornea.

Coloboma of the lid is a notch in the edge of the lid. The gap is usually situated to the inner side of the middle line, generally affecting the upper lid. Two or more defects may occur in the same lid. Sometimes a bridge of skin links the coloboma to the globe, or there is a dermoid astride the limbus at the site of the coloboma. There are often other congenital defects of the eye or other parts of the body, *e.g.*, coloboma of the iris, accessory auricles, &c. Some cases are due to incomplete closure of the fœtal facial cleft, others probably to pressure of amniotic bands. Occasionally there is a notch at the outer part of the lower lid, associated with maldevelopment of the malar bone.

Cryptophthalmia (κρυπτός, hidden) is a very rare condition in which there is total ankylo- and sym- blepharon, associated with abnormality of the eye and often of the orbit. The skin passes continuously from the brow over the eye to the cheek.

Microblepharon is the condition in which the lids are abnormally small. They may be absent—*ablepharon*. These conditions usually occur only in cases of microphthalmia, or congenitally small eyes. Microphthalmia may be associated with a congenital *orbito-palpebral* cyst, causing a swelling of the lower lid. The cyst is connected with the eye-ball, contains retinal tissue in its lining, and is due to defective closure of the fœtal fissure—an extreme case of ectatic coloboma of the choroid (*q.v.*). The eye-ball may be apparently absent (*congenital anophthalmia*), but there are always microscopic vestiges of ocular tissues.

Epicanthus is a semilunar fold of skin, situated above and sometimes covering the inner canthus. It is usually bilateral, the eyes are far apart, and the bridge of the nose is flat. It may disappear as the nose develops. It is normal in Mongolian races.

Elephantiasis neuromatodes or **plexiform neuroma** may affect the lids and orbit. In all cases the temporal region is also affected. The swollen lid and temporal region form a characteristic picture. The hypertrophied nerves can be felt through the skin as hard cords or knobs. The nerve fibres are little changed, the hyperplasia affecting the endo- and peri- neurium. In several cases the ciliary nerves have been found affected, both outside and inside the globe, and in some cases the eye has been buphthalmic. Operative measures are seldom satisfactory.

CHAPTER XXXII.

DISEASES OF THE LACRYMAL APPARATUS.

Anatomy and Physiology.—The lacrymal apparatus consists of the lacrymal glands and the lacrymal passages.

The *lacrymal glands* of each eye consist of the superior or orbital gland, the inferior or palpebral gland, and the accessory lacrymal glands or Krause's glands. All are serous acinous glands, scarcely distinguishable, microscopically, from serous salivary glands, with which they are morphologically identical. The superior gland, about the size of a small almond, is situated in the lacrymal fossa at the outer part of the orbital plate of the frontal bone. Ten or twelve *lacrymal ducts* pass from it to open upon the surface of the conjunctiva at the outer part of the upper fornix. The inferior gland consists of only one or two lobules situated upon the course of the ducts of the superior portion. It can be seen when the eye looks down and in after the upper lid has been everted. The accessory or Krause's glands are microscopic acini, lying below the surface between the fornix and the edge of the tarsus. There are about forty-two in the upper, six to eight in the lower, fornix. The ducts of numerous acini unite to form a larger duct which opens on to the fornix.

The lacrymal passages consist of the puncta lacrymalia, the canaliculi, the lacrymal sac, and the nasal duct (Fig. 310). The *puncta lacrymalia* lie near the posterior border of the free margin of the lid about 6 mm. from the inner canthus. Each lid has one punctum and one canaliculus. The punctum is situated upon a slight elevation, larger in elderly people, the papilla lacrymalis. As already mentioned, this is visible under normal circumstances only when the lid is slightly everted (*vide* p. 79). The *canaliculus* passes from the punctum to the lacrymal sac. It is first directed vertically for about 1 to 2 mm., then horizontally for 6 to 7 mm. The canaliculi usually open separately through the outer wall of the lacrymal sac. The *lacrymal sac* lies in the lacrymal fossa formed by the lacrymal bone. When distended it is about 15 mm. long vertically, and 5 to 6 mm. wide. The fundus extends slightly

above the level of the inner tarsal ligament. The sac is surrounded by fibres of the orbicularis. The lower end narrows as it opens into the nasal duct. The *nasal duct*, varying much in size (12 to 24 mm. long, 3 to 6 mm. in diameter) passes downwards and slightly outwards and backwards, bounded by the superior maxilla and inferior turbinate, to open at the anterior part of the outer wall of the inferior meatus of the nose. The line of the duct is given by a point just outside the inner canthus and the groove between the ala of the nose and the cheek. The upper end of the nasal duct is the narrowest part. The canaliculi are lined by stratified epithelium, the lacrymal sac and nasal duct by columnar epithelium, lying



FIG. 310.—Diagram of lacrymal apparatus.

upon a corium which contains a venous plexus. The mucous lining forms an imperfect valve at the orifice into the nose.

The *lacrymal secretion* is a slightly alkaline fluid containing sodium chloride as its chief constituent. The ordinary amount secreted is just sufficient to moisten the eyeball, and is lost by evaporation. Only under reflex irritation, psychical or peripheral, is an excess secreted, and this is forced into the lacrymal sac and through the nasal duct into the nose during the act of winking, when the fibres of the orbicularis contract around the sac. It must be remembered that xerosis or dryness of the conjunctiva does not result from extirpation of the superior and inferior lacrymal glands, the moistening of the conjunctiva by Krause's glands and its own mucous cells being

sufficient to prevent it. *Per contra*, epiphora does not result from extirpation of the lacrymal sac, except in the presence of psychical or peripheral stimuli to increased secretion. It is possible that the tears have some slight antiseptic properties.

Diseases of the Lacrymal Gland.

Diseases of the lacrymal gland are rare. *Dacryo-adenitis* occurs occasionally, usually going on to suppuration. Tubercle also occurs here. A permanent *fistula* may result from rupture of an abscess in the gland. Spontaneous and traumatic *dislocation* of the gland have been described, a swelling being formed under the outer part of the upper lid.

Dacryops is a cystic swelling in the upper fornix, due to retention of secretion owing to blockage of one of the lacrymal ducts. It can only be distinguished from retention cysts of Krause's glands by its position.

Tumours of the lacrymal gland show a very marked resemblance to those of the parotid. In *Mickulicz' disease* there is symmetrical enlargement of the lacrymal and salivary glands, probably of a lymphomatous nature. Mixed tumours, in reality endotheliomata, containing cartilage, myxomatous material, &c., are the commonest form. Carcinomata and sarcomata are very rare.

All conditions which cause swelling of the gland may lead to impairment of movement of the eye. The globe is pushed downwards and inwards; movement outwards, and especially outwards and upwards, is limited. There may be some proptosis.

The rare diseases mentioned above must be treated on general principles.

Diseases of the Lacrymal Passages.

Eversion of the lower punctum occurs from laxity of the lids in old age, from chronic conjunctivitis, blepharitis, and any cause leading to ectropion (*q.v.*). It causes epiphora, which in turn aggravates the condition (*vide* p. 575).

Treatment.—In slight cases, especially in old people, the eversion may be sufficiently counteracted by making a small scar in the fornix just behind and below the position of the punctum. This is best done with the actual cautery, a fairly deep gutter being made. As the cicatricial tissue contracts the punctum is pulled inwards towards the eye.

If this procedure fails the canaliculus should be slit up or a radical operation for ectropion performed. In the former case the greatest care must be exercised that the canaliculus is slit on its posterior wall, so that the open duct will lie against the globe. It is not necessary, nor advisable, in these cases to slit the canaliculus completely into the sac, but it must be incised so far as is necessary to bring the slit part in contact with the eye.

Fig. 311. — Nettleship's canaliculus dilator.

Slitting the lower canaliculus is performed as follows. Instruments required: Nettleship's dilator (Fig. 311), canaliculus knife. The best form of canaliculus knife is the modification of Weber's in which the probe point is straight, not curved forwards as in the original instrument (Fig. 312). The eye should be cocained, though this will not eliminate all pain. The surgeon stands behind the patient. In operating upon the right side he everts the lower lid with his left thumb. With the right hand he inserts the point of Nettleship's dilator into the punctum, passing it directly downwards as far as it will go easily, then rotating it outwards and pushing it inwards along the canaliculus. In this manner the punctum is dilated. The knife is then taken and the probe point is passed into the punctum in the same manner, first downwards, then inwards. The back of the knife is directed forwards and slightly downwards. In this manner, as the knife is pushed inward, the posterior wall of the canaliculus is incised. In bad cases of eversion or when it is desired to slit the canaliculus completely into the sac it is necessary to rotate the handle of the knife upwards and towards the nose as soon as the probe point is stopped by the lacrymal bone. While this manœuvre is being performed the lid is kept stretched outwards, so that the wall of the duct is kept taut against the edge of the knife. Care must be taken that the edge of the knife, which is directed towards the globe, does not injure the eye, though there is little danger of such an accident. A Stilling's knife (Fig. 312) is useful for completing the incision into the sac.

A probe should be passed along the incised canaliculus on the day following the operation, and occasionally on succeeding days, so as to prevent closure of the incision.

Under no circumstance should the canaliculus be slit up unless it is absolutely necessary. It should never be slit up *more* than is absolutely necessary.

Occlusion of the puncta may be congenital, which is extremely rare, or cicatricial. Epiphora is caused. These cases are very difficult to treat. An endeavour should be made to slit up the occluded punctum -not the whole canaliculus. On inspection no trace of the punctum may be visible, but it is rare that some evidence of its presence cannot be seen on minute examination of the normal site with a loupe. The point of the dilator is inserted at this site, and may succeed in opening up the punctum sufficiently to admit the probe point of the canaliculus knife. There is usually no difficulty in knowing when the knife is in the duct, as it passes on in the proper direction quite easily. If this method fails to permit an entrance the canaliculus may be cut across vertically. When bleeding has stopped the inner cut end is examined with a loupe, and the probe point of the knife is inserted into it. If this also fails, and the upper punctum is patent, an attempt may be made to pass a small curved probe by the upper punctum into the sac and out into the lower canaliculus. This procedure requires an anæsthetic.

FIG. 312. — Tweedy's canaliculus knife.

Occlusion of the canaliculus may be due to a scar (*vide* p. 584) or to a foreign body. Of the latter an eyelash is the commonest, less frequent a "concretion." An eyelash usually projects somewhat from the punctum, and is easily removed with forceps. Concretions are masses of the mycelium of a fungus, usually a streptothrix. They are removed by dilating the canaliculus and injecting 10 per cent. protargol.

FIG. 313. — Stilling's canaliculus knife.

Congenital anomalies of the puncta and canaliculi are occasionally met with. The puncta may

be absent or constricted ; there may be two puncta in a lid, generally opening into the same canaliculus. Sometimes a groove is found instead of a canaliculus.

Dacryocystitis or inflammation of the lacrymal sac is not uncommon, especially among the lower classes. It is generally chronic. There is epiphora, aggravated by exposure to wind, &c. Usually there is swelling at the site of the sac. Often the caruncle and neighbouring parts of the conjunctiva are inflamed. On pressure over the sac fluid regurgitates through the puncta, or more rarely passes down into the nose. The fluid may be tears, mucus, or muco-pus ; the swelling is often called a *mucocele.* Bacteriological examination of the fluid demonstrates the presence of an extraordinary number of bacteria—staphylococci, pneumococci, streptococci, &c. Of these the pneumococcus is very frequently present in virulent form. This fact is of supreme importance, since it explains the frequency with which hypopyon ulcer arises in these cases, and the danger of panophthalmitis if any intra-ocular operation is undertaken. Dacryocystitis is a constant menace to the eye, since minute abrasions of the cornea are of almost daily occurrence, and such an abrasion is liable at any moment to become infected and give rise to an hypopyon ulcer.



Fig. 314.—Couper's lacrymal probes.

Chronic dacryocystitis is commonly attributed to the effects of stricture of the nasal duct. It is by no means certain that the stricture is primary in all these cases ; it is not unlikely that it sometimes results from the inflammation of the sac or from the treatment applied to remedy the dacryocystitis. It might be anticipated that the infection was frequently derived from the nose in cases of eczema, &c., but investigation tends to negative this conjecture. There are, however, many undoubted cases in which intractable dacryocystitis has been cured by treatment of a coincident nasal inflammation. Obstruction to the lower end

of the nasal duct may be caused by the pressure of nasal polypi, an hypertrophied inferior turbinate bone, extreme deviation of the septum, and so on.

Untreated chronic dacryocystitis never undergoes spontaneous resolution. The condition tends to progress, the walls of the sac ultimately become atonic, the contents never being evacuated except by external pressure. In any case an acute inflammation may arise, a lacrymal abscess being formed. This sequel may be caused by treatment, an abrasion of the epithelial lining leading to infection of the pericystic tissues.



Fig. 315. — Lüer's syringe, with nozzle, for syringing the lacrymal passages.

The patients are usually elderly, and such as are exposed to dirt in the course of their daily occupations. Want of personal cleanliness is probably an important factor. Dacryocystitis may, however, occur in the new-born. In these cases it is generally due to adhesion of the epithelial lining, or to imperfect canalisation of the epithelial cord in which the nasal duct is formed. The careful passage of a small probe *once* will cure these cases. Occasionally dacryocystitis in babies is extremely intractable : I am of the opinion that most of these cases are tuberculous or syphilitic, usually originating in caries of the surrounding bones.

Tubercle of the lacrymal sac also occurs in adults as a rare form of dacryocystitis.

Treatment.—In the new-born a simple boric lotion should be ordered, and minute directions should be given for expressing the contents of the sac, which should be done very frequently. Many cases will be cured by this treatment. If it fails after a fortnight, an anæsthetic should be given and a small probe passed down the nasal duct, the greatest care being exercised to avoid injuring the walls of the duct. It is unnecessary to slit up the canaliculus. The punctum and canaliculus are dilated with a Nettleship's dilator. A small probe (No. 1 or 2) is inserted vertically downwards into the canaliculus, then passed gently but firmly inwards until the point is felt against the lacrymal

bone. The probe is then rotated upwards and towards the middle line, and pushed down the nasal duct until it touches the floor of the nose. It should be remembered that the duct is short in the new-born. The force required is quite slight if rightly applied in the line of the duct (*vide* p. 590). Since much harm may be done by bad probing, these cases should be treated by an expert.

In adults the conjunctival sac should be cocained. The punctum is dilated and the sac syringed out with a lacrymal syringe. I prefer the glass syringes of Lüer (Fig. 315). A moderately fine straight nozzle should be used. The point is inserted into the canaliculus; it need not pass into the sac. Two or three syringefuls of boric lotion are passed. Probably the whole of the fluid will regurgitate through the upper canaliculus. The operation should be repeated every day for a fortnight or longer. In the majority of cases the fluid will



Fig. 316.—Müller's retractor.

pass freely down into the nose in a few days. When this occurs the syringing should be repeated at constantly increasing intervals. A great number of previously untreated cases can be cured in this manner. The patient should be told to squeeze out the contents of the sac frequently in the intervals between syringing.

The rationale of this treatment depends upon the fact that the walls of the sac and upper part of the duct are inflamed. The swollen mucous membrane prevents the fluid in the sac from passing into the nose. The treatment reduces the swelling and restores the communication. If it is unsuccessful after trial for a week or fortnight protargol (10 to 20 per cent.) should be used for syringing occasionally instead of boric lotion.

If syringing fails the condition of the nasal fossæ should be thoroughly investigated by an expert, and any pathological condition likely to cause inflammation or obstruction of the nasal duct treated.

If no cause is discovered in the nose it is best to excise the lacrymal sac. This radical operation, properly performed, completely removes the disease, with a minimum of inconvenience to the patient.

Many surgeons, perhaps the majority, treat chronic dacryocystitis by probing. The canaliculus is slit up, and probes of increasing calibre are passed down the nasal duct into the nose (Fig. 314). The objections to this method of treatment are—(1) it is impossible to probe the swollen and inflamed duct without injuring the walls; (2) such injury may lead to infection of the surrounding tissues and an acute cellulitis; (3) in any case healing of the abrasions is accompanied by the formation of connective tissue, which contracts when it organises and leads to fibrous stricture instead of obstruction by swollen mucous membrane; (4) probing is always painful, and when once begun has to be continued for a prolonged period; (5) most cases are alleviated only temporarily, fresh courses of probing being required at intervals. Protargol should never be injected immediately after slitting up the canaliculus or probing. If an abrasion of the mucous membrane has been caused the protargol may be injected into the subcutaneous tissues and violent cellulitis follows. Orbital cellulitis and atrophy of the optic nerve have been known to result from neglect of this rule. Even if these serious results do not follow there is unsightly and permanent staining of the skin. These facts are proof of the injury done to the mucous membrane by probing.

FIG. 317.—Lang's dissector and scoop.

Some surgeons introduce styles into the nasal duct in order to keep it patent. After obtaining a passage with probes a lead or silver style is passed into the duct, the upper end being bent outwards so that it lies upon the lid or cheek, thus preventing the style from being lost. This treatment has all the objections which have been adduced against probing. Occasionally it is useful

in intractable cases in young children, in whom it is difficult to extirpate the lacrymal sac efficiently. Even in such cases it is better to extirpate the sac as well as possible and pack the cavity with antiseptic gauze, so that the wound heals from the bottom.

Excision of the lacrymal sac is performed as follows. Instruments required: small scalpel, dissecting forceps, toothed (fixation) forceps, retractors (preferably Müller's) (Fig. 316),



Fig. 318.—Incision for excision of the lacrymal sac. The broken lines indicate the bones and orbital margin; also the limits of the conjunctival sac.

small blunt-pointed (tenotomy) scissors, blunt dissector and small sharp spoon (Fig. 317).

The operation can be performed with local anæsthesia; the best solution to employ is novocain (1 per cent.) and suprarenin (0·00125 per cent.), of which up to two drachms may be injected around sac. The lids are held stretched outwards by a finger applied near the outer canthus. This throws the internal palpebral ligament into prominence. A curved incision is made, commencing just above the ligament and extending downwards and slightly outwards to the inferior margin of the orbit and for a short distance along it (Figs. 318—319).

The incision should be only 3–4 mm. internal to the inner canthus, otherwise there is danger of wounding the angular

vein or its larger branches. The internal palpebral ligament should be severed and the sac opened, but the incision should not be so deep in its inferior part. Müller's retractors are then inserted, and any hæmorrhage is stopped by firm pressure with a small swab dipped in adrenalin (1 in 1,000). The mucous membrane can now be seen protruding from the opening made in the sac. It is seized on the inner side with fixation forceps and separated from the surrounding tissues by the blunt dissector. After the inner wall has been freed the fundus and outer wall are similarly treated. The sac is then



Fig. 319.—Excision of the lacrymal sac. Müller's retractor *in situ*.

pulled forwards and freed behind. Finally the sac is pulled well up, so as to drag out as much of the mucous lining of the nasal duct as possible, and is removed by a snip of the scissors, applied as close as possible to the bone. The wound is carefully inspected to see that all of the mucous tissue is excised. The upper part of the nasal duct is then thoroughly scraped out with the sharp spoon. The wound is completely sutured. A small spherical pad is applied, and over this an ordinary pressure pad and bandage. The wound generally heals by first intention ; it should not be pressed with the finger to see if there is any regurgitation during the first few days, as it is easily reopened.

Difficulty is experienced in excising contracted sacs. It

may then be found that the incision has not opened the sac, which is firmly adherent to the surrounding tissues. The best guide to the position of the sac in these cases is the crest of the lacrymal bone, which can be easily felt with the finger-nail. The sac must then be separated from the bone on the inner side by scalpel or scissors. The fundus, outer wall, and posterior attachment are successively dissected free. There is usually much more hæmorrhage when this method has to be adopted. It is best controlled by firm pressure with small swabs soaked in adrenalin.

There is no objection to removing both lacrymal sacs at the same operation if there is bilateral mucocele. When the operation is satisfactorily performed there is no regurgitation on pressure over the scar. If after a week or two there is still some regurgitation, part of the mucous membrane has been left behind, and the operation must be repeated. Usually it is the fundus of the sac which has been left. This lies above the palpebral ligament; hence the advisability of dividing the ligament in order that a good view may be obtained. Not the slightest evil results from this procedure. Sometimes regurgitation is due to leaving the mucous membrane of the upper part of the nasal duct; it will not occur if the duct is well curetted.

In all cases of cataract in which there is a mucocele the lacrymal sac should be excised as a preliminary to extraction. Only some weeks after this operation, when there is no trace of regurgitation, is it permissible to proceed with the extraction. A more difficult problem is the presence of a mucocele in a case of acute glaucoma. Here immediate iridectomy is indicated and admits of no delay. In these cases the sac must be completely isolated from the conjunctival sac. This is best effected by passing a ligature round each canaliculus and tying it firmly. Some surgeons cauterise the punctum, thus sealing it up with a cicatrix. Either procedure may be followed by the development of a lacrymal abscess, but in the meantime the iridectomy wound has probably healed, and in any case the pus is evacuated through the skin and not into the conjunctival sac.

Epiphora usually persists for a time after excision of the sac. This is due to the chronic conjunctivitis set up by the mucocele. Under normal conditions the tears evaporate from the surface of the globe, and only when there is excessive secretion from reflex irritation or emotional causes do tears pass down the nasal duct. Hence the post-operative epiphora should be treated by astringent lotions, &c.; in no case is it

necessary in my experience to remove the lacrymal gland, as has been advocated. Epiphora will, however, still occur on exposure to wind, &c., and this cannot be avoided.

Lacrymal Abscess may be due to acute dacryocystitis or to suppuration starting in the pericystic tissues. The skin over the sac becomes red and swollen. The redness and swelling rapidly extend to the lower lid and upper part of the cheek, so that the condition may be easily mistaken for erysipelas. There is severe pain, and often some fever. The abscess usually points below and to the outer side of the sac owing to gravitation of the pus to margin of the orbit. If it opens spontaneously pus continues to be discharged for some time, and a permanent fistula is likely to result.

Treatment.—If seen at the commencement of the process an attempt should be made to prevent the abscess pointing on the face by freely slitting up the lower canaliculus well into the sac. Hot fomentations are then applied frequently.

If the abscess is fully developed it should be freely opened at the site where it is pointing. The pus is squeezed out, and the cavity is thoroughly plugged with cyanide gauze. A hot fomentation is applied and repeated frequently. The plugging is repeated daily, so that the cavity granulates from the bottom. If the discharge continues for a long period the cavity should be well curetted and again plugged. Usually the epithelial lining of the sac is destroyed by the purulent inflammation, the sac is permanently destroyed, and the cure is complete. In other cases some of the mucous membrane escapes destruction, and a fistula may follow. It may sometimes be closed by cauterising the edges with the galvanocautery, but it is better to reopen the sac along a director introduced through the fistula and extirpate the remnants, but this procedure should not be adopted until several weeks after the acute inflammation has subsided.

Stricture of the Nasal Duct has already been referred to incidentally. It is probable that most intractable fibrous strictures are caused by probing, though it cannot be asseverated that they may not arise spontaneously as the result of destruction of the epithelium by extension of inflammation from the nose or lacrymal sac. Occasionally bony strictures occur, usually caused by caries.

Treatment.—The usual treatment of stricture of the nasal duct is dilatation with probes. The objections to this treatment have been mentioned. I advocate excision of the lacrymal sac in these cases.

CHAPTER XXXIII.

DISEASES OF THE ORBIT.

It is unnecessary to describe the anatomy of the orbit and its contents here. The student is recommended to revise his knowledge of the subject, paying special attention to the relations of the nasal cavities and their accessory sinuses, and to the communications with the interior of the cranial cavity by way of the optic foramen and sphenoidal fissure. The intimate adhesion of the dural sheath of the optic nerve to the walls of the optic foramen is of great pathological importance, and the relations of the intraorbital to the intracranial circulation must be thoroughly appreciated. The eye is slung in position in the orbit by fascia, one sheet of which, Tenon's capsule, forms a socket in which the globe moves. This, with the sclerotic, forms a lymphatic space, lined completely with endothelium. The extrinsic muscles of the eye do not perforate this capsule, but invaginate it, the fascia being reflected from their surfaces.

The normal position of the eye is such that a straight-edge applied vertically to the middle of the upper and lower margins of the orbit just touches the closed lids over the apex of the cornea. There are individual variations which are of no pathological importance when symmetrical; in all cases of doubt the two sides should be compared.

Abnormal protrusion of the globe is called *exophthalmos* (more accurately, *exophthalmia*) or *proptosis*. It is much commoner than abnormal retraction or *enophthalmos*. The former condition is due to many causes, among which increase in the orbital contents and loss of tone of the extrinsic ocular muscles are the most important. Slight prominence of the eyes accompanies high myopia, paralysis of the extrinsic muscles, stimulation of Müller's muscle by cocain, and as an idiosyncrasy, especially in very obese people. Unilateral exophthalmos occurs in orbital cellulitis from any cause, thrombosis of the orbital veins with or without implication of the cavernous sinus, arterio-venous aneurysm, tumours of the orbit and its contents, and orbital hæmorrhage or

emphysema. Bilateral proptosis occurs in exophthalmic goître, the later stages of thrombosis of the cavernous sinus, empyema of the accessory sinuses of the nose, symmetrical orbital tumours (lymphoma, pseudoleukæmia), and as a result of diminished orbital volume in oxycephaly or "tower-skull" and leontiasis ossium. Enophthalmos is generally due to severe injury in which the orbital bones are fractured, or to orbital cellulitis with mechanical retraction by fibrous tissue. Slight degrees of exophthalmos or enophthalmos are best diagnosed by the test with a straight-edge. Accurate estimates of the amount can be obtained only by special mechanical devices (exophthalmometers). A convenient test is the following: The patient is seated, the surgeon standing behind him. The surgeon holds the patient's head in such a manner that he looks straight down the nose. He then rotates the head backwards until he can just see the apex of one cornea. If he can see more of the other cornea, that eye is relatively proptosed.

Orbital Inflammation.

Periostitis is not uncommon, particularly affecting the margin. It is most often due to injuries, extension of inflammation from neighbouring parts, tubercle or syphilis. Tuberculous periostitis is most frequent in children, syphilitic in adults: in the former, caries of the bone results; the latter is gummatous. In traumatic cases the margin is naturally most affected, but a traumatic element is often an exciting cause in the other cases, so that in them also the margin most frequently suffers.

When situated at the margin, the inflamed part is swollen and tender; the swelling is intimately connected with the bone, so that it cannot be moved over it. Syphilitic cases usually respond well to treatment. The other types generally go on to suppuration. An abscess is formed, and when it discharges or is opened rough bone can be felt with a probe. In tuberculous cases particularly a fistula may result, the edges of the aperture being bound down to the bone, so that a depressed cicatrix is formed. The fistula remains open until all the necrosed bone is extruded. The cicatrisation may lead to displacement of the lid—ectropion, lagophthalmia, and so on.

Periostitis of the deeper parts of the orbit causes less definite signs. There is more pain of a deep-seated character. There may be proptosis with deviation in the direction of the

eye. In the case of gumma the roof of the orbit is generally involved, the deviation of the eye is downwards, and there is rapid loss of movement owing to involvement of the extrinsic muscles. There is severe supraorbital neuralgia, which is worse at night. Often the true nature of the disease is only discovered by an exploratory operation, or by the evacuation of pus. The case may present all the features of orbital cellulitis (*q.v.*). If the roof of the orbit is involved, the pus may discharge into the cranial cavity, life being endangered by meningitis or cerebral abscess.

Treatment is determined by the ætiological factor. In syphilitic cases, mercury and iodide of potassium are pushed rapidly. In traumatic cases, if suppuration supervenes, the abscess is opened, hot fomentations being applied previously if necessary. In tuberculous cases an incision should be made early and any carious bone removed, care being taken not to encroach upon the cranial cavity.

In deep-seated periostitis an exploratory operation may be necessary, and should not be too long delayed. An incision is made through the skin at the margin of the orbit, the knife being passed cautiously deep into the orbit along the wall. The site of the incision is determined by the signs present. Sinus forceps are passed down the track of the wound and opened. The greatest care should be exercised to avoid unnecessary damage to the orbital contents, and this is best accomplished by keeping closely to the bony walls. Special care must be taken not to injure the pulley of the superior oblique. If pus is found, a small drainage tube or a long strip of cyanide gauze is inserted. In periostitis of the inner wall, the bone may be extensively diseased. Severe operations, involving the opening of the frontal or ethmoid sinuses, may be essential, with or without drainage through the nose. These cases often do remarkably well. They are usually tuberculous, and occur most frequently in children. Exploration of the orbit in children is much more difficult than in adults. The eye is relatively much larger in comparison with the size of the orbit, so that there is very little room between the globe and the orbital wall. In some cases it may be advisable to perform Krönlein's operation (*vide* p. 613).

Orbital Cellulitis is purulent inflammation of the cellular tissue of the orbit. It is due to deep injuries, especially those with retained foreign body, or septic operations, *e.g.*, enucleation of the eyeball; extension of inflammation from neighbouring parts, especially the nasal sinuses and teeth; facial

erysipelas; metastasis in pyæmia, meningitis, infective fevers, &c.

There is great swelling of the lids, with chemosis. The eye is proptosed, and its mobility impaired. Pain is severe, increased by movement of the eye or pressure upon it. Fever is present, and cerebral symptoms may arise. Movement of the eye is painful, and there may be diplopia owing to limitation of movement. Vision may not be affected, or it may be reduced owing to retrobulbar neuritis. The fundus is difficult to examine; it may be normal or show engorgement of the veins and optic neuritis, passing later into optic atrophy. An abscess is formed which usually points somewhere in the skin of the lids near the orbital margin, or it may empty into the fornix conjunctivæ. Panophthalmitis may supervene. There is grave danger of extension to the meninges and brain, leading to a fatal issue from purulent meningitis or cerebral abscess. Thrombosis of the cavernous sinus (*q.v.*) may result from orbital cellulitis, and is always difficult to diagnose from it.

Treatment.—Hot fomentations are applied, but must not be relied upon too long. An early incision as in orbital periostitis (*q.v.*) is imperative. Even if pus is not reached, the tension is relieved and a track is prepared for its evacuation. Several incisions along the walls of the orbit are indicated if there is no sufficient guide to the position of the focus. If the source of infection is obscure, the nose and other likely seats must be investigated, and the primary focus treated.

Thrombosis of the Cavernous Sinus may be due to extension of thrombosis from various sources.

The anatomy of the venous channels which communicate with the cavernous sinus is of prime importance for the comprehension of thrombosis affecting it. The superior and inferior ophthalmic veins enter it in front and the superior and inferior petrosal sinuses leave it behind. It communicates directly with the pterygoid plexus through the middle meningeal veins and the vein of Vesalius, and indirectly through a communicating vein from the inferior ophthalmic to the pterygoid plexus. The anastomoses of the ophthalmic veins with the frontal and angular open up a communication with the face. Labyrinthine veins opening into the inferior petrosal sinus afford a communication with the middle ear. Numerous tributaries throw it into direct or indirect communication with most parts of the cerebrum. The mastoid emissary vein places the sinus in communication with the sub-

cutaneous tissues behind the ear through the sigmoid sinus and superior petrosal sinus: it is this communication which is of great diagnostic importance, since swelling behind the ear may decide the question of thrombosis in each direction along them. The sinus of one side communicates with that of the other by two (or sometimes three) transverse sinuses which surround the pituitary body.

Infection may occur viâ the orbital veins *e.g.*, erysipelas of the face, orbital cellulitis, and mouth and pharynx; from the ear, nose and accessory sinuses; or as a metastasis in infectious diseases or septic conditions.

The patient presents almost the same symptoms and signs as in orbital cellulitis. If in addition there is œdema in the mastoid region behind the ear the diagnosis is certain, for this



FIG. 320.—Tributaries of the cavernous sinus.

is due to thrombosis of the emissary vein. A further point of diagnostic importance is transference of the symptoms to the opposite eye, which occurs in 50 per cent. of cases, whereas bilateral orbital cellulitis is very rare. The first sign is paralysis of the opposite external rectus, and this sign should be carefully watched for in any suspicious case of inflammatory unilateral exophthalmos. It must be remembered, however, that thrombosis of the sinus may be a complication of cellulitis.

There is severe supraorbital pain, owing to implication of the branches of the ophthalmic division of the fifth nerve, and the motor ocular nerves are paresed or paralysed. In the later stages the eye is immobile, the pupil dilated, and the cornea anæsthetic. Proptosis occurs in nearly all cases, but is of late onset in those of otitic origin.

It is commonly stated that the retinal veins are greatly engorged, but in many cases this is certainly not true. When it occurs it is usually accompanied by pronounced papillitis, and both signs indicate extensive implication of the orbital veins and tissues. Simultaneous thrombosis of both cavernous sinuses, with proptosis and papillitis, occurs in diseases of the sphenoidal sinuses. Typical papillœdema is commonest in otitic cases, and indicates meningitis or cerebral abscess: it is bilateral, and more pronounced on the side of the aural lesion.

Thrombosis of the cavernous sinus is accompanied by



FIG. 321.—Tributaries of the cavernous sinus.

serious general symptoms—septic temperature, rigors, vomiting, and severe cerebral symptoms. The patient always dies.

Tenonitis is inflammation of Tenon's capsule. It may be serous or purulent. There is exophthalmos straight forwards, with limitation of movement of the globe and pain on attempted movement. There may be some œdema of the lids, and chemosis. It may occur in severe iridocyclitis, and is constant in panophthalmitis. It may also follow tenotomy, &c. Simple serous tenonitis is rare, and has been attributed to influenza, gout, rheumatism, &c.

Treatment consists in the application of hot fomentations, and the evacuation of pus, if it forms. When it occurs as part of panophthalmitis, &c., it requires no special treatment.

Distension of the Accessory Sinuses of the Nose.

The accessory sinuses of the nose—the frontal, ethmoidal and sphenoidal sinuses, and the antrum of the superior maxilla are separated from the orbit only by thin plates of bone. The orifices which form the communication between these cavities and the nose are liable to become occluded by catarrh, polypi, &c. The normal sero-mucous discharge is thus unable to drain into the nose. The cavities become distended with fluid, and owing to the presence of pyogenic organisms pus may be formed. The treatment of the conditions thus set up

Fig. 322.—Distension of the frontal sinus.

cannot be considered part of the functions of the ophthalmic surgeon, but he must be prepared to diagnose them since they not infrequently appear for the first time in the ophthalmic clinic. This is particularly the case in distension of the frontal, ethmoidal and sphenoidal sinuses. Of these the frontal sinus suffers most often.

Distension or empyema of the frontal sinus causes bulging at the upper and inner part of the orbit (Fig. 322). There may be some proptosis and displacement of the eyeball downwards and outwards, but these features are more marked when the ethmoidal sinus is involved. Œdema of the upper lid or slight ptosis may be the only external sign. There is considerable pain and tenderness, with severe headache. There is often discharge from the nostril of the same side, or manifest disease

of the nasal cavities. Owing to erosion of the walls of the sinus the fluid may extend under the periosteum, causing bulging into the posterior part of the orbit. It may escape into the opposite sinus and through the infundibulum of that side; or it may rupture into the orbit, through the skin, forming a sinus, or even into the cranial cavity. Orbital cellulitis may be set up in this manner.

The frontal sinus is not developed until about the sixth year; the disease occurs most commonly between twenty-five and thirty, and more cases occur in men than in women.

Treatment of distended frontal sinus consists in providing free discharge of the contents through the nose. In most cases a radical cure is effected only by laying open the sinus, scraping away completely the diseased mucous membrane, and passing a drain down into the nose. The disease is tedious to treat and much disfigurement may follow.

Distension of the ethmoidal cells by polypi, new growths or inflammatory products may also cause bulging into the orbit and displacement of the globe. Diplopia, chemosis, venous engorgement and ptosis may be caused. Ethmoiditis is usually associated with nasal discharge. It may give rise to orbital cellulitis, or in less severe cases to retrobulbar neuritis. The latter is probably more commonly associated with inflammation and distension of the sphenoidal cells, which lie in close proximity to the optic nerve, being sometimes separated from it by a very thin lamina of bone. In doubtful cases help may be afforded by a skiagram. It has already been mentioned that the accessory sinuses of the nose are not infrequently the foci from which toxins are disseminated, leading to iridocyclitis and other metastatic septic processes in the eye (*vide* pp. 259, 318).

Injuries of the Orbit.

Injuries to the soft parts usually arise from penetration of a foreign body, which may be retained. The lids and eyeball are frequently implicated. The signs depend upon the particular structures injured. In most cases there is considerable hæmorrhage; as the blood does not find a ready exit exophthalmos may result. Extravasation of blood under the conjunctiva and into the lids is common. Hæmorrhage may result from pressure with forceps at birth. It also occurs in some cases of fracture of the base of the skull. Paralysis of extrinsic muscles may be due to direct injury or to injury of

the motor nerves. The optic nerve may be severed or retrobulbar neuritis may ensue; in either case atrophy, involving the optic disc, follows (*vide* p. 365). The eyeball may be perforated or contused (*vide* p. 397) or dislocated *en masse*. Dislocation forwards between the lids occurs most often when the blow is directed from the outer side, where the orbital margin affords least protection. Insane patients sometimes enucleate their eyes by gouging them out with their fingers. Sight is not necessarily lost after dislocation forwards. Retained foreign bodies are extremely liable to set up suppuration and orbital cellulitis (*q.v.*).

Injuries to the bone most commonly affect the margin of the orbit. Fractures in this locality are easy to diagnose from the unevenness of the margin, sensitiveness to pressure, and sometimes crepitation. Emphysema (*q.v.*) may occur. The soft parts may be injured by splinters of fractured bones. Deep fractures may be caused by penetrating wounds or by severe contusions, falls, &c. Fracture of the base of the skull may involve one or both optic foramina, in which case the optic nerve is often severed, lacerated, or compressed by clot. Blindness without ophthalmoscopic signs may be caused in this manner; atrophy of the disc follows in three to six weeks (*vide* p. 365).

Treatment.—If there is a wound it must be cleansed, and, if necessary, probed. The treatment of a retained foreign body depends upon its situation and the probability of suppuration occurring. If the foreign body cannot be extracted with ease a skiagram should be taken. If the position is such that very serious manipulation would be requisite for removal, and if there is evidence that the substance is aseptic, expectant treatment may be adopted. If suppuration occurs the foreign body must be removed and the case treated as one of orbital cellulitis (*q.v.*).

Tumours of the Orbit.

Orbital tumours are rare. Benign growths include dermoid cyst, dermo-lipoma (*vide* p. 185), angioma, osteoma (Fig. 323), plexiform neuroma (*vide* p. 588), meningo-encephalocele. Of these, dermoid cysts appear as swellings under the lid, usually at the upper and outer angle; they contain sebaceous material derived from sebaceous glands in the walls, which are lined with epithelium and possess hair follicles. Clinically they may be mistaken for meningo-encephaloceles, which usually occur at the upper and inner angle, where there are most

sutures between bones. In the latter (1) the tumour is immovably attached to the bones; (2) the hole in the bone may be palpable; (3) pulsation, synchronous with respiration and the pulse and increasing on straining, can be seen); (4) pressure may cause diminution in size due to fluid being pressed back into the cranium; (5) exploratory puncture produces clear fluid with the characteristics of cerebro-spinal fluid. Osteomata start usually from the frontal bone (Fig. 323); they are intensely hard and often large, producing great displacement of the globe.

Malignant tumours of the orbit are usually sarcomata, though carcinoma derived from the lacrymal gland (*vide* p. 591) or by extension from the nasal mucous membrane also occurs. All types of sarcoma, including endothelioma, may occur. The small round-celled growths include cases of lymphoma, leukæmic tumours, chloroma, &c.

FIG. 323.—Orbital osteoma. (Tweedy.)

Primary tumours of the optic nerve manifest themselves clinically as orbital tumours. They are rare, and are found on anatomical examination to consist of two groups—intra- and extradural. Of these the former are more numerous (Fig. 324). They rarely spring from the true nervous tissue (gliomata, resembling cerebral glioma, not retinal glioma), but usually originate in the connective tissue septa derived from the pia mater and from the arachnoid sheath. They often contain tissue of mucoid type, and are hence described as myxosarcoma, &c. Probably most are endotheliomata which have undergone degenerative changes. Extradural tumours spring from the dural sheath and are also probably endotheliomata (Fig. 325). All these optic nerve tumours are most common in children and are locally malignant, but show little or no tendency to metastasis. They may kill the patient by intracranial extension.

Most orbital tumours cause proptosis, which is very rarely straight forwards except in the case of optic nerve tumours. This is an important diagnostic feature. The exophthalmos increases slowly and gradually, and is nearly always unilateral : in rare cases of lymphoma it is bilateral. The mobility of the eyeball is impaired in the direction towards the position of the tumour. There is usually diplopia from this cause. Papillitis may be present, especially with optic nerve tumours. Optic atrophy from pressure on the nerve is common in the other



Fig. 324.—Intradural tumour of the optic nerve. (Mayou.) Longitudinal section, section stained by Weigert-Pal, to show distribution of the nerve fibres.

forms. The tumour may be palpable by the finger pushed back between the globe and the orbital wall. The lymphatic glands are seldom affected.

Careful examination of neighbouring parts nose, antrum mouth (especially the naso-pharynx, and line of the teeth) must be made to determine whether the invasion of the orbit is secondary or whether the growth is primarily orbital. In doubtful cases an X-ray examination should be made.

Treatment.—An exploratory operation and removal of a portion of the growth for microscopic examination may be a necessary preliminary to radical treatment. It may be

feasible to remove dermoid cysts and some other benign tumours without injury to the globe, though its mobility is likely to be impaired in extensive operations. As already mentioned, many malignant orbital growths show little tendency to metastasis, so that their treatment may be more conservative than is usual in other parts of the body. Thus it is possible in some cases of optic nerve tumour to remove the growth while retaining the eyeball. It can be effected by *Krönlein's operation*, which is also of utility as an exploratory procedure in some cases. In it a semilunar incision is made vertically just outside the outer canthus, the convexity being directed forwards. The bone is chiselled through at the upper and lower outer angles of the orbit, and bone, muscle and skin are reflected backwards in one flap. The posterior part of the orbit is thus exposed in a manner which is impossible by any other method. The greatest care must be exercised that infection does not occur, since the spongy bone is laid open, and there is also danger of meningitis.



FIG. 325.—Extradural tumour of the optic nerve (× 1·4).

In the case of more malignant types of tumour their complete removal is imperative at all costs, and the eye, which may be quite normal, may have to be sacrificed. In these cases, as well as in recurrence or in orbital extension of malignant intraocular growths (glioma of the retina, sarcoma of the uveal tract), it may be necessary to remove the whole contents of the orbit.

In *exenteration of the orbit* the lids may be retained if they are not implicated in the growth, but the free margins, carrying the cilia, should always be removed. If it is not done the lashes are troublesome when the lids become retracted into the orbit, as invariably follows. If the lids are removed the incision is carried through the skin at the margin of the

orbit in its whole circumference. The orbital contents are separated from the walls by a periosteal elevator, so that they remain attached only at the apex of the orbit. The pedicle is then severed with strong scissors, hæmorrhage being controlled by pressure. The orbit is packed with gauze and firm pressure is kept up. At a later stage it may be advisable to apply Tiersch grafts to the walls, since the lids and conjunctiva never afford sufficient epithelial covering, and the extension of the epithelium over so large a surface is a tedious process.

Special Forms of Exophthalmos.

Exophthalmic Goitre (*Syns.—Graves's or Basedow's Disease*) is one of the commonest causes of exophthalmos (Fig. 326). The symptom complex includes, besides proptosis, enlargement of the thyroid gland, tachycardia and muscular tremors. The proptosis is almost always bilateral and may be extreme, leading to lagophthalmia and its deleterious consequences (*vide* p. 214). There is a peculiar stare, with retraction of the upper eyelid, so that there is an unnatural degree of separation between the margin of the two lids (Dalrymple's sign). Normally, when vision is directed downwards, the upper lid moves concordantly with it. In this disease the upper lid follows tardily or not at all (von Graefe's sign): this symptom is not always present and may occur in other forms of exophthalmos. There is diminished frequency of winking and imperfect closure of the lids during the act (Stellwag's sign). There may be imperfect power of convergence (Möbius' sign), and often the skin of the eyelids shows pigmentation. Ophthalmoscopically veins and arteries may be somewhat distended, but specific signs are absent. One or more of the cardinal symptoms may be absent. Paralysis of extrinsic ocular muscles, usually the external rectus, sometimes occurs. The cause of the disease is unknown. The most probable theory attributes it to over-production or other disorder of thyroid secretion, but it has also been attributed to disease of the cervical sympathetic ganglia, of the central nervous system, &c.

It is necessary for the ophthalmic surgeon to be able to recognise the disease. Further details of its usual course and treatment must be sought in medical text-books.

Pulsating Exophthalmos is generally due to arterio-venous aneurysm, the communication taking place between the internal cartoid artery and the cavernous sinus. The eyeball

is protruded and the blood vessels of the conjunctiva and lids are widely dilated. The angular vein and its branches near the inner canthus are very prominent, and they can be seen, or more easily felt, to pulsate synchronously with the arterial pulse, since, owing to the arterio-venous communication, they are under arterial pressure. The patient complains of continual rumbling, as of a waterfall, and this can be heard on auscultation over the eye or orbit by the surgeon. The proptosis is diminished by steady pressure on the globe, and may be diminished or abrogated by pressure on the common carotid artery of the same side, or sometimes only by pressure on the carotid of the

FIG. 326.—Exophthalmic goître.

opposite side. Ophthalmoscopically the veins of the retina are greatly distended : there may be papillitis with defective vision, which may amount to complete blindness. There is often considerable pain from stretching of the branches of the fifth nerve.

The cause of the arterio-venous aneurysm is usually a severe blow or fall upon the head, but probably in all cases the walls of the artery are already degenerated. It may occur from syphilitic or other arteriosclerosis, without discoverable traumatism. The exophthalmos in rare cases subsides spontaneously. More commonly it increases, and may end in hæmorrhage or death from cerebral causes.

Treatment.—Continuous pressure applied to the carotid

artery which stops the pulsation usually fails to effect a cure. Ligature of the carotid has been more successful, but recurrence of pulsation not infrequently occurs. In these cases the opposite carotid may also be tied, but this should not be done for some weeks after the first operation, owing to risk to life from cerebral anæmia. This procedure also may fail to relieve the condition, and in these cases the distended veins have been dissected out, an operation of considerable danger.

Intermittent Exophthalmos occasionally occurs, generally when the head is depressed, enophthalmos not infrequently being present in the erect position. The proptosis is increased by pressure on the corresponding jugular vein. It is ascribed to varicosity of the orbital veins.

SECTION VIII.

PREVENTIVE OPHTHALMOLOGY.

CHAPTER XXXIV.

THE CAUSES AND PREVENTION OF BLINDNESS.

The previous chapters have dealt chiefly with the diagnosis and treatment of already established diseases of the eye. An equally important branch of medical science is concerned with the prevention of disease, and although this aspect of ophthalmology has hitherto received less attention than it merits, it ought not to be ignored by the medical student or practitioner.

The most disastrous result of ocular disease, short of the relatively rare loss of life, is blindness. A study of the causes of blindness will enable the student to form a judgment as to the comparative danger of various ocular diseases.

The term "blindness" implies inability to perceive light; but it is obvious that many people who yet retain some slight degree of visual capacity are helpless from the economic standpoint. The Advisory Committee on the Welfare of the Blind, therefore, included among the blind all those who are "too blind to perform work for which eyesight is essential." The Register of the Blind for England and Wales, compiled on this basis, shows that there were 34,894 blind persons in those areas in 1921. Statistics as to age incidence are somewhat unreliable, but such as they are, they show that about 20 per cent. became blind within the first year of life, and that about 10 per cent. became blind in each decade from one year to seventy years of age.

The factors producing blindness have a different rate of incidence at different ages. Thus, in a home for blind *infants* Harman found about 50 per cent. due to ophthalmia neonatorum, 11 per cent. to intraocular inflammations, and 30 per cent. to congenital defects. Statistics of *children of school age* show that 20—30 per

cent. were blind from ophthalmia neonatorum, 10—20 per cent. from interstitial keratitis, and 15—20 per cent. had optic atrophy due to various causes, including disseminated choroiditis. The statistics for *adults* are very unreliable, owing to defective case histories, the impossibility in many cases of determining the causes of blindness from examination of the patients, and other reasons. In middle life the high incidence of ophthalmia neonatorum is still noticeable, choroiditis and optic atrophy are important, while iritis and iridocyclitis are markedly advanced in relative position. Many of these are symptomatic conditions, the underlying cause being often syphilis. Myopia is a prominent factor, accidents assume a high proportion, and glaucoma appears, increasing considerably in later life.

The importance of *ophthalmia neonatorum* as a cause of blindness is so great that it has been deemed advisable to discuss the measures for its prevention earlier in this book (*see* p. 155). While there is unfortunately no evidence of any reduction in the incidence of this disease in recent years there is some evidence of a reduction in the amount of blindness caused by it.

Syphilis, both in the congenital and in the acquired form, is responsible for a large amount of blindness. Harman found definite signs of congenital syphilis in one-third of a group of 1855 blind children, and in most of these cases congenital syphilis, was the undoubted cause of the blindness. At least 10—15 per cent. of cases of blindness in adults are probably due to syphilis, and these figures do not include cases due to vascular disease of possible syphilitic origin.

Phlyctenular keratitis was found to be the cause of blindness in 3·56 per cent. of 1855 blind children (Harman). This disease and such conditions as blepharo-conjunctivitis are largely due to insanitary conditions of life. They might probably be eliminated as causes of blindness by the adequate provision for the education and treatment of the children in special residential schools, a method which has proved eminently successful for trachomatous children. *Measles* is another not uncommon cause of blindness through corneal ulceration; it emphasises the importance of proper treatment of the eyes by the medical practitioner. *Trachoma* (*q.v.*) is, fortunately, now a rare cause of blindness in this country.

Myopia was the cause of blindness in 3 per cent. of 1855 blind children (Harman), and in 14 per cent. of 601 blind persons of all ages (Harman). These figures underestimate the serious distress and economic loss due to this cause. It is generally agreed that myopia is increased by near work (*vide* Chap. XXIV.). Special "myope" classes have been instituted for the education of short-sighted children. "It is convenient to classify myopia in two

divisions - a 'school' myopia and a 'pernicious' myopia. The former is, as a rule, low in degree, does not progress beyond a certain extent, and is but rarely associated with other ocular changes. The fact that ophthalmic surgeons recognise a school myopia is the strongest argument for its prevention by the provision of all those measures which are summed up in the term 'school hygiene'; such as good lighting in classrooms, good print in books, regulated needlework, a maximum of oral instruction, and, above all, the early correction of errors of refraction and the special supervision of those who show signs of being or becoming short-sighted. But it is the pernicious myopia which figures in these tables of blindness or partial blindness; a variety which commences at an early age, may progress rapidly, is very liable to be associated with serious intraocular disease, and therefore necessitates the provision of special methods for the safe education of the subjects of it. The cardinal aim should be to endeavour to prevent the continued development of the disease in the children affected; such prevention requires the complete elimination of any form of eye strain such as is involved in close work, and the provision of educational facilities under the best hygienic conditions, such as is aimed at in the myope classes." There is evidence that habitual close eyework, such as that of clerks, sempstresses, compositors, &c., is disastrous to the eyesight of myopes, and every effort should be made to direct these people into suitable occupations.

Glaucoma is a serious factor in the production of blindness after middle life; in a home for the aged blind the percentage of such cases was 29·82. "Cases of glaucoma frequently come in the first instance under the cognisance of the general practitioner. The acute form is apt to be mistaken for iritis, with disastrous results if the treatment appropriate to this latter disease is adopted; on the other hand, as the disease is often ushered in by severe headache and vomiting, the essential ocular condition may be overlooked, and the case regarded as a simple 'bilious attack.' The chronic form is so insidious in its onset that it may easily be overlooked." It is unfortunately by no means rare to meet with cases of acute or chronic glaucoma which have been treated with atropin or allowed to progress without proper treatment. It is of the utmost importance that the student should pay special attention to this disease, and especially to the diagnostic features which distinguish it from iritis (*vide* p. 246). If there is the slightest doubt as to the presence of glaucoma the case should be referred *immediately* to an ophthalmic surgeon.

Industrial conditions may cause blindness, either by disease or accident. The chief diseases, such as poisons (lead, derivatives of benzene, &c.), glass- and iron-workers' cataract, miners'

nystagmus, &c., have already been dealt with. Blindness due to industrial accidents is commonest among miners and in the engineering trades. Of 5,575 blind persons of all ages the blindness was due to industrial accidents in 7·2 per cent. (Scottish Register of the Blind, 1922). More striking is the enormous economic loss entailed by relatively minor accidents, such as foreign bodies in the eye (*vide* p. 409), &c. Many of these could be entirely prevented by the use of appropriate guards, screens and goggles. These matters require special attention from factory medical officers; but of far greater importance to the ordinary medical practitioner is the grave risk of blindness due to sympathetic ophthalmia produced by penetrating injuries of the eye. "The danger of the development of sympathetic ophthalmia should always be present in the mind of the medical practitioner, and all cases of penetrating wounds of the eye should be placed immediately under the observation of an ophthalmic surgeon; in no eye condition is prompt co-operation between the general practitioner and the expert more essential."

(Further information and recommendations referring to the subject of this chapter will be found in the Report of the Departmental Committee of the Ministry of Health on the Causes and Prevention of Blindness. H.M. Stationery Office, 1922.)

CHAPTER XXXV.

THE HYGIENE OF VISION.

Apart from the conditions which seriously endanger the eyesight, discussed briefly in the last chapter, there are many others which are liable to impair the efficiency of vision or the health of the individual. It is well known that the use of the eyes with uncorrected errors of refraction or muscle balance, or under unsuitable conditions of illumination, &c., cause ocular pain and discomfort (commonly known as "eye-strain"), headaches, migraine, and general malaise. More serious disorders and diseases have been attributed by some to these causes. The exact pathology of "eye-strain" is unknown, and the *rationale* of visual fatigue in the production of ocular and systematic disorders is largely a matter of conjecture. It is, in the first place, a safe principle to make the ametropic eye approximate by artificial means as nearly as possible to the emmetropic eye. This is effected by suitable spectacles. In the next place it is necessary to study the normal limits of adaptability of the eye to various conditions of illumination, &c., and to use the knowledge thus obtained to prevent these limits being transgressed. When we bear in mind the evolution of the visual apparatus in man, and the immense increase in the amount and nature of the work which it is called upon to perform in modern civilised life, it is surprising that eyes are capable of withstanding the strain.

Errors of Refraction.—The correction of ametropia by glasses has already been discussed. It is evident that theoretically this correction should be made as early as possible, and especially before the increased strain of school life is encountered. Much advance has been made in this direction in recent years, and the routine examination of the eyes of young school children ensures the discovery of serious errors. Facilities for their correction and for the supply of suitable glasses are now prevalent. The most difficult problem in this connection is that of myopia, which has already been mentioned in Chap. XXXIV.

Illumination.—Normal vision is capable of adaptation to very wide ranges of intensity and quality of illumination. It has been found that form vision is very defective under dark adaptation and with low intensity of illumination. As the intensity is increased and the eye becomes light-adapted visual acuity

increases – rapidly at first, and then only very slowly. Almost maximum visual acuity is reached with Snellen's test types when the illumination is about 2 foot-candles on the types. For very fine work, however, it is desirable that the illumination should be higher. The Departmental Committee of the Home Office on the Lighting of Factories and Workshops recommends that for "fine-work" the illumination on the working plane shall not fall below 3 foot-candles, and for "very fine work" not below 5 foot-candles. They have scheduled the various processes in a large number of occupations under these two headings. At extremely high illuminations, visual acuity is diminished owing to glare. It has been found that discrimination is increased by using monochromatic light, chiefly owing to the elimination of chromatic aberration in the eye: it is best with yellow light, the brightest part of the photopic spectrum.

There are many factors, however, which influence visual acuity besides the intensity of the light. Among these is the size of the pupils, but more important is the amount and character of the light falling upon peripheral areas of the retina. Thus, it is undesirable that there should be too great contrast between the areas under observation and surrounding areas. A brilliantly illuminated field of work in an otherwise dark room causes rapid alterations of adaptation which are deleterious. Hence a moderate amount of general illumination is preferable, and this has the additional advantage that it prevents the formation of very sharply defined shadows. On the other hand, it is very important that there should be no glaring lights in the field of vision; such lights should be carefully shaded. Care, too, should be taken to avoid direct reflection of light into the eyes. Thus, in reading, especially books written on shiny paper, and in working on bright metals, &c., if the source of light is in front of the eyes light is reflected directly into them. This light is useless for visual purposes, and indeed diminishes the contrasts which are the basis of discrimination. Hence the source of light should be placed laterally, and preferably to the left-hand side and somewhat behind the worker. Flickering lights should be avoided.

Various sources of light differ much in intensity and quality. The natural criterion is sunlight, which we are accustomed to regard as white light. Sunlight differs much, however, in "whiteness," and in intensity on different days, at different times of the day, and whether direct or diffuse. Owing to the adaptability of the eye it is difficult to judge the intensity of a given illumination. Measurements show that bright direct sunlight may give several thousand foot-candles, and an illumination of several hundred foot-candles on a well-placed desk is quite common. One great advantage of daylight is its diffusion; the illumination of a

room usually comes, not directly from the sun but from a considerable area of sky, and is reinforced by innumerable reflections from buildings and other objects. Sunlight is much richer in radiation of short wave length—blue and violet—than any artificial illuminants. Most modern illuminants have continuous spectra derived from incandescent solids; the higher the temperature the more nearly the energy distribution of the spectrum approximates to that of sunlight. Incandescent gases —such as used in the mercury-vapour lamp—have line spectra; they therefore more nearly approximate monochromatic light.

Glare may be regarded as light in the wrong place. The more concentrated the light the more disturbing is the effect. Glare, therefore, varies rather with the intrinsic brilliancy of the light than with its intensity. Intrinsic brilliancy is defined as candle-power per square inch. Clear sky has a very low intrinsic brilliancy—about the same as the candle, viz., 2·5 candles per square inch. A metal filament has an intrinsic brilliancy of 800 c. per square inch, and an arc light 20,000 c. per square inch. In general the ratio of the intrinsic brilliancy of a source of light to that of the surrounding field should not exceed 100. Glare is diminished in artificial interior illumination by the use of indirect lighting. In this method the light is reflected from the ceiling and suitably curved cornices, so that no direct light reaches the eye. By it shadows are almost eliminated. It is a restful, but monotonous mode of illumination; it is quite unsuited for certain purposes. Thus, sewing is very difficult with it, especially the sewing of monochromatic material, because the threads of the texture throw no shadows, and consequently their discrimination is made very difficult. In semi-indirect lighting the use of opalescent bowls permits of a certain amount of direct illumination.

Many modern illuminants emit a considerable amount of ultra-violet radiation, which is deleterious (*vide* pp. 179, 346). Most of this is absorbed by glass, so that the dangers arising from this cause are slight and have been much over-rated. It must be remembered, however, that globes absorb an appreciable amount of the luminous energy; even clear glass globes absorb 5—15 per cent., and opal globes as much as 10 —40 per cent. The distribution of light from artificial sources varies greatly. It can be modified by the use of reflectors and prismatic (holophane) globes. Too little attention has hitherto been paid by architects and others to the position and characters of light sources from the hygienic point of view. It is of great importance in the lighting of factories and workshops, and especially in that of schools. A committee of the Illuminating Engineering Society (1911), recommended the following standards for school rooms:—

"(I.) *Desk Illumination.*—(*a*) For ordinary clerical work

(reading and writing, &c.), the minimum illumination measured at any desk where the light is required should not fall below 2 foot-candles. (*b*) For special work (art classes, drawing offices, workshops, and stitching with dark materials, &c.), a minimum of 4 foot-candles is desirable. (*c*) For assembly rooms, &c., and for general illumination, a minimum of 1 foot-candle measured on a horizontal plane 3 feet 3 inches from the ground.

"(II.) *Blackboard Illumination.*—In rooms where students are distant more than 20 feet from the blackboard, and where it is customary to use diagrams in coloured chalk, an illumination on the blackboard of 60 per cent. in excess of that in the rest of the room is desirable. The board should be maintained a dead black and repainted at regular intervals."

Reading and Writing.—Considering the vast importance of reading and writing in modern life it is surprising that they have been so little investigated by physiologists and ophthalmologists. The forms of printed types are derived from manuscripts, and have been modified for technical reasons. Further advance has been almost entirely empirical, and even in the best presses more care has been exercised in obtaining æsthetic effects than in fostering legibility.

If we consider ordinary Roman printed characters we find that all capital letters extend above the line. Of the small letters thirteen are short, eight extend above the line (ascending letters), and four below the line (descending letters). There are thus twice as many ascending as descending letters, and in an ordinary page of print it will be found that of the long letters about 85 per cent. are ascending and only 15 per cent. descending. Examination of the short letters shows that their most characteristic features are in the upper parts. Hence in reading, attention is specially directed to the upper parts of the letters, as is strikingly demonstrated by covering the lower parts of a line of print with a card. The print is almost as legible as if it were uncovered. If, however, the upper halves of the letters are covered, it is almost, if not quite, impossible to read the print.

The ends of the lines of which letters are composed are commonly emphasised by means of serifs. These were doubtless introduced empirically, but the advantage in sharpness of definition has a physiological basis. They counteract irradiation, and hence the visibility of letters is improved if the serifs are triangular.

The tendency of typefounders has been to minimise the differences between letters, probably with a view to greater regularity of line and uniformity in appearance. For example, round letters have been flattened laterally and square letters rounded. The loops of b, d, p, and q, have been equalised to o. If the lower parts of short letters are covered, the similarity in the

topmost curves of a, c, e, o, s, of n and r, of h and b, or of n and p, is much greater in modern print than in some early samples.

Legibility is not determined solely by visibility in the physiological sense of the term. Thus, the emphasis of some lines in letters, which originated in the use of reeds and pens for writing, increases legibility whilst diminishing visibility. A child learning to read depends upon physiological visibility ; hence there should be little difference between the breadth of the thick and slender strokes. As facility in reading is acquired, legibility is increased by diminishing the breadth of the slender strokes, and as smaller letters are used the diminution must be more rapid than that of the heavy strokes, so that the interspaces may not be unduly contracted. At the same time, the slender strokes must not transgress the limits of visibility at reading distance, and their distribution should be emphasised by suitably formed serifs. Hence, Jaeger small types are more legible than Snellen's.

The spacing of the letters and words has a considerable effect upon legibility. Irradiation plays an important part here. Roughly speaking, the interspace between letters should be at least as broad as the blanks in m or n, but round letters like o and e should have slightly less interspace than square letters. Owing to irradiation the interspaces in general look larger than they really are, and two o's separated by a space look farther apart than two n's separated by the same space. Javal attributes a large part of the "remarkable legibility of English books" to the shortness of most English words and the consequent multiplication of blank interspaces. Of course, the spacing of words, and to a less degree of letters, in ordinary printing is very largely haphazard as far as legibility is concerned, the main object of the printer being to obtain general uniformity of appearance with rigid equality in the lengths of the lines. There is some difference of opinion as to whether "leading" or interlinear spacing is beneficial. Owing to the design of the blocks of type there is always a small space between the lower limits of descending and the upper limits of ascending letters, even without leading.

A line of print is read in a series of small jumps. At each pause a group of about ten letters is more or less accurately visualised ; the movements are too rapid to permit of visualisation whilst they are occurring. The number of leaps taken by the eye remains the same irrespective of the distance of the book, so long as this is consistent with legibility. A child reading makes more jumps in a line than the average, and the same applies to people reading a foreign language or correcting proofs. Attention is directed chiefly to the commencements of words, and words are not read by letters but by their general configuration.

There is, therefore, a very important psychological factor involved in the act of reading, quite apart from the interpretation of the meaning of the words.

Enough has been said to show that reading is a highly complex act, and the rules which can at present be devised for the avoidance of strain and discomfort involve a multiplicity of factors which have not yet been satisfactorily correlated.

Handicrafts.—The same visual principles as have been discussed above underlie the carrying on of many handicrafts and industrial processes, but each provides specific problems. These have hitherto been submitted to little scientific investigation.

No attempt has been made in this Section to deal exhaustively with so extensive a subject as Preventive Ophthalmology, but it has been deemed advisable to indicate to the student how innumerable and complex are the applications of ophthalmology to everyday life.

APPENDIX I.

PRELIMINARY INVESTIGATION OF THE PATIENT.

Scientific observation can only be fostered by methodical investigation. The student is recommended to study each patient according to the following scheme :—

(1) **General condition of the patient.**

(2) **Position of the head.**—Characteristic in paralyses of extraocular muscles (*vide* p. 506).

(3) **Face.**—Note asymmetry, facial paralysis, affections of the skin, *e.g.*, herpes ophthalmicus, &c.

(4) **Position of the eyebrows.**—Vicarious action of the frontalis in ptosis (*vide* p. 581), &c.

(5) **Orbits.**—Exophthalmos, enophthalmos, &c.

(6) **Eyeballs.**—(*a*) Position and direction—strabismus.

(*b*) Movements (*vide* p. 502).

(*c*) Size and shape—microphthalmia, glaucoma (*vide* p. 268) ; myopia (*vide* p. 472), buphthalmia (*vide* p. 283), staphyloma, &c.

(7) **Lids.**—(*a*) Position—ptosis, ectropion, entropion, &c.

(*b*) Palpebral aperture—ptosis, exophthalmos, &c.

(*c*) Movements—ptosis, exophthalmic goître, &c.

(*d*) Margins—blepharitis, tumours, &c.

(*e*) Lashes—trichiasis, distichiasis, &c.

(*f*) Glands—hordeolum, chalazion, &c.

(*g*) Puncta lacrymalia—eversion, occlusion, &c.

(*h*) Lacrymal sac—swelling, regurgitation, &c.

(8) **Conjunctiva.**—(*a*) Ocular — congestion (conjunctival and ciliary), secretion, phlyctenules, growths, wounds, &c.

(*b*) Palpebral—congestion, granulations, scars, concretions, ulcers, &c.

(*c*) Plica semilunaris—displacement in pterygium growths, &c.

(*d*) Caruncle—inflammation, granulations, growths, &c.

(9) **Cornea.**—(*a*) Size—glaucoma, buphthalmia, &c.

(*b*) Curvature—conical, buphthalmia, anterior staphyloma, abrasions (*vide* p. 81), &c.

(*c*) Surface—corneal reflex, abrasions, ulcers, foreign bodies, &c.

(*d*) Transparency—ulcers, nebulæ, keratitis (superficial and deep vascularisation, *vide* p. 86), "k.p.," striate opacity, &c.

(10) **Sclerotic.**—(*a*) Curvature and colour—myopia, staphyloma, episcleritis, &c.

(*b*) Vessels—ciliary injection, episcleritis, scleritis, &c.

(11) **Anterior chamber.**—(*a*) Depth—(α) shallow—perforating wound glaucoma, dislocation of lens, &c.; (β) deep—buphthalmia, iridocyclitis, dislocation of lens, &c.; (γ) irregular—iris bombé, dislocation of lens, &c.

(*b*) Contents—cloudy aqueous, hypopyon, hyphæma, foreign bodies, dislocated lens, &c.

(12) **Iris.** (*a*) Colour—muddy in iritis, heterochromia (congenital and in iridocyclitis), atrophy, ectropion of uvea, melanomata, &c.

(*b*) Position, especially plane of surface—iris bombé, retraction in iridocyclitis, pseudo-glioma, &c.

(13) **Pupils.**—(*a*) Relative size (*vide* p. 88).

(*b*) Reaction to light—direct, consensual, maintenance of constriction (*vide* p. 88).

(*c*) Reaction on convergence.

(*d*) Synechiæ—anterior and posterior.

(14) **Intraocular tension.**—Increased in glaucoma, iridocyclitis; diminished in iridocyclitis, perforation or rupture of the globe, &c.

(15) **Central vision** (*vide* p. 125).

(16) **Lens.**—By oblique illumination and the ophthalmoscope.

(17) **Vitreous.**—Opacities, fluidity, foreign bodies, hæmorrhage, "retinitis proliferans," persistent hyaloid artery, &c.

(18) **Fundus.**—(*a*) Optic disc—blurring of the edges, swelling, cupping, colour, crescents, &c.

(*b*) Retinal vessels—size, contour, tortuosity, &c.

(*c*) General view—retinitis, choroiditis, &c.

(*d*) Periphery.

(*e*) Macula.

(19) **Field of vision.**

APPENDIX II.

THERAPEUTIC NOTES.

(*The strengths of lotions, &c., are given in percentages: 1 per cent. = gr. v. to ʒ i. (approximately).*)

Lids.

Lotions.—3 per cent. Sodium bicarbonate.
3 per cent. Borax.

These are used for dissolving the crusts in blepharitis. They may be used in conjunction with 1 to 2 per cent. salicylic acid, 1 to 2 per cent. resorcin, liquor carbonis detergens (♏ i.—iv. to ʒ i.), &c.

Pigments.—10 per cent. Silver nitrate.
1 per cent. Picric acid.
1 per cent. Salicylic acid.

These are used for ulcerative blepharitis; the excess should be removed with cotton wool.

Ointments.—2 per cent. Ammoniated mercury.
3 per cent. Yellow oxide of mercury.

These are well rubbed into the lashes for five minutes three times a day after removing the crusts in ulcerative blepharitis. Alternative preparations are 1 per cent. salicylic acid, 3 per cent. airol, 2 per cent. resorcin, 3 per cent. ichthyol, &c.

Hypodermic Injections for Local Anæsthesia:
1 to 2 per cent. Novocain.
1 per cent. Stovain.
1 per cent. Alypin.

These drugs are less toxic than cocain. They may be combined with advantage with a hæmostatic, adrenalin or suprarenin (0·0025 per cent.). Novocain-adrenalin solution injected deep into the orbit may be used for excision of the eye (*vide* p. 459). Alypin-adrenalin solution similarly injected has been advocated for intraocular operations on inflamed eyes, and is said to cause no serious results (Pooley).

Conjunctiva.

Lotions.—3 per cent. Boric acid.
1 in 5000 Perchloride of mercury.

These are used as cleansing lotions in acute conjunctivitis, and have slight antiseptic properties; they should be mixed with an equal quantity of hot water as a rule.

2 per cent. Borax.

This, combined with dilute hydrocyanic acid (1 per cent.), relieves irritation in mild conjunctivitis.

1 per cent. Alum.

1 per cent. Zinc sulphate or chloride.

These are astringent lotions used in chronic conjunctivitis. Zinc lotion is specially indicated in angular conjunctivitis (*q.v.*). Alternative preparations are 1 per cent. tannin, $\frac{1}{4}$ per cent. copper sulphate, 3 to 6 per cent. sodium sozoiodol, $\frac{1}{2}$ per cent. zinc sulphocarbolate, &c.

1 per cent. Quinine hydrochloride or sulphate.

This is recommended for membranous conjunctivitis. The least possible amount of dilute sulphuric acid should be used to dissolve the salt if the sulphate is used.

Pigments.—2 per cent. Silver nitrate.

This is used in acute conjunctivitis. Alternative preparations are 10 to 40 per cent. argyrol, 10 to 40 per cent. protargol, 5 to 10 per cent. argentamin, &c.; these are probably not so effectual.

2 to 4 per cent. Perchloride of mercury in glycerin and water.

This is occasionally used in trachoma; it must not be allowed to touch the cornea. Iced compresses should be applied immediately after the application is made.

Drops.—The astringent lotions may be used in the form of drops. Silver nitrate should not be ordered in this form on account of the staining which may occur from prolonged use.

2 to 4 per cent. Cocain hydrochloride.

This is used for producing local anæsthesia. Conjunctival and corneal anæsthesia is complete after instilling four drops at intervals of five minutes. During the intervals the eyes should be kept closed on account of the desiccating action on the corneal epithelium. In intraocular operations one drop should be instilled into the opposite eye (*vide* p. 447).

2 per cent. β-Eucain hydrochloride.

2 per cent. Alypin.

These may be used instead of cocain. They do not dilate the pupil and are less toxic than cocain, but they cause considerable smarting.

1 to 10 per cent. Dionin.

This causes intense œdema of the conjunctiva; after prolonged use the reaction is slight or absent. The patient complains of a burning sensation which quickly disappears. The first application should be made by the surgeon. The drug stimulates the lymph flow, and has therefore been recommended in a large variety of cases, *e.g.*, corneal nebulæ, scleritis, iridocyclitis, &c. It must be used with caution.

Subconjunctival Injections.—These have been recommended for the same reason as dionin, and their value, if any, is still unproved. Many solutions have been used. Sterile 2 to 10 per cent. salt solution is the best; the others probably possess no advantage, and are in some cases liable to cause necrosis. The injection is made with a hypodermic needle under the bulbar conjunctiva as far back as possible above the globe. There is considerable reaction and pain. Not more than 5 to 10 minims should be injected every other day.

Ointments.—Vaseline.

15 per cent. Boric acid ointment.

These are used to prevent the lids from sticking together and thus causing retention of secretion.

1 to 3 per cent. Yellow oxide of mercury.

This is used as a stimulant and antiseptic in phlyctenular and chronic conjunctivitis.

Cornea.

Lotions.—The same collyria as for conjunctival conditions are used as cleansing and antiseptic measures. The astringent collyria are seldom indicated.

Pigments.—1 per cent. Silver nitrate.

This is indicated in some cases of marginal ulceration (*vide* p. 212).

Pure carbolic acid.

This is used as a cauterising agent in hypopyon ulcer (*vide* p. 206).

Drops.—These are the ordinary mydriatics and cycloplegics, and in rare cases miotics (*vide infra*).

Dionin or subconjunctival injection is sometimes used with a view to clearing corneal nebulæ. Two per cent. fluorescein in 3 per cent. bicarbonate of sodium solution is used for staining ulcers and abrasions for diagnostic purposes.

Ointments.—These are the same as for conjunctival conditions.

2 to 5 per cent. iodoform, xeroform, or airol ointment may be used in corneal ulceration. Dionin (5 to 10 per cent.) and mydriatics may be prescribed in ointment form. 1 to 5 per cent. yellow oxide of mercury ointment is used in gradually increasing strength to aid in the clearing of nebulæ and the opacity of interstitial keratitis; it may be combined with dionin.

Mydriatics and Cycloplegics; Miotics.

Mydriatics and Cycloplegics (*vide* p. 57).—$\frac{1}{2}$ to 1 per cent. Atropin sulphate. $\frac{1}{4}$ to $\frac{1}{2}$ per cent. Hyoscin or Scopolamin hydrobromide. $\frac{1}{4}$ to $\frac{1}{2}$ per cent. Duboisin sulphate. $\frac{1}{2}$ to 1 per cent. Daturin sulphate.

These are used as drops or ointment in corneal ulcers, iritis, iridocyclitis, &c. Atropin is used generally, the others being employed as substitutes when atropin causes irritation. Hyoscin occasionally causes delirium and should be watched.

1 per cent. Homatropin hydrobromide, with or without 2 per cent. cocain hydrochloride.

1 to 2 per cent. Euphthalmin hydrochloride.

These are used for investigation of refraction and for ophthalmoscopic examination; also for the diagnosis of synechiæ (*vide* p. 246) and occasionally for the diagnosis of increased tension (*vide* p. 246). For refraction solutions of the same strength of the bases in castor oil are more certain in their action.

2 per cent. Cocain hydrochloride.

This is used occasionally in old people as a mydriatic for ophthalmoscopic purposes, being less likely to raise the tension, the effect being readily counteracted by miotics.

Miotics (*vide* p. 58).—$\frac{1}{2}$ to 1 per cent. Pilocarpin hydrochloride. $\frac{1}{4}$ to 1 per cent. Eserin or Physostigmin sulphate or salicylate, with or without 1 per cent. cocain hydrochloride.

These are used in glaucoma (*q.v.*), and occasionally in other cases (*vide* p. 212). The effect of cocain is to assist the absorption of the other drug.

Vaso-Constrictors.

1 in 1000 adrenalin hydrochloride or suprarenin may be added with advantage to the cocain solution in performing tenotomy, advancement, &c. It is used alone in excision of the lacrymal sac, and occasionally in conjunctivitis (*vide* pp. 165, 178). It may be combined with $\frac{1}{2}$ per cent. chloretone.

The Treatment of Syphilitic Affections.

In cases of suspected syphilitic disease of the eye the possibility of demonstrating the presence of the spirochæte should be borne in mind. More generally useful is the application of Wassermann's test, for which the aid of a bacteriologist is advisable. No anti-syphilitic remedies should be used until the test has been applied. When the lesion is probably syphilitic the patient should be brought rapidly under the influence of mercury. For this purpose inunctions, intramuscular injections, or intravenous injections may be used, supplemented if necessary by administration by the mouth. Inunction is generally employed, and for this purpose the oleate of mercury is preferable to the ordinary mercury ointment. Intramuscular and intravenous injections require special technique. Intravenous injection of salvarsan ("606") or its equivalent is sometimes remarkably successful, especially in the acute stage of syphilitic manifestations. When the acute ocular condition is subsiding a course of treatment at Aix-la-Chapelle may be recommended when feasible. Some recent substitutes for the ordinary iodides are sajodin or iodoglidine tablets (1 to 3, *i.e.*, 7 to 15 gr. three times a day), iodipin (30 gr. of 25 per cent. = about 10 gr. of potassium iodide), &c.

Serum and Vaccine Treatment.

In diphtheritic conjunctivitis antitoxic serum must be used. The indications for other sera and vaccines in the domain of ophthalmology are much less precise. Thus Römer has introduced an antipneumococcic serum for the treatment of hypopyon ulcer, but the results obtained have been disappointing. Antigonococcic serum has been advocated in ophthalmia neonatorum and gonorrhœal ophthalmia. In desperate cases of septic infection of the eye after perforating wound, accidental or operative, a polyvalent serum may be employed, or if possible a vaccine made from a culture taken from the eye: oral administration of antidiphtheritic serum has been recommended in these cases. The vaccine treatment of tuberculosis is much used. Many obscure pathological conditions in the eye suggest the possibility of a tuberculous origin. Diagnosis may be facilitated in some, probably a minority of cases, by the application of von Pirquet's cutaneous test. Wolff-Eisner and Calmette's conjunctival reaction should

not be used. Occasionally, as in conjunctival tuberculosis, it is possible to place the diagnosis beyond doubt by inoculation experiments on animals; the incubation period, however, is lengthy (20 to 30 days). Tuberculin treatment is best carried out by a bacteriologist who has familiarised himself with the technique.

Care of Instruments.

Ophthalmic instruments should be kept in an air-tight glass cabinet, or when not constantly in use in velvet-lined cases.

All instruments should be sterilised before use by boiling in 3 per cent. sodium carbonate solution (*not* bicarbonate), made with *distilled* water. This procedure does not impair the cutting edges, but knives and scissors should not be boiled more than three to five minutes: this is amply sufficient if the surfaces are bright and free from tarnish, as they ought to be. If distilled water cannot be obtained the cutting instruments should be well soaked in pure carbolic acid before being transferred to the dish.

The instruments may be removed from the steriliser *immediately* before operating and used dry. More commonly they are immersed in fluid in a dish. The fluid should be 3 per cent. sodium carbonate which has been boiled. *In no case must boric lotion be used*, since it tarnishes the steel.

It is, however, much safer to use the instruments dry. It is almost impossible to sterilise the skin of the hands efficiently, and if the instruments are wet fluid from the fingers is liable to run along them into the eye. The points of knives, &c., should be dipped in sterile saline immediately before use to facilitate their passage through the tissues.

The surgeon should wear a sterilised gown, or at least sterilised sleeves, and also a mask covering the nose and mouth, for all intraocular operations.

APPENDIX III.

REQUIREMENTS OF CANDIDATES FOR ADMISSION INTO THE PUBLIC SERVICES.

Commissions in the Army.

The minimum standards of acuteness of vision with which a candidate for a commission will be considered fit are as follows :—

Standard I.

Right Eye.	*Left Eye.*
Distant vision.—V = 6/6.	V = 6/6.
Near vision.—Reads 0, 6.	Reads 0, 6.

Standard II.

Better Eye.	*Worse Eye.*
Distant vision.—V = 6/6.	V, without glasses = not below 6/60; and after correction with glasses = not below 6/24.
Near vision.—Reads 0, 6.	Reads 1.

Standard III.

Better Eye.	*Worse Eye.*
Distant vision. — V, without glasses = not below 6/24, and after correction with glasses = not below 6/6.	V, without glasses = not below 6/24; and after correction with glasses = not below 6/12.
Near vision.—Reads 0, 8.	Reads 1.

Each eye must have a full field of vision as tested by hand movements.

Squint or any morbid condition of the eyes or of the lid of either eye liable to aggravation or recurrence will cause the rejection of the candidate.

Each eye will be examined separately, and the lids must be kept wide open during the test. The candidate will be required to read the tests in ordinary daylight.

Inability to distinguish the principal colours will not be regarded as a cause for rejection, but the fact will be noted in the proceedings, and the candidate will be informed.

The degree of acuteness of vision of all candidates for commissions will be entered in the proceedings as follows :—

V.R. = ... ; with glasses = ... Reads .....
V.L. = ... ; with glasses = ... Reads .....

No relaxation of the standard of vision will be allowed.

Commissions in the Royal Navy.

Blindness or defective vision, squint, imperfect perception of colours, fistula lachrymalis, or any chronic disease of the eyes or eyelids renders the candidate unfit. Candidates for first appointments in the Royal Marines (R.M.A. and R.M.L.I.) must possess full normal vision as determined by Snellen's tests, each eye being separately examined. Candidates for Naval Cadetships at the Royal Naval College, Dartmouth, must possess full normal vision, viz., 6/6, and be able to read D = 0·6 as determined by Snellen's test types, each eye being separately examined. A high degree of hypermetropia will disqualify. Candidates for Naval Cadetships (Special Entry): The standard of distant vision on entry is—one eye 6/6, and the other not less than 6/12 without glasses. The defect must be due to refractive errors. For near vision they must be able to read D = 0·6 without glasses with each eye. A high degree of hypermetropia will disqualify. This lower standard will only be accepted on the clear understanding that these officers will undergo a further test of eyesight on the completion of their Sub-Lieutenant's examination at the schools, and a further examination if considered necessary on attaining the age of twenty-five.

Recruits for the Royal Navy and Royal Marines.

1. For candidates for the Seaman Class (including Boys and Youths), Royal Marines (excluding Marine Bandsmen), Engine Room Artificers, Ordnance Artificers, Boy Artificers and Ship's Cooks, full normal vision is required, viz., 6/6 each eye tested separately.

2. For candidates for Electrical Artificers, other Artisan Ratings (Shipwrights, Joiners, Blacksmiths, Plumbers, Painters, &c.), and Stokers, the vision must be 6/9 each eye tested separately.

3. For Sick Berth Staff and Royal Marine Band Boys, 6/12 each eye tested separately.

4. For all other ratings, including Writers, Victualling Assistants, Boy Writers, Victualling Boys, Officers' Stewards and Cooks, and Boy Servants, the vision must not be less than 6/12 both eyes.

5. For all ratings, except Writers, Officers' Stewards and Cooks, or Royal Marine Bands, the colour sense must be normal.

6. Marine Bandsmen, Sick Berth Staff, Writers, Victualling ratings, and Officers' Stewards and Cooks are allowed to wear glasses, but defects of vision must only be due to errors of refraction, and must be capable of correction to 6/6 Snellen by means of glasses, and the candidate must be able to read D = 0. 6 without the aid of glasses.

7. To determine the acuity of vision, Snellen's letter types are used, and the colour sense is determined by means of the Edridge Green Colour Perception Lantern; if not available, by means of the bead test.

Appointments under the Government of India.

The Ecclesiastical, Education, Geological Survey, Agricultural, Indian Finance, Customs, Civil Veterinary, and other Departments not specially provided for in the following pages.

1. A candidate may be admitted into the Civil Services of the Government of India if ametropic in one or both eyes, provided that, with correcting lenses, the acuteness of vision be not less than 6/9 in one eye and 6/6 in the other; there being no morbid changes in the fundus of either eye.

2. Cases of myopia, however, with a posterior staphyloma, may be

admitted into the service, provided the ametropia in either eye does not exceed 2·5 D, and no active morbid changes of choroid or retina be present.

3. A candidate who has a defect of vision arising from nebula of the cornea is disqualified if the sight of either eye be less than 6/12 ; and in such a case the acuteness of vision in the better eye must equal 6/6, with or without glasses.

4. Squint or any morbid condition, subject to the risk of aggravation or recurrence, in either eye, may cause the rejection of a candidate. The existence of imperfection of colour sense will be noted on the candidate's papers.

*The Departments of Forest, Survey, Telegraph, Factories, and for various Artificers.**

1. If myopia in one or both eyes exists, a candidate may be passed, provided the ametropia does not exceed 2·5 D, and if with correcting glasses, not exceeding 2·5 D, the acuteness of vision in one eye equals 6/9 and in the other 6/6, there being normal range of accommodation with the glasses.

2. Myopic astigmatism does not disqualify a candidate for service, provided the lens or the combined spherical and cylindrical lenses required to correct the error of refraction do not exceed — 2·5 D ; the acuteness of vision in one eye, when corrected, being equal to 6/6, and in the other eye 6/9, together with normal range of accommodation with the correcting glasses, there being no evidence of progressive disease in the choroid or retina.

3. A candidate having total hypermetropia not exceeding 4 D is not disqualified provided the sight in one eye (when under the influence of atropin) equals 6/9 and in the other eye equals 6/6, with + 4 D or any lower power.

4. Hypermetropic astigmatism does not disqualify a candidate for the service, provided the lens or combined lenses required to cover the error of refraction do not exceed 4 D, and that the sight of one eye equals 6/9 and of the other 6/6, with or without such lens or lenses.

5. A candidate having a defect of vision arising from nebula of the cornea is disqualified if the sight of one eye be less than 6/12. In such a case the better eye must be emmetropic. Defects of vision arising from pathological or other changes in the deeper structures of either eye, which are not referred to in the above rules, may exclude a candidate for admission into the service.

6. Squint or any morbid condition, subject to the risk of aggravation or recurrence, in either eye, may cause the rejection of a candidate. The existence of imperfection of colour sense will be noted on the candidate's paper.

Public Works Department and Superior Establishments, Railway Department.

1. If myopia in one or both eyes exists, a candidate may be passed, provided the ametropia does not exceed 3·5 D, and if, with correcting glasses not exceeding 3·5 D, the acuteness of vision in one eye equals 6/9 and in the other 6/6, there being normal range of accommodation with the glasses.

2. Myopic astigmatism does not disqualify a candidate, provided the lens, or the combined spherical and cylindrical lenses, required to correct the error of refraction, does not exceed 3·5 D ; the acuteness of vision in one eye, when corrected, being equal to 6/9 and in the other 6/6, together with

* Artificers engaged in map and plan drawing may be considered separately, and this standard relaxed if it appears to be desirable.

normal range of accommodation with the correcting glasses, there being no evidence of progressive disease in the choroid or retina.

3. A candidate having total hypermetropia not exceeding 4 D is not disqualified, provided the sight in one eye (when under the influence of atropin) equals 6/9, and in the other eye equals 6/6, with + 4 D glasses, or any lower power.

4. Hypermetropic astigmatism does not disqualify, provided the lens or combined lenses required to cover the error of refraction do not exceed 4 D, and that the sight of one eye equals 6/9 and the other 6/6, with or without such lens or lenses.

5. A candidate having a defect of vision arising from nebula of the cornea is disqualified if the sight of that eye be less than 6/12. In such a case the better eye must be emmetropic. Defects of vision arising from pathological or other changes in the deeper structures of either eye, which are not referred to in these rules, may exclude a candidate.

6. Squint or any morbid condition, subject to the risk of aggravation or recurrence, in either eye, may cause the rejection of a candidate. Any imperfection of the colour sense is a disqualification for appointment to the Engineering Branch of the Railway Department, or as Assistant Superintendent in the Traffic Department. In all other cases a note as to any imperfection of colour sense will be made on the candidate's papers.

The Indian Medical Service and the Police Department.

1. Squint, or any morbid condition of the eyes or of the lids of either eye liable to the risk of aggravation or recurrence, will cause the rejection of the candidate.

2. The examination for determining the acuteness of vision includes two tests; one for distant, the other for near vision. The Army Test Types will be used for the test for distant vision, without glasses, except where otherwise stated below, at a distance of 20 feet; and Snellen's Optotypi for the test for near vision, without glasses, at any distance selected by the candidate. Each eye will be examined separately, and the lids must be kept wide open during the test. The candidate must be able to read the tests without hesitation in ordinary daylight.

3. A candidate possessing acuteness of vision, according to one of the standards herein laid down, will not be rejected on account of an error of refraction, provided that the error of refraction, in the following cases, does not exceed the limits mentioned, viz.:—(*a*) in the case of *myopia*, that the error of refraction does not exceed 2·5 D; (*b*) that any correction for *astigmatism* does not exceed 2·5 D; and, in the case of myopic astigmatism, that the total error of refraction does not exceed 2·5 D.

4. Subject to the foregoing conditions, the standards of the minimum acuteness of vision with which a candidate will be accepted are as follows:—

Standard I.

Right Eye.	*Left Eye.*
Distant vision.—V = 6/6.	V = 6/6.
Near vision.—Reads 0, 6.	Reads 0, 6.

Standard II.

Better Eye.	*Worse Eye.*
Distant vision.—V = 6/6.	V, without glasses, = not below 6/60; and after correction with glasses, = not below 6/24.
Near vision.—Reads 0, 6.	Reads 1.

Standard III.

Better Eye.	*Worse Eye.*
Distant vision.—V, without glasses, = not below 6/24; and after correction with glasses, = not below 6/6.	V, without glasses, = not below 6/24; and after correction with glasses, = not below 6/12.
Near vision.—Reads 0, 8.	Reads 1.

N.B. In all other respects candidates for these two branches of the service must come up to the standard of physical requirements laid down for candidates for commissions in the army.

The Indian Pilot Service, and Candidates for appointments as Guards, Engine-drivers, Signalmen, and Pointsmen on Railways.

1. A candidate is disqualified unless both eyes are emmetropic, his acuteness of vision and range of accommodation being perfect.
2. A candidate is disqualified by any imperfection of his colour sense.
3. Strabismus, or any defective action of the exterior muscles of the eyeball, disqualifies a candidate for these branches of service.

The Indian Marine Service, including Engineers and Firemen.

1. A candidate is disqualified if he have an error of refraction in one or both eyes which is not neutralised by a concave or by a convex 1 D lens, or some lower power.
2. A candidate is disqualified by any imperfection of his colour sense.
3. Strabismus, or any defective action of the exterior muscles of the eyeball, disqualifies a candidate for this branch of service.

Special Duty.

Candidates for special duty under Government must possess such an amount of acuteness of vision as will, without hindrance, enable them to perform the work of their office for the period their appointment may last. In all cases of imperfection of colour sense a note will be made on the candidate's papers.

Pilot Service.

1. A candidate shall be examined both as to his physical fitness and as to his acuteness of vision and colour perception.
2. The sight tests shall be carried out in accordance with the Board of Trade Regulations applying to examinations for certificates of competency in force for the time being.
3. A candidate must have no defect of sight, and must have full normal vision in both eyes, each eye being examined separately without the aid of spectacles.
4. A candidate shall be disqualified by any imperfection of his colour sense or by squint, or any defective action of the eye muscles or any disease of the eye.
5. A candidate must be also otherwise physically fit for the duties of a pilot.

Masters and Mates of the Mercantile Marine.

I.—Form Vision Test.

1. *Form Vision Test to be passed first.*—The first test which the candidate is required to undergo is the test for form vision, and until he has passed

this test he must not be allowed to proceed further with the examination.

2. *Apparatus Used.*—The form vision test to be used for all candidates is that conducted on Snellen's principle by means of sheets of letters.

3. *Object of the Test.* The object of the form vision test is to determine whether the candidate can reach a sufficient standard of visual acuteness, or, in other words, to find out whether his eyesight is good or bad.

4. *Standard of Vision Required.*—Every candidate for a first certificate of competency will be required to possess normal vision. With the exceptions indicated below (*see* paragraph 8), every candidate for a second or higher certificate will be required to possess normal vision.

"Normal vision" is defined, for the purpose of these Regulations, as ability to read correctly nine of the twelve letters in the sixth line and eight of the fifteen letters in the seventh line of a test sheet placed in a good light at a distance of 16 feet from the eye.

The candidate will have the option of using either eye separately or both eyes together.

5. *Spectacles not Allowed.*—During the examination for form vision candidates must not be allowed to use spectacles or glasses of any kind, or any other artificial aid to vision.

6. *Method of Testing.*—The test sheets should be hung on the wall, *in a good light, but not in direct sunlight*, at a height of five or six feet from the ground. The candidate should be placed at a distance of exactly 16 feet from the sheets, and exactly opposite them. This distance should be carefully measured, and should never in any circumstances be varied.

One of the sheets should then be exposed, and the candidate should be asked to read the letters on each sheet, beginning at the top and going downwards. Any mistakes which he makes should be carefully noted. If then it is found that he has read correctly at least nine letters in the sixth line and eight letters in the seventh line of a sheet, the candidate may be considered to have normal vision, and should be marked "passed" in the appropriate column of the form of application (Exn. 2 or Exn. 2A, as the case may be).

7. *Passing or Failure.*—If at the conclusion of the test the candidate is found to reach the required standard, he may be considered to have passed, and the Examiner should proceed to test him for colour vision. If the candidate fails to reach the standard required for the certificate entered for, he should be tested with at least four sheets, and the Examiner should fill in a Form Exn. 17b, and should forward it with any remarks he may wish to make, to the Principal Examiner for his instructions as to whether the candidate is to be regarded as passing or as failing in form vision.

8. *Lower Standard Required in Certain Cases.*—Candidates who are in possession of certificates obtained before January 1st, 1914, may be regarded as passing in form vision if they can read correctly with both eyes at least five of the eight letters in the fifth line of a test sheet.

9. *Tests to be Varied.*—The Examiner should take care, by varying the order of the test sheets and by every other means in his power, to guard against the possibility of any deception on the part of the candidate.

10. *Result of Examination to be Reported.*—The result of every examination in form vision should be reported, in the case of a candidate for a certificate of competency, to the Registrar-General of Shipping and Seamen on Form Exn. 2, and to the Principal Examiner of Masters and Mates on Form Exn. 14; and, in the case of a candidate for the sight tests only, to the Registrar-General of Shipping and Seamen on Form Exn. 2A.

II.—Colour Vision Test.

11. *Apparatus.*—A special lantern and a mirror have been provided for this test. The lantern should be placed directly in front of the mirror, so

that the front part of the lantern is exactly 10 feet from the mirror. Care should be taken that the lantern is properly placed, that is to say, the lights reflected in the mirror must show clearly when viewed through the rectangular aperture on the left of the lantern. The Examiner should always satisfy himself that these conditions are fulfilled before commencing the examination.

12. Darkness Adaptation.—*It is essential that a candidate should be kept in a room which is either completely or partially darkened for at least a quarter of an hour before he is required to undergo this test.*

Before the examination commences the Examiner must satisfy himself that the room in which it is conducted is so darkened as to exclude all daylight.

13. *Method of Testing.*—The lantern supplied for the examination is so constructed as to allow one large or two small lights to be visible, and is fitted with 12 glasses of three colours—red, white and green. At the commencement of the examination the Examiner should show to the candidate a series of lights through the large aperture, and should require him to name the colours as they appear to him. Care should be taken in showing the white light to emphasise the fact that the light is not a pure white. If a candidate makes a mistake of calling the light " red," a proper red light should be shown immediately after and the candidate's attention directed to the difference between the two.

After a series of lights through the large aperture has been shown, the Examiner should make a complete circuit with the two small apertures, requiring the candidate to name the colours of each set of two lights from left to right. To prevent any possibility of the order in which the lights are arranged from being learnt, the Examiner should at least twice in each circuit go back a varying number of colours.

A record of any mistakes made with either the large aperture or the two smaller apertures should be kept on Form Exn. 17b in accordance with the instructions thereon.

14. *Passing or Failure.*—If a candidate with either the large aperture or the two smaller apertures of the lantern mistakes red for green or green for red, he should be considered to have " failed " in Colour Vision.

If the only mistake made by the candidate with the lantern is to call the white light " red," and if after his attention has been specially directed to the difference between the two he makes no further mistake of this nature, he should be considered to have passed in Colour Vision.

If a candidate makes any other mistake with the lantern, *i.e.*, if he calls white " red " repeatedly or red " white " at all, or confuses green and white, his case should be reported to the Principal Examiner of Masters and Mates and he should be told that the decision as to whether he is passed or failed, or a further examination is necessary, will be communicated to him in due course. Pending the receipt of the Principal Examiner's instructions, such a candidate should only be allowed to proceed with the remainder of the examination for a Certificate of Competency on the express understanding that the latter examination will be cancelled in the event of failure in the Sight Tests.

15. *Further Examination and Appeals.*—If in the cases covered by the preceding paragraph the Principal Examiner decides that a further examination is necessary, arrangements will be made for a special examination to be held in London, and the third class travelling expenses necessarily incurred by a candidate in attending such an examination will be paid by the Board of Trade, together with a subsistence allowance at a rate which will be notified to the candidate, but which will not in any circumstances exceed 12*s.* 6*d.* for each night's absence from home necessarily entailed by attendance at the Special Examination. If a night's absence is not involved expenses incurred during the day will be paid up to a maximum of 5*s.* for each day

necessarily occupied in attending the examination. In these cases the above expenses will be paid whatever may be the result of the final examination.

If, however, on the report of the local Examiner the Principal Examiner decides that the nature of the mistakes made shows conclusively that a candidate is so colour-blind as to be unfit to hold a certificate, the candidate shall be considered to have failed.

If during the progress of a special examination a candidate with defective Form Vision is found to be suffering from a permanent defect of the eyes such as to render him unfit for a sea career, the Special Examiners may finally reject such candidate, and he will not be allowed to be again examined in the Sight Tests on any future occasion.

In cases where, upon the report of the local Examiner, a candidate is failed by the Principal Examiner, as well as in the cases covered by paragraph 14, the Board will be prepared to allow a candidate who is dissatisfied with this decision to appeal for a special examination in London, but the Board will not pay the travelling expenses of any such candidate unless he is reported by the Special Examiners conducting the appeal examination to have passed.

INDEX.

(*The principal page references are printed in heavier type.*)

Aberration,
chromatic, 53
spherical, 52
Ablatio retinæ, 350
Ablepharon, 588
Abscess,
intracranial, ocular manifestations of, 553
lacrymal, 601
Accessory sinuses of nose, distension of, 608
Accommodation, 45, 499
age and, 47, 49
amplitude of, 47
and convergence, 499, 522
anomalies of, 489
effect of, 45
paralysis of, 489
range of, 49, 132
relaxing of, 116
spasm of, 128, 490
Achromatopsia, 380
Acne rosacea, 146, 221
Acuity of vision, 62, 125
Adaptation, retinal, 63
in glaucoma, 278
Adnexa, diseases of, 561
Adrenalin, 142, 165, 536, 598, 632
Advancement operation, 512, 528, **535**
Age and accommodation, 47, 49
and glaucoma, 267
errors of refraction and, 486
incidence of ametropia, 475
influence on hypopyon ulcers, 204
influence on refraction, 50
Albinism, 94, 120, **355**, 514
Albuminuric neuro-retinitis, 340
Alcohol, amblyopia due to, 349
Alypin, 630
Amaurosis, 376
fugax, 377
uræmic, 377
Amaurotic cat's eye, 93
Amaurotic family idiocy, 333
Amblyopia, **376**
congenital, 376, 521
ex anopsia, 376, 521
heteronymous, 376
hysterical, 377
quinine, 348
tobacco, 347
toxic, 347, 349
Ametropia, **35, 472,** 621
age incidence of, 475
axial, 41, 110, 472, 480
curvature, 41, 110, 472, 480, 483
glasses for, 38
index, 41, 111, 472, 480, 483
Amotio retinæ, 350
Amplitude
of accommodation, 47
of convergence, 500
Anæmia
of retina, 329
secondary, ocular complications of, 558
Anæsthesia, local, hypodermic injections for, 629
Anaphylaxis, 225
Anatomy
of cornea, 1
of extrinsic ocular muscles, 492
of eye, 1
of lacrymal apparatus, 589
of the lids, 561
Angioid streaks, 335
Angioma
of lids, 586
of orbit, 610
Angle
of anterior chamber, 3, 18, 258, 264, 432
metre, 499
minimum visual, 62
visual, 34, 125
Angular conjunctivitis, 166
Aniridia,
congenital, 261
traumatic, 400
Anisometropia, 51, **485**

Ankyloblepharon, 580, 587
Annular scleritis, 239
Anterior chamber,
anatomy of, 3
delayed reformation of, 457
examination of, 86
investigation of, 628
Aphakia, 111, 480, **483**
Aqueous humour, 3, 17
refractive index of, 32
secretion of, 17
Arcus
juvenilis, 228
senilis, 228
Argyll Robertson pupils, 89, 541
Argyrol, 148
Argyrosis, 165, 184, 235
Army, requirements of candidates for commissions, 634
Arteries,
ciliary, 11
of eye, 10
ophthalmoscopic appearance of, 120
retinal, degenerative changes, 334
Artificial eye, 462
Aspergillus fumigatus, 210
Asthenopia, 481, 482, 531
Astigmatism, 41, 43, 114, **482**
causes of, 51
hypermetropic,
compound, 43
simple, 43
irregular, 45
lenticular, 51
mixed, 43
myopic,
compound, 43
simple, 43
regular, 44
Astringent lotions, 630
Atheromatous ulcers, 214
Atrophy, optic, 365
Atropin, **57**, 253
dangers of, 165
in corneal ulceration, 195
irritation, 176
ointment, 195
routine use of, 209
Auditory nerve, tumours of, 548

Bacillus coli, 154, 158
Bacillus,
diphtheria, 156
Koch-Weeks', 144, 158
Morax-Axenfeld, 166
tubercle, 161, 175, 252, 316, 595
xerosis, 158, 183
Bacterial infection of the eye, 318
Bacteriology
of conjunctiva, 139
of cornea, 187, 203
of mucocele, 594
of sympathetic ophthalmia, 428
Bandage, Moorfield's, 450
Band-shaped opacity, 228
Basedow's disease, 614
Beer's
discission needle, 438
knife, 569
Binocular
diplopia, **505**
vision, 498
field of, 137
Birth injuries, 234
Bjerrum's
screen, 137
test types, 138
Blennorrhœa, acute, 148
Blepharitis, 564
Blepharophimosis, 581
Blepharospasm, 83, 573
"Blight," 143
Blindness, causes and prevention of, 617
Blind spot, 60
Blood,
diseases of, 558
in cornea, 235, **398**
supply of eye, 10
vessels
of cornea, 85, 86
of retina, 8
degenerative changes, 334
disorders of, 329
of uveal tract, 12
ophthalmoscopic appearance, 120
Blue sclerotic, 80, **240**
Bowman's
discission needle, 438
membrane, 1
Bruch, membrane of, 7
Buller's shield, 150
Buphthalmia, 282
Burns, injuries caused by, 396
Burow's operation, 574

Calcareous film, transverse, 228
Canal
of Cloquet, 17
of Schlemm, 4, 13
Canaliculus, 589
congenital anomalies of, 593
occlusion of, 593
slitting of, 592

Candidates for public services, requirements of, 634
Canthoplasty, 573
Capillary pulsation of retinal vessels, 121
Capsulo-pupillary membrane, 261
Carbolic acid, application of, 207
Carcinoma
of choroid, 389
of lacrymal gland, 591, 611
of lids, 586
" Cartella " shield, 451
Cataract, **285**
after-, 301
anterior
capsular, 194, **297**
cortical, 297
polar, 297
axial, 299
black, 289
central, 299
complicated, 302
concussion, 403
congenital, 295, 299
coralliform, 299
diabetic, 303
discoid, 299
extraction,
combined, 447
complications after, 456
without iridectomy, 455
fusiform, 299
glass-workers', 304, 619
hard, 289
hypermaturity of, 288
immature, 293
incipient, 291
infantile, 295
ironworkers', 305, 619
lamellar, treatment of, 295, 296
lightning, 403
mature, 288, 292
Moorfield's bandage, 450
Morgagnian, 288
nuclear, 299
perinuclear, 295
posterior
cortical, 308, **298**
polar, 298, **302**, 310
punctate, 299
pyramidal, 299
rate of development of, 290
secondary, 301
discission of, 443
senile, **285**
extraction of, 445
intranuclear, 286
subcapsular, 286
supranuclear, 286
soft, 289
Cataract—*contd.*
symptoms of, 289
total, 288, 300, 302
traumatic, 403, **408**
treatment of, 291, 293
unilateral, 293, 301
zonular, 295
Catarrhal conjunctivitis, 143
Caustics, injuries caused by, 396
Cauterisation of cornea, 206
Cautery, galvano-, 206
Cavernous sinus,
thrombosis of, 605
tributaries of, 606, 607
Cellulitis, orbital, 604
Cerebral
peduncle, diseases of, 549
syphilis, 551
Chalazion, 568
Chemosis, 141, 182
Chlorosis, 558
" Choked disc," 358
Cholesterin, 288, 291, 308
Choriocapillaris, 6
Choroid,
anatomy of, 6
carcinoma of, 389
coloboma of, 354
detachment of, 327
diseases of, **312**
injuries to, 403
lymph production of, 18
sarcoma of, 383
tubercle of, 316
Choroiditis, **312**
anterior, 315
central, 316, 321
areolar, 322
guttate, 321
circumpapillary, 323
diffuse, 316
disseminated, 313
Förster's areolar, 322
juxtapapillaris, 316
metastatic, 320
myopic, 474
purulent, 423
syphilitic, atrophy after, 314
Tay's, 321
Chromatic aberration, 53
Chromatopsia, 380
Cicatrices, corneal, 191
Ciliary
arteries, 11
body, 17, 46
anatomy of, 5
diseases of, 241
gumma of, 260
inflammation of, 256

Ciliary—*contd.*
body—*contd.*
injuries to, 400, 425
sarcoma of, 388
tubercle of, 260
congestion, 78
muscle, 5
processes, 6
staphylomata, 273
veins, 11
Circle of least diffusion, 42
Circulatory system, ocular complications of, 558
Circulus arteriosus iridis major and minor, 13
Cloquet, canal of, 17
Cocain, **58**, 142, 630
"Colloid bodies," 322
Coloboma
of choroid and retina, 354
of iris, 261
of lens, 305
of lid, 588
of optic disc, 369
Colour
blindness, 64, **380**
congenital, 380
tests for, 381
sense, 64
vision,
Hering's theory, 63
Young-Helmholtz' theory, 63
Coloured vision, 379
Commotio retinæ, 404
Concave mirrors, 22
Concomitant strabismus, **517**
Concretions, 179
Congenital
abnormalities
of choroid, 354
of iris, 261
of lens, 305
of lids, 587
of nasal duct, 593
of optic disc, 369
of puncta, 593
of retina, 354
amblyopia, 521
anomalies of canaliculi, 593
colour blindness, **380**
defects of extrinsic muscles, 538
spastic paralysis, ocular manifestations of, 542
Congestion, conjunctival, 77
Conical cornea, 231
Conjugate
deviation, 494, 496
paralysis, 509
Conjunctiva, 1
bacteriology of, 139
congestion of, 77
cysts of, 184
degenerative changes in, 179
disease of, 139
examination of, 75
hyperæmia, 77, 140
inflammation of, 140
investigation of, 627
syphilis of, 176
therapeutics of, 629
tubercle of, 175
tumours of, 184
wounds, 405
Conjunctivitis, **140**
acute simple, 143
angular, 166
catarrhal, 143
chronic,
in cataract operation, 292
simple, 164
treatment of, 165
differential diagnosis from iritis, 246
diphtheritic, 156
diplobacillary, 166
due to drugs, 176
eczematous, 159
follicular, 167
gonorrhœal, 148
treatment of, 150
granular, 168
membranous, 156
treatment of, 158
metastatic gonorrhœal, 152
muco-purulent, 143
pathology of, 144
treatment of, 146
Parinaud's, 178
phlyctenular, 159
treatment of, 162
purulent, 148
secretion in, 143
Contusion
of eye, 397
of lids, 585
Convergence, 499
amplitude of, 500
and accommodation, **499**, 522
insufficiency of, 534
Converging power of eye, 45
Convex mirrors, 23
Copper foreign bodies, 411
Copper sulphate, 172
Corectopia, 261
Cornea,
anatomy of, 1
birth injury of, 234

Cornea—*contd.*
blood in, 235
blood-staining of, 398
degenerative changes in, 228
diseases of, 187
examination of, 80, 102
facets of, 191
fistula of, 194
foreign bodies in, 394
inflammation of, 187
in glaucoma, 233, 268
injuries of, 397
investigation of, 627
marginal atrophy of, 229
opacities of, 234
operations on, 431
perforation of, causing glaucoma, 264
perforation of ulcer of, 192
pigmentation of, 235
recurrent erosion of, 398
refractive index of, 32
rupture of, 399
sensibility of, 86
symptomatic conditions of, 232
therapeutics of, 631
tumours of, 235
ulcers of, 171, **186**
vessels in, 85, 86
wounds of, 405
Corneal loupe, 83
Corneo-sclera, wounds of, 406
Corpora quadrigemina, diseases of, 549
Corresponding points, 498
Couper's
capsule forceps, 203
lacrymal probes, 594
Credé's method, 154
Crescent,
inferior, 369
myopic, 474
Critchett's lens spoon, 448
Cryptophthalmia, 588
Crypts of iris, 4
Crystalline lens. *See* Lens.
Cup, glaucomatous, 269
Curette evacuation, 444
Cyclitis, **256**
chronic, 257
plastic, 260
purulent, 194
simple, 257
sympathetic, 426
treatment of, 259
Cycloplegia, 489
glaucomatous, 270
traumatic, 400
Cycloplegics, 57, 631
Cystotome, Moorfield's, 446
Cysts,
implantation, 184
of conjunctiva, 184
of iris, 262
of Meibomian, 568
of Moll's glands, 585
of orbit, 610
of retina, 354
orbito-palpebral, 588
tarsal, 568

Dacryo-adenitis, 591
Dacryo-cystitis, 203, 594
Dacryops, 591
Daturin, 58, 631
Day-blindness, 379
Degenerative changes
in choroid, 321
in conjunctiva, 179
in cornea, 228
in lens, 285
in vitreous, 307
in retinal vessels, 334
Dendritic ulcer, 217
Dermoid
cyst, 610
of conjunctiva, 185
Dermo-lipomata of conjunctiva, 185
Descemet's membrane, 1
rupture of, 234, 283
Desmarres'
forceps, 576
paracentesis needle, 199
retractor, 84
Deuteranopes, 380
de Wecker's scissors, 201
Diabetes,
cataract complicating, 303
mellitus, ocular complications of, 559
Diabetic
iritis, 252
retinitis, 343
Digestion, organs of, diseases of, ocular complications of, 559
Dilator
iridis, 4
Nettleship's canaliculus, 592
Dimmer's operation for ectropion, 578
Dionin, 182, 198, 630
Dioptre, 29
Diphtheria bacilli, 158
Diphtheritic conjunctivitis, 156
Diplobacillary
conjunctivitis, 166
ulcers, 206, 212
Diplobacilli, 166, 205, 212, 565

Diplopia, **504**
binocular, 505
chart, 508, 510
heteronymous, 505
homonymous, 505
uniocular, 402
Discission, 441
in lamellar cataract, 297
needles, 438
of secondary cataract, 443
Dislocation
of lacrymal gland, 591
of lens, 265, 288, **401**
Disparate points, 498
Distichiasis, 588
Donders
on convergent strabismus, 522
theory, 523
Drops, 630
Drugs
causing conjunctivitis, 176
toxic effects of, 349
use and action of, 57
Duboisin, 58, 631

Ecchymosis, subconjunctival, 181
"Eclipse blindness," 346
Ectasia
of cornea, 191, **229**
sclerotic, 273
Ectopia lentis, 305
Ectropion, 565, **575**
of uveal pigment, 273
Eczematous
conjunctivitis, 159
keratitis, 210
Electric light ophthalmia, 179
Electrolysis, 571
Elephantiasis neuromatodes, 588
Elliot's disc forceps, 439
Embolism of central artery, 330
Emmetropic eye, 33, 473
Emphysema of lids, 585
Empyema of frontal sinus, 608
Encephalitis lethargica, 555
Endophthalmitis, metastatic, 318
Enophthalmos, 602
Entropion, 571
cicatricial, 572
Epicanthus, 588
Epiphora, 565, 575, 591
Episcleritis, 236
Epithelioma, 186, 587
Ergot, 291
Erysipelas, 558, 563
Erythropsia, 379
Eserin, **58**, 274, 282
in corneal ulceration, 195
Esophoria, 530
Ethmoiditis, 609
Ethyl-hydrocuprein, 208, 349
Eucain, 630
Euphthalmin, 631
Eversion of punctum, 591
Evisceration of eyeball, 461
Examination,
external, 74
functional, 124
of anterior chamber, 86
of conjunctiva, 75
of cornea, 80
of iris, 87
of lacrymal apparatus, 79
of lens, 90
of pupils, 87
of sclerotic, 80
of tension, 91
ophthalmoscopic, 93
preliminary, 627
Excision
of eyeball, 459
of lacrymal sac, 598
scissors, 461
Exenteration of orbit, 613
Exophoria, 533
Exophthalmia, 602
Exophthalmic goitre, 614
Exophthalmometers, 603
Exophthalmos, 602
intermittent, 615
pulsating, 614
Expression in trachoma, 174
External rectus, paralysis of, 510
Extraction,
linear, 444
of senile cataract, 445
Eyeball,
excision of, 459
investigation of, 627
Eyelids, investigation of, 627

Facets of cornea, 191, 213
Family
idiocy, amaurotic, 333
maculo-cerebral degeneration, 333
retinitis pigmentosa, 323
Far point, 36
Fascicular ulcer of cornea, 210
Fibro-fatty tumours of conjunctiva, 185
Fibromata of conjunctiva, 185
Field of vision, 64
binocular, 137
in hemianopia, 371, 374
testing of, 133
uniocular, 64, 133
Filtering scar, 280

Fistula,
corneal, 194
lacrymal, 591
Fixation,
eccentric, 520
development of, 516, 521
"Fixation abscess," 430
Fluorescein, 207, 631
Focal
illumination, 81
interval of Sturm, 42
Follicular conjunctivitis, 167
Fontana, spaces of, 4
Forceps,
capsule, Couper's, 203
Desmarres', 576
disc, Elliot's, 439
entropion, 576
fixation, 197
Graddy's, 173
iris, 200
Knapp's, 174
Prince's, 535
ring, Griffith's, 571
T-shaped, 573
Wilde's, 576
Foreign bodies,
injuries caused by, 393
intraocular, 409
size and velocity of, 410
war injuries, 422
Form sense, 61
Förster's areolar choroiditis, 316
Fovea centralis, 8
Foveal reflex, 119
Friedreich's disease, 542
Frontal sinus, distension of, 608
Functional examination, 124
Fundus,
investigation of, 628
oculi, 117
ophthalmoscopic examination of, 119, 121
"Fusion centre," 524

Generative organs, disorders of, ocular complications of, 560
Glands,
Krause's, 561, 589
lacrymal, 589
Meibomian, 562
Moll's, 562
Zeis's, 562
Glare, 623
Glass rod, 147
Glass-workers' cataract, 305, 619
Glasses
for ametropia, 38
tinted, in cataract, 291
Glaucoma, 263, 619
absolute, 272
acute, 270
age and, 267
both eyes involved, 281
causes of secondary, 264
chronic, 276
congestive, 271
differential, diagnosis from iritis, 246
hæmorrhagic, 266
infantile, 283
iridectomy for, 275, 432
primary, 266
secondary, 263
after cataract extraction, 458
treatment of, 274
trephining in, 281
Glioma of retina, 389
Globe,
anatomy of, 1
injuries involving the, 585
rupture of, 399
Goitre, exophthalmic, 614
Gonococci, 149, 152, 158, 206
Gonorrhœal
conjunctivitis, 148
iritis, 251
Graddy's forceps, 173
Graefe
knife, 208, 446
sign, 614
Granular conjunctivitis, 168
Granulomata of conjunctiva, 184
Graves's disease, 614
Green-blind, 381
Griffith's ring forceps, 571
Gumma
of ciliary body, 260
sclerotic, 239
Gummatous iritis, 250

Haab's giant magnet, 417
Hæmangioma, 586
Hæmorrhage,
expulsive, after cataract extraction, 457
into the vitreous, 308
intraocular, 194
cause of glaucoma, 265
retinal, 336
subhyaloid, 337
Helmholtz'
ophthalmoscope, 94
theory of accommodation, 46
Hemiachromatopsia, 372
Hemianopia, 370
altitudinal, 374

Hemianopia—*contd.*
associated with cortical and subcortical lesions, 371
associated with lesions of optic tract, 372
binasal, 376
bitemporal, 373
fields of vision in, 371, 373
homonymous, 370
quadrant, 372
Hemianopic pupil reaction, 56, 88
Hemicrania, 378
Herbert's operation, 281
Hereditary ataxy, 542
Hering's theory of colour vision, 63
Herpes
corneæ, 218
ophthalmicus, 220
Hess's operation for ptosis, 583
Heterochromia iridis, 87, 261
Heterophoria, 530
Hexamin, 184
Hippus, 57
Hole at macula, 404
Holmgren's wools, 382
Homatropin, **58**, 290, 292, 631
Hordeolum, 567
internum, 568
Hutchinson's teeth, 225
Hyaloid
artery, persistent, 310
membrane, 7
Hydrocephalus, ocular manifestations of, 556
Hydrophthalmia, 282
Hygiene of vision, 621
Hyoscin, 58, 631
Hyperæmia
of conjunctiva, 140
treatment of, 142
optic disc, 357
retina, 335
Hypermetropia, 36, 49, 479
axial, 480
curvature, 480
examination in, 98
index, 480
latent, 128
manifest, 128
total, 128
Hypermetropic eye, 34, 36, 37
Hyperopia, 479
Hyperphoria, 531
Hypopyon, **189**, 206, 243
ulcer, 203
treatment of, 206

Illumination, 621
focal or oblique, 81
Index, refractive, 24
India, appointments in, requirements of candidates for, 635
Infections,
intraocular, 413
septic, after cataract extraction, 457
Infectious diseases, ocular manifestations of, 558
Inflammation
in sympathising eye, 427
of ciliary body, 241
of conjunctiva, 140
of glands of lids, 567
of iris, 241
of lids, 563
of orbit, 603
of sclerotic, 236
Influenza, ocular complications of, 558
Injuries
caused
by burns, 396
by caustics, 396
by foreign bodies, 393
of lids, 584
of orbit, 609
perforating, 405
to choroid, 403
to ciliary body, 400
to iris, 399
to lens, 401
to retina, 404
to the eye, 392, 585, 620
Insane, general paralysis of, 550
Instruments, care of, 633
Intracranial
abscess, 553
disease, papillitis due to, 361
tumours, 552
Intraocular
hæmorrhage, 194
cause of glaucoma, 265
pressure, 14, 91, 263, 269
tumours, 385
Iridectomy, **431**
broad, 280
for glaucoma, 275, 432
for iritis, 255
for prolapsed iris, 201
optical, 431
peripheral, 456
preliminary, 294
Irideremia, 261
traumatic, 400
Iridocyclitis,
plastic, 426
sympathetic, treatment of, 429
See also Cyclitis.

Iridodialysis, 105, 400
Iridodonesis, 87, 283, 288, 401
Iridotomy, 437
Iris, 4
  abnormalities of, 261
  anatomy of, 4
  bombé, 248
  cysts of, 262
  diseases of, 241
  examination of, 87
  forceps, 200
  inflammation of, 241
  injuries to, 399
  melanoma of, 261
  movements of, 52
  operations on, 431
  prolapse of, 192, 202, 456
    iridectomy of, 201
  repositor, 202
  tremulous, 87, 283, 288, 401
  tumours of, 385
Iritis, **241**
  after cataract extraction, 458
  diabetic, 252
  differential diagnosis from conjunctivitis, 246
  gonorrhœal, 251
  gummatous, 250
  plastic, 243
  rheumatic, 251
  serous, 257, 427
  symptoms of, 246
  syphilitic, 249
  treatment of, 253
  tuberculous, 252
  varieties of, 249
Iron foreign bodies, 411

Jaeger test-types, 131
Jaesche-Arlt operation, 574

Keratalgia, traumatic, recurrent, 398
Keratectasia, 231
  from ulcer, 191
Keratic precipitates ("K.p."), 224, 234, 243, 257, 426
Keratitis, **187**
  bullous, 233, 273
  disciformis, 209
  e lagophthalmo, 214, 581
  filamentary, 234, 273
  interstitial, 222
  neuroparalytic, 214
  non-suppurative, 218
  parenchymatous, 222
  phlyctenular, 210, 618
    treatment of, 211
  profunda, 227
Keratitis—*contd.*
  punctata ("K.p."), 224, 234, 243, 257, 426
  purulent, 187
  rare forms of, 227
  sclerosing, 234, 238
  superficial punctate, 221
  vesicular, 233, 273
Keratocele, 192
Keratoconus, 231
Keratoglobus, 232, 283
Keratomalacia, 213
Keratome, 199
Kinetic strabismus, 512
Knapp's roller forceps, 174
Knife,
  cataract, Graefe's, 208, 446
  Beer's, 569
  Graefe, 208
  Lang's, 439
  Stilling's canaliculus, 593
  Tweedy's canaliculus, 593
  Weber's, 592
  Ziegler's, 442
Koch-Weeks' bacilli, 145
"K.p.," 224, 234, 243, 257, 426
Krause's glands, 561, 589
Krönlein's operation, 613

Lacrymal
  abscess, 601
  apparatus,
    anatomy of, 589
    diseases of, 591
    examination of, 79
    physiology of, 590
  ducts, 589
  glands, 589
    diseases of, 591
    dislocation of, 591
    fistula of, 591
    tumours of, 591
  passages, diseases of, 591
  probes, Couper's, 594
  sac, 589
    excision of, 598
    inflammation of, 594
  secretion, 590
Lagophthalmia, 214, **581**
Lagrange's operation, 281
Lamellar cataract, 295
Lamina
  cribrosa, 9, 118, 269, 366
  plane, 24
  vitrea, 6
Lamp, Gullstrand's slit, 83
Lang's
  dissector and scoop, 597
  knife, 439

Lang's speculum, 196
Lantern test for colour blindness, 381
Lawford's speculum, 433
Lead
deposits, 235
poisoning, 349
Leber's disease, 365
Legibility, 625
Lens,
anatomy of, 9
coloboma of, 305
congenital, abnormalities of, 305
curvature of, 46
diseases of, 285
See also Cataract.
dislocation of, 193, 265, 288, 305, 401
cause of glaucoma, 265
examination of, 90, 104, 106, 285, 628
foreign bodies in, 417
injuries to, 401, 406, 408
intumescence of, 287
opacities of, 90, 286
operations on, 441
reflexes, 484
spoon, Critchett's, 448
wounds of, 406, 408
cause of glaucoma, 265
Lenses, 25
correcting, effect of, on retinal image, 38
cylindrical, 43
measurement of, 29
refractive power of, 29, 47
types of, 25–31
Lenticonus, 305
Leucæmia, retinitis in, 344, 559
Leucoma, 190
adherent, 205
Levator palpebræ, 563, 581
Lid
spatula, 574
upper, eversion of, 75
Lids,
anatomy of, 561
anomalies of position of, 570, 588
carcinoma of, 587
coloboma of, 588
congenital abnormalities of, 588
contusions of, 585
diseases of, 561
glands of, inflammation of, 567
inflammation of, 563
Lids—*contd.*
injuries of, 584
œdema of, solid, 564
sarcoma of, 587
syphilis of, 566
therapeutics of, 629
tumours of, 585
wounds of, 584
Ligamentum pectinatum iridis, 4
Light,
bright, retinitis from, 346
difference, 61
minimum, 60
projection of, detection of, 290
reaction to, 56, 88
sense, 60, 138
transmission of, 20
Lime burn, 396
Line
of fixation, 518
of vision, 64, 518
Lithiasis, 179
Little's disease, 542
Liver, diseases of, 559
Lotions, 629
Loupe, corneal, 82
Luer's syringe, 595
Lymph,
of eye, 3
secretion of, 15
Lymphangioma
of conjunctiva, 184
of lids, 586

McHardy's perimeter, 134
Macropsia, 315
Macula lutea, 8, 119, 122
cortical representation of, 69
of cornea, 190
Maculo-cerebral family degeneration, 333
Madarosis, 564
Maddox rod, 532
Magnet,
Haab's, 417
hand, 416
Mellinger's ring, 420
Malaria, 217
Malingering, 383
Manometer, diagram of, 15
Margo-intermarginalis, 561
Maunoir's angular iris scissors, 447
Maxwell's operation, 463
Measles, 558
Meibomian glands, 562
inflammation of, 568
Melanoma of iris, 261

Melliner's ring magnet, 420
Membrane of Bruch, 6
Membranous conjunctivitis, 156
Meningitis,
epidemic cerebro-spinal, 555
purulent, 555
tuberculous, 554
Meningococcus, 149, 320
Meningo-encephalocele, 610
Mercantile marine, sight tests for masters and mates, 638
Metabolic diseases, ocular complications of, 559
Metallic foreign bodies, 411
Metamorphopsia, 315, 328
Metre angle, 499
Microblepharon, 588
Micrococcus catarrhalis, 149
Microphthalmia, 311, 354, 588
Micropsia, 315, 328
Migraine, 330, **378**
ophthalmoplegic, 379
Mikulicz's disease, 559
Miners' nystagmus, 516
Miosis, 59, 544
traumatic, 400
Miotics, 58, 632
Mirror, preliminary examination with, 97, 101
Mirrors,
concave, 22
convex, 23
plane, 21
Möbius' sign, 614
Moles, congenital,
of conjunctiva, 185
of lid, 586
Moll's glands, 562
cysts, of, 585
Molluscum contagiosum, 586
Mooren's ulcer, 247
Moorfield's
cataract bandage, 450
cystotome, 446
strabismus hook, 460
Morax-Axenfeld diplobacillus, 166, 205, 212, 565
Morgagnian
cataract, 288
globules, 285
Mucocele, 594
contraindication to cataract operation, 293
Muco-purulent conjunctivitis, 143
Mules's operation, 461
Müller, fibres of, 7
Müller's
muscles, 563
retractor, 596, 599
Muscæ volitantes, 289, 306, 475
Muscles,
extrinsic,
anatomy and physiology of, 492
congenital defects of, 538
intrinsic, 5
Müller's, 563
operations on, 534
paralysis of, 502, 541, 544
Myasthenia gravis, 545
Mydriasis, traumatic, 400
Mydriatics, 57, 631
Myelitis, 542
Myopia, 36, **472**, 618
axial, 472
curvature, 472
examination in, 99
high, 478, 618
operative treatment of, 479
index, 472
over-correction in, 478
spherical, 469
Myopic eye, 35, 36, 38
Myotonia atrophica, 546

Nævi
of conjunctiva, 185
of lid, 586
Nagel's anomaloscope, 382
Nasal duct, 590
stricture of, 601
Navy, Royal, requirements of candidates for commissions, 635
Near point, 46, 132
Nebula, 190
Necrosis, corneal, 188, 200
Needle holder, Silcock's, 536
Needles,
broad, 198
curved, 536
discission, 438
paracentesis, Desmarres', 199
Needling, 441
Nephritis, albuminuric retinitis in, **341**, 377
Nerve, optic,
anatomy of, 8
diseases of, 357
tumours of, 611
Nerves, motor, 495
of iris, 55
Nervous system, diseases of, 540
Nettleship's canaliculus dilator, 592
Neuritis,
optic, 329, 340, **357**, 365, 553, 555, 612
retrobulbar, 363, 543
Neurology of vision, 65

Neuroma, plexiform, 588
Neurons, orders of, 67
Neuroparalytic keratitis, 214
Neuro-retinitis, 329, **340**, 361
Night-blindness, 213, 327, **379**
Nodal point, 34, 518
Nose, accessory sinuses of, distension of, 608
Novocain, 629
Nystagmoid jerks, 514
Nystagmus, **514**
  labyrinthine, 515
  miners', 516

Oblique illumination, 81
Occlusio pupillæ, 245
Occlusion
  of canaliculus, 593
  of punctum, 593
Occupations, eye tests for various, 634
Ocular paralysis, varieties of, 509
Œdema of lids, 563
Ointments, 631
Opacities,
  congenital, of cornea, 234
  diagnosis of, 102
  in cataract, 291, 295
  of cornea, 196, 229, 234, 398
  of lens, 285
  of vitreous, 306
  striate, of cornea, 234
Opaque nerve fibres, 356
Operations
  on extrinsic muscles, 534
  on the eyeball, 431
Ophthalmia,
  catarrhal, 143
  due to electric light, 179
  gonorrhœal, 148
  neonatorum, 152, 618
    prevention of, 155
  nodosa, 178
  phlyctenular, 159
  sympathetic, 393, 425
Ophthalmology, preventive, 617
Ophthalmoplegia, 509, 546
Ophthalmoscope, 94
Ophthalmoscopic examination, 93
  by direct method, 111
  by distant direct method, 101
  by indirect method, 106
  preliminary, 97
Optic
  atrophy, 365
    post-neuritic, 366
    primary, 365
  axis, 33, 518
  disc, 117
Optic—*contd.*
  disc—*contd.*
    coloboma of, 369
    hyperæmia of, 357
  nerve,
    anatomy of, 8
    diseases of, 357
    extradural tumour of, 611
    intradural tumour of, 611
  neuritis, 329, 340, **357**, 365, 553, 555, 612
  tract, lesions of, 372
Optical
  centre, 27, 34
  iridectomy, 431
Optics, 19
  physiological, 32
Optochin, 208, 349
Ora serrata, 6
Oral sepsis, ocular complications of, 559
Orbicularis palpebrarum, 561, 563
Orbit,
  bones of, injuries to, 584
  cellulitis of, 604
  diseases of, 602
  exenteration of, 613
  inflammation of, 603
  injuries of, 584, 609
  tumours of, 610
Orientation, 497
  false, 505
Orthophoria, 530
Orthoptic treatment, 527
Osteoma, orbital, 610
Oxycephaly, 554

Pannus,
  phlyctenular, 211
  trachomatous, 170
Panophthalmitis, 194, **319**, 393, **423**, 564, 605
  metastatic, 319
Papillitis, 341, **357**, 359, 543, 551–557, 612
Papillœdema, 359, 549, 551–557
Papillo-macular fibres, **69**, 348, 364
Papilloma of conjunctiva, 184
Paracentesis, **199**, 207, 430
Parallactic displacement, 102, 111, 115
Paralysis,
  conjugate, 509
  ocular, 507, 542
    varieties of, 509
  of accommodation, 489
  of external rectus, 508, 510
  of insane, general, 550
  of superior oblique, 510

Paralysis—*contd.*
of third nerve, 510
progressive, 550
spastic congenital, 542
Paralytic dementia, 550
Parinaud's conjunctivitis, 178
Parotitis, 559
Patient, preliminary investigation of, 627
Pemphigus, 179, 183
Perimeter, 133
chart, 136
McHardy's, 134
Periostitis, 603
Peritomy, 174
Phlyctenular
conjunctivitis, 159
keratitis, 210, 618
Photophobia, 162, 564
Photophthalmia, 179
Photopic vision, 63
Photopsiæ, 315, 350
Photo-retinitis, 346
Phthiriasis palpebrarum, 565
Phthisis
bulbi, 256
ocular complications of, 558
Physiological
cup, 118
optics, 32
Physiology
of extrinsic ocular muscles, 492
of eye, 14
of lacrymal apparatus, 589
Physostigmin, **58**, 282, 632
Pigment, retinal, 7, 313, 324
Pigmentation
of cornea, 235
of retina, 323, 356
Pigments, 630
Pilocarpin, **58**, 282, 632
Pilot service, requirements of candidates for, 638
Pineal glands, diseases of, 549
Pinguecula, 179
"Pink eye," 144
Plane
lamina, 24
mirrors, 21
Plastic iritis, 243, 426
Plexiform neuroma, 588
Pneumococci, 145, 158, 203, 594
Pneumococcic ulceration, 204, 208
Pneumonia, 558
Points,
corresponding, 498
disparate, 498
near and far, 46
nodal, 34, 518
Polioencephalitis, 554
Polycoria, 261
Polyopia, uniocular, 289
Pons, diseases of, 547
Porus opticus, 9
Posterior
chamber, 10
longitudinal bundle, 496
Precipitates, keratic ("K.p."), 224, 234, 243, 257, 426
Pre-retinal hæmorrhage, 336
Presbyopia, 50, **489**
Pressure,
intracapillary, 15
intraocular, 14
Priestley Smith's
tape, 525
theory of glaucoma, 266
Prince's forceps, 535
Prisms, 24
use of, 499, 533
Probes, lacrymal, 594
Projection
of light, 290
of objects, 497
false, 506
Prolapse
of ciliary body, 407
of iris, 201, 405, 456
of vitreous, 407, 452
Proptosis, 602
Protanopes, 380
Protargol, 148, 630
in blepharitis, 566
Pseudoglioma, 309, 391
Pseudo-neuritis, 359
Pseudo-pterygium, 180
Pterygium, 180
Ptosis, 545, **581**, 587
trachomatous, 171
Public services, requirements for candidates, 634
Pulsation of retinal vessels, 120, 269
Punch forceps, 441
Puncta lacrymalia, 589
Punctum,
congenital anomalies of, 593
displacement of, 79
lower, eversion of, 591
occlusion of, 593
proximum, 46, 132
remotum, 36
Pupillary membrane,
inflammatory, 245, 249
persistent, 261
Pupils, 52
contraction of, 51, 53
dilatation of, 89
examination of, 87

Pupils—*contd.*
innervation of, 53
investigation of, 628
reactions, 56, 57, 87, 132
Pus in the vitreous, 309

Quinine
amblyopia, 348
lotions, 159, 630

Radium, 587
Railway employees, eye tests for, 636, 638
Range
of accommodation, 49, 132
of convergence, 500
Reaction to light, 56
Reactions of pupils, 56, 57, 132
Reading and writing, 624
Recruits, eye tests for, 634
Red-blind, 380
Reflection, 21
Reflex, foveal, 119
Reflexes,
during ophthalmoscopic examination, 107
pupillary, 56
Refraction, 23
dynamic and static, 50
errors of, 35, 464, 472
correction of, 486
of divergent rays, 44
of parallel rays, 43
static, 44
Refractive
index, 24
of cornea, 32
power of lenses, 47
Remote or far point, 37
Repositor, iris, 202
Respiratory tract, diseases of, 558
Retina, 6
adaptation of, 60
anatomy of, 6
blood-supply of, 11
coloboma of, 354
congenital pigmentation of, 356
cysts of, 354
detachment of, **350**
recognition of, 104
diseases of, **312**, 328
examination of, 104
fibres of, 69
glioma of, 389
hole in, 404
injuries to, 404
operations on detached, 353
primary affections of, 328
vascular disorders of, 329
Retina—*contd.*
vessels of, 11, 118
Retinal vessels, perivascular changes in, 329
Retinitis, **328**
albuminuric, 340
arteriosclerotic, 335
circinata, 345
diabetic, 343
exudative, 346
from bright light, 346
hæmorrhagic, 344
leucæmic, 344
pigmentosa, 323
sine pigmento, 327
primary, 328
proliferans, 308, 345
punctata albescens, 327
purulent, 346, **423**
septic, 344
syphilitic, 339
Retinoscopy, 464
Retractor,
Desmarres', 84
Müller's, 596
wire, 84
Retrobulbar neuritis, 363, 543
Rheumatic iritis, 251
Rheumatism, ocular complications of, 559
Rodent ulcer of cornea, 217
Ronne's step, 278
Rupture
of choroid, 403
of cornea, 398
of iris, 400
of retina, 404
of sclerotic, 399

Saemisch's section, 207
Salmon patches, 222
Salvarsan in treatment, 254, 430, 512
Sarcoma
of choroid, 385
of ciliary body, 388
of conjunctiva, 186
of iris, 385
of lids, 586
of orbit, 611
Saunders's discission needle, 438
Scaphocephaly, 554
Scarlet fever, 558
Schiötz's tonometer, 92
Schlemm, canal of, 3, 13, 18, 283
Scissors,
de Wecker's, 201
excision, 461
Maunoir's, 447
tenotomy, 459

Scleral
puncture, 353
ring, 117
Sclerectomy, posterior, 441
Scleritis, 237
annular, 239
Sclerosing keratitis, 234, 238
Sclerosis,
combined, 542
disseminated, 543
Sclerotic, 1
blue, 240
diseases of, **236**
examination of, 80
gumma of, 239
inflammation of, 236
injuries of, 399
operations on, 437
tubercle of, 239
ulceration of, 239
wounds of, 408
Sclerotomy,
anterior, 284, 439, 441
posterior, 275, 441
Scopolamin, 631
Scotoma,
absolute, 135, 330
central relative, 137
negative, 315
positive, 315
ring, 325
scintillating, 378
sector-shaped, 331
Scotopic vision, 61, 63
Screen test in strabismus, 519
Seclusio pupillæ, 245
Secretion, lacrymal, 590
Seidel's sign, 278
Septic infection after cataract extraction, 457
Serpiginous ulcer, 204, 212
chronic, 212, 217
Serum treatment, 632
Shadow test, 464
Short sight, 40, 472
Siderosis bulbi, 411
Sight tests for candidates for public services, 634
Silcock's needle holder, 536
Silver nitrate,
application of, 147, 151, 158, 165, 172, 196
salts, 148
wire arteries, 331
Skiascopy, 464
Skull, base of, fractures of, 556
Small-pox, 558
Snellen's
" reform " eye, 462
Snellen's—*contd.*
sutures, 578
test types, 125, 129
Snow blindness, 179
Socket, contracted, 462
Spasm of accommodation, 128, 490
Spasmus nutans, 514
Spatula, 199
lid, 574
Spectacles, 38, 488
Spectrum, formation of, 19
Speculum,
Lang's, 196
Lawford's, 433
Spherical aberration, 52
Sphincter iridis, 4
Spoon,
Critchett's, 448
sharp, 570
Spring catarrh, 177
Spud, 395
Squint. *See* Strabismus.
Staphylococcus, 160, 206, 457, 568, 594
Staphyloma,
anterior, 193, 229
ciliary, 238, 273
equatorial, 273
intercalary, 273
posterior, 474
posticum verum, 474
Statistics of eye symptoms in disease of nervous system, 557
Steel particles in the eye, 395
Stellwag's sign, 614
Stilling's
canaliculus knife, 593
tests for colour blindness, 382
Stone particles in the eye, 395
Strabismus, **502**
alternating, 519
apparent, 518
concomitant, 517
treatment of, 526
convergent, 522
divergent, 523, 529
hook, 460
kinetic, 517
latent, 530
paralytic, 502
periodic, 520, 531
unilateral, 519
Streptococcus, 157, 158, 206, 594
Striate opacity, 234
Stricture of nasal duct, 601
Sturm's
conoid, 42
focal interval, 42

Stye, 567
Styles, lacrymal, 595
Subconjunctival
ecchymosis, 181
injections, 353
Subhyaloid hæmorrhage, 336
Sulcus subtarsalis, 139, 561
Superior oblique, paralysis of, 510
Suprarenin, 598, 632
Suspensory ligament, 46
rupture of, 402
Symblepharon, 579
prevention of, 397
Sympathetic
ophthalmia, 393, 425
paralysis of cervical, 58
Symptomatic
conditions of cornea, 232
diseases of eye, **540**
disturbances of vision, **370**
Synchisis, 307
scintillans, 307
Synechia,
annular, 245, 265
posterior, cause of glaucoma, 265
anterior, 192
cause of glaucoma, 265
posterior, 246
ring, 245
Synkinesis, 51, 493, 513
Syphilis,
cerebral, ocular manifestations of, 551
of conjunctiva, 176
of cornea, 222
of lids, 566
Syphilitic
affections, treatment of, 632
choroiditis, 315
iritis, 249
retinitis, 339
tarsitis, 566
Syringe, lacrymal, Luer's, 595
Syringomyelia, 545

Tabes dorsalis, 541
Tarsal cyst, 568
Tarsitis, syphilitic, 566
Tarsorrhaphy, 216
Tattooing, 198
Tay's choroiditis, 321
Tears, 591
Teeth, hypoplasia of, lamellar cataract with, 296
Teichopsia, 378
Telangiectases of lids, 586
Tenonitis, 607
Tenotomy, 512, **534**
scissors, 459
Tension,
intraocular, 14, 91, **263**, 628
testing of, 91
Test types,
Bjerrum's, 138
for far vision, 125
for near vision, 130, 131
illumination of, 127
Jaeger's, 130
Snellen's, 125, 130
Tests
for candidates for public services, 634
colour blindness, 381
Therapeutic notes, 629
Thrombosis
of cavernous sinus, 605
of central vein, 266, **337**, 338
Tiersch graft, 575
Tigroid fundus, 120
Tobacco amblyopia, 347
Tonometer, Schiötz's, 92
Torticollis, ocular, 506
Torus, 40
Toxic amblyopia, 347
Trachoma, **168**
complications of, 170
treatment of, 172
Transillumination, 80, 388
Treatment
by serums and vaccines, 632
of blepharitis, 565
of cicatricial entropion, 574
of complicated corneal ulcers, 198
of concomitant strabismus, 526
of detachment of retina, 353
of dislocation of lens, 402
of ectropion, 577
of follicular conjunctivitis, 167
of foreign bodies in the eye, 394, 416
of glaucoma, 274, 280, 284
of gonorrhœal conjunctivitis, 150
of hyperæmia of conjunctiva, 142
of hypopyon ulcer, 206
of injuries caused by caustics, 397
of irido-cyclitis, 259
of iritis, 253
of membranous conjunctivitis, 158
of muco-purulent conjunctivitis, 146
of ophthalmia neonatorum, 154

Treatment—*contd.*
of phlyctenular conjunctivitis, 162
of phlyctenular keratitis, 211
of ptosis, 582
of simple chronic conjunctivitis, 165
of spastic entropion, 572
of symblepharon, 580
of sympathetic ophthalmia, 428
of syphilitic infections, 632
of trachoma, 172
of uncomplicated corneal ulcers, 194
Tremulous iris, 87, 283, 288, 401
Trephine, 439
Trephining, 439
for chronic glaucoma, 281
Trichiasis, 565, **570**
T-shaped forceps, 573
Tubercle
of choroid, 316
of conjunctiva, 175
of iris, 252
of lacrymal gland, 591
of sclerotic, 239
Tuberculous iritis, 252
Tumours,
intracranial, 547, 552
intraocular, **385**
cause of glaucoma, 265
of auditory nerve, 548
of choroid, 385
of ciliary body, 388
of conjunctiva, 184
of cornea, 235
of iris, 385
of lacrymal gland, 591
of lids, 585
of optic nerve, 611
of orbit, 610
of pons, 547
of retina, 389
Tweedy's canaliculus knife, 593
Tylosis, 565
Typhoid fever, 558

Ulceration of sclerotic, 239
Ulcers,
atheromatous, 214
central, 213
chronic serpiginous, 217
complicated, treatment of, 198
dendritic, 217
fascicular, 210
herpetic, 218
hypopyon, 203
marginal, 212
Mooren's, 217
Ulcers—*contd.*
mycotic, 210
of cornea, 171, **187**
perforating, 192
ring, 212
rodent, 217
trachomatous, 171
treatment of, 194
Ulcus serpens, 204, 212, 217
Ultra-violet rays, 19, 179, 346
Undine, 146
Uræmic amaurosis, 342, 376
Urotropin, 184
Uveal tract, 3, 4
blood supply of, 11, 12
Uveitis, 241

Vaccine treatment, 632
Vaccinia, 566
of eyelids, 558
Vascularisation in corneal ulcer, 188, 191
Vaso-constrictors, 632
Veins,
ciliary, 11
of eye, 10
ophthalmoscopic appearance of, 120
retinal, degenerative changes in, 334
vorticose, 13
Vernal catarrh, 177
Vertigo, 507
Vision,
acuity of, 61, 125, 289, 294
binocular, 498
colour, 68, 380
coloured, 379
field of, 64, 133
binocular, 137
in glaucoma, 276, 277, 278, 281
line of, 64
neurology of, 65
photopic, 63
scotopic, 61, 63
symptomatic disturbances of, 370
testing of, 133
Visual
acuity, 61, **125**
angle, 34
minimum, 62
centre, 68
perceptions, 59
colour sense, 63, 380
form sense, 61
light sense, 60

Vitreous,
blood in, 308
blood vessels in, 309, 339
diseases of, **306**
fibrous tissue in, 308
fluidity of, 307
foreign bodies in, 414
humour, 10
opacities of, **306**
parasites in, 311
prolapse of, 408, 452, 453
pus in, 309
Vossius' ring, 403
V-Y operation, 579

War injuries of the eye, 421
Weber's law, 61
Werner's mnemonic for diplopia, 509
Wharton-Jones operation, 579
Whooping cough, 558
Wilde's forceps, 576
Wire retractors, 84
Wolfe graft, 463
Word-blindness, 383
Wounds
of conjunctiva, 405
of cornea, 405
of lens, 408
of lids, 584
of sclerotic, 408
perforating, 405
with retention of foreign bodies, 409

Xanthelasma, 585
Xerophthalmia, 183
Xerosis, 183

Young-Helmholtz' theory of colour vision, 64

Zeis's glands, 561, 562, 567
Ziegler's knife, 442
Zinc salts, 165, 166, 212, 630
Zinn, zonule of, 9
Zonular
cataract, 295
opacity of cornea, 229

PRINTED IN GREAT BRITAIN BY THE WHITEFRIARS PRESS, LTD.,
LONDON AND TONBRIDGE.

Malay Poisons and Charm Cures.

By JOHN D. GIMLETTE, M.R.C.S., L.R.C.P., Residency Surgeon of Kelantan, one of the Un-Federated Malay States. With foreword by Sir W. H. WILLCOX, K.C.I.E., C.B., C.M.G., M.D., F.R.C.P. New (2nd) Edition. 8s. 6d. net. Postage 6d.

Eden's Manual of Midwifery.

5th Edition. 5 Plates (4 in Colour), 269 Text-figures. 24s. net. Postage 1s.

"*We have nothing but praise for this book.*"—THE LANCET.

Wood Jones' Principles of Anatomy as Seen in the Hand.

With 2 Plates and 123 Text-figures. 15s. net. Postage 7d.

Whittaker's Surface Anatomy.

3rd Edition. 17 Plates. 7s. 6d. net. Postage 6d.

Williams' Minor Surgery and Bandaging.

17th Edition. 239 Illustrations. 10s. 6d. net. Postage 6d.

"*One of the best of its kind . . . . . contains a wealth of information.*"—BRITISH JOURNAL OF SURGERY.

Taylor's Operative Surgery.

300 Illustrations. 32s. net. Postage 1s. 3d.

Taylor's Principles and Practice of Medical Jurisprudence.

Edited by FRED. J. SMITH, M.D., F.R.C.P. 7th Edition. With 20 Engravings. Two Vols. £3 3s. net. Postage 1s. 3d.

"*For the Lawyer as well as the Medical Practitioner this standard work is invaluable. In every branch of law into which medical considerations enter it may always be consulted with profit.*"—THE JUSTICE OF THE PEACE.

Arvedson's Notes on Diseases treated by Medical Gymnastics and Massage.

Translated by MINA L. DOBBIE, M.D. 8s. 6d. net. Postage 4d.

Kleen's Massage and Medical Gymnastics.

Translated by MINA L. DOBBIE, M.D., B.Ch., Med. Officer, Chelsea Phys. Training Coll. 2nd Edition. 182 Illus. 32s. net. Postage 1s.

"*The book will be found a most valuable contribution to physico-therapeutic literature. It is admirably produced.*"—THE LANCET.

Mennell's Massage : Its Principles and Practice.

2nd Edition. 167 Illustrations. 21s. net. Postage 1s.

Burns' Introduction to Biophysics.

By DAVID BURNS, M.A., D.Sc., Professor of Physiology, University of Durham. With a foreword by Prof. D. NOEL PATON, LL.D., F.R.S. 80 Illustrations. 21s. net. Postage 1s.

J. & A. CHURCHILL

Anrep and Harris' Practical Physiology.

Introduction by Prof. E. H. STARLING, C.M.G., F.R.S. With 197 Illustrations. 10s. 6d. net. Postage 6d.

Taylor and Poulton's Practice of Medicine.

New (12th) Edition. 30s. net. Postage 1s.

"*Wisely and extremely thoroughly edited.*"—BRITISH MEDICAL JOURNAL.

"*Dr. E. P. Poulton and his collaborators are to be congratulated on their revision of this famous text-book. Without disturbing the essential nature of the book, Dr. Poulton has managed to re-arrange its contents and bring it into harmony with the latest conceptions of pathology. The section dealing with diseases of the nervous system has now been placed towards the end of the volume, a new section has been introduced dealing with diseases due to dietetic deficiencies, and diabetes mellitus is placed in its true position—with other diseases of the endocrine organs . . . . . New articles are devoted to œrophagy, coeliac disease, and diverticulitis, and those dealing with achylia gastrica and the X Ray examination of the stomach have been largely revised. Dr. Poulton is particularly happy in his articles on diseases of metabolism, those concerned with basal metabolism and diabetes being specially learned; his description of ketosis marks a great advance in our knowledge . . . . . we feel sure that for many years to come the revised 'Taylor's Medicine' will enjoy the popularity it deserves.*"—THE LANCET.

Greenish's Microscopical Examination of Foods and Drugs.

3rd Edition. 200 Illustrations. 18s. net. Postage 9d.

Clayton's Theory of Emulsions and Emulsification.

Foreword by Prof. F. G. DONNAN, C.B.E., D.Sc., F.R.S. 22 Illustrations. 9s. 6d. net. Postage 6d.

Berkeley and Bonney's Difficulties and Emergencies of Obstetric Practice.

3rd Edition. 309 Illustrations. 36s. net. Postage 1s.

Cushny's Pharmacology.

7th Edition. 71 Illustrations. 24s. net. Postage 1s.

Sequeira's Diseases of the Skin.

3rd Edition. 52 Coloured Plates. 257 Text-figures. 38s. net. Postage 1s. 3d.

Lee's Microtomists' Vade-Mecum.

8th Edition. Revised and Edited by Dr. J. BRONTË GATENBY. 28s. net. Postage 1s.

Marshall's Microbiology for Agricultural and Domestic Science Students.

3rd Edition. 21s. net. Postage 1s.

"*We know of no other book which in so limited a space gives such an excellent account, general and special, of micro-organisms in all their aspects. The text contains numerous illustrations.*"—NATURE.

J. & A. CHURCHILL

Coles' Critical Microscopy.

How to get the best out of the Microscope. 8 Illustrations. 7s. 6d. net. Postage 5d.

Eden and Lockyer's Gynæcology.

2nd Edition. 24 Coloured Plates and 513 Text-figures. 36s. net. Postage 1s. 3d.

"*In every way the new edition maintains the high standard of its predecessor.*" —BRITISH MEDICAL JOURNAL.

Sterility in Woman, Its Causes and Treatment.

By R. A. GIBBONS, M.D., F.R.C.S.Edin. 44 Illustrations. 12s. 6d. net. Postage 6d.

Fleming's Medicine.

3rd Edition. 64 Illustrations. 21s. net. Postage 1s.

Goodhart and Still's Diseases of Children.

11th Edition. With 60 Illustrations. 32s. net. Postage 1s.

Hale-White's Materia Medica.

17th Edition. 10s. 6d. net. Postage 5d.

Gask and Wilson's Surgery.

A Text-book by Various Authors. With 30 Plates, 20 in colour, and 467 Text-figures. 30s. net. Postage 1s. 3d.

"*The book is good, not too big, well printed and illustrated, and worthy of the school from which it comes.*"—THE BRITISH MEDICAL JOURNAL.

Jellett's Practice of Gynæcology.

4th Edition. 374 Illustrations. 11 Coloured Plates. 21s. net. Postage 1s.

Jellett's Short Practice of Midwifery.

8th Edition. 207 Text-figures. 4 Coloured Plates. 18s. net. Postage 8d.

Hewlett's Bacteriology.

7th Edition. 31 Plates and 63 Text-figures. 21s. net. Postage 1s.

Hewlett's Pathology, General and Special.

5th Edition. 37 Plates and 15 Text-figures. 18s. net. Postage 1s.

Hewlett and Nankivell's Preventive Medicine.

17 Illustrations. 18s. net. Postage 1s.

Applied Pharmacology.

By A. J. CLARK, M.C., M.D., F.R.C.P., Professor of Pharmacology in the University of London. 46 Illustrations. 15s. net. Postage 1s.

Collis and Greenwood's Health of the Industrial Worker.

With Special Chapter on Reclamation of the Disabled. Introduction by Sir GEORGE NEWMAN, K.C.B., D.C.L. 38 Illustrations. 30s. net. Postage 1s.

"*This is the first adequate modern treatise upon the hygiene of industry in general. A more ideal combination of authors for this purpose it would be difficult to find. The book reaches an exceptionally high standard.*"—C. S. MYERS, F.R.S., *in* NATURE, 19th May, 1921.

Starling's Physiology.

3rd Edition. With 579 Illustrations. 25s. net. Postage 1s.

"*A monument to the masterly presentment of the subject by one of the most eminent of British Physiologists.*"—PHYSIOLOGICAL ABSTRACTS.

Bowlby and Andrewes' Surgical Pathology.

7th Edition. With 200 New Illustrations. 30s. net. Postage 1s.

Jex-Blake's Physical Signs in the Chest and Abdomen.

27 Illustrations. 9s. 6d. net. Postage 4d.

Craig's Psychological Medicine.

3rd Edition. 27 Plates. 18s. net. Postage 9d.

Craig's Nerve Exhaustion.

6s. net. Postage 4d.

Pauli's Colloid Chemistry of the Proteins.

Translated by P. C. L. THORNE, M.A., A.I.C. 8s. 6d. net. Postage 4d.

Crowther's Principles of Radiography.

55 Illustrations. 7s. 6d. net. Postage 4d.

Moore's Medical Ophthalmology.

80 Illustrations. 15s. net. Postage 8d.

"*The whole book is excellent . . . . . One of the most useful to the Physician and to the Ophthalmic Surgeon alike that has appeared in this country in recent years.*"—BRITISH MEDICAL JOURNAL.

Rowlands and Turner's Operations of Surgery.

797 Illustrations, 40 in colour. 2 volumes. £2 10s. net. Postage 1s. 6d.

"*The very best book of its kind in the British language.*"—THE PRACTITIONER.

J. & A. CHURCHILL

USE A SEPARATE SLIP FOR EACH WORK WANTED

Read the Rules on the other side

BOSTON MEDICAL LIBRARY

AUTHOR	TITLE Brief Title. Give place and date If a periodical, give year, vol. and page	SHELF NO.

Date *Name of reader*

Address

FORM 3. 9-38-10M-EDA